

AMATERASU: a tool to track stellar activity

Telmo Monteiro^{1,2*}¶, João Gomes da Silva^{2*}, Elisa Delgado-Mena^{3*}, and Nuno C. Santos^{1,2*}

¹ Departamento de Física e Astronomia, Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre, 4169-007 Porto, Portugal ² Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, 4150-762 Porto, Portugal ³ CAB-CSIC ¶ Corresponding author * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Open Journals](#) ↗

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))⁰.

Summary

Stellar variability can impact planetary signals detected via the RV method. This is often addressed by tracking spectral lines sensitive to magnetic or/and temperature changes in the stellar atmosphere. With the growing use of NIR instruments like NIRPS, understanding NIR activity indicators is crucial, as their sensitivity may vary with stellar properties.

AMATERASU (AutoMATIC Equivalent-width Retrieval for Activity Signal Unveiling) is a Python tool to check for periods in spectral activity indices similar to an input period. This way, by running AMATERASU for a specific spectral line, the user can see if the input period may be correlated with activity. AMATERASU follows a methodology similar to ("Blind search of activity-sensitive lines in the NIR using HARPS and NIRPS observations of Proxima and Gl 581," 2025, accepted) and is inspired by ACTIN ((João Gomes da Silva et al., 2018), (J. Gomes da Silva et al., 2021)).

Statement of need

Description

AMATERASU computes the equivalent-width (EW) of a spectral line in a normalization independent way, by using the 90th percentile of the flux in a given window as the continuum level. It then computes the EW for an array of bandpasses, going from 0.1 Å up to a user defined width. This way, the input includes the spectral line center, maximum bandpass width window and a window that includes both the line and some continuum flux. The flux is interpolated inside the window using the `specutils` package, with a step similar to the original spectrum's step, to align the spectrum portion with the bandpass edges.

The maximum bandpass and interpolation window can be given manually or automatically. In the automatic way, the spectra are coadded into a master spectrum, using the first spectrum as the common reference grid, which is then smoothed with a moving average. Then, the `find_peaks` function from the `scipy` package is used to find the spectral lines and then retrieves the FWHM of the closest line to be studied (with a threshold of 0.1 Å). The bandpass window is a multiple (rounded) of the FWHM retrieved (by default 5 times) and the interpolation window is triple that (15 times the FWHM). Having retrieved a time series of EWs measurements for a given bandpass, AMATERASU cleans the data by applying a 3- σ sequential clipping and binning the data by night.

Finally, AMATERASU runs Generalized Lomb-Scargle (GLS) periodograms (Zechmeister & Kürster, 2009) and extracts the significant peaks. These are the ones not close to peaks in the

40 window function periodogram and with a false alarm probability (FAP) under the specified
41 FAP threshold. If any of the significant peaks is close within the specified threshold to the
42 input period, the program warns the user.

43 The user can choose one of the predefined indices in the `ind_table.csv` table or test a new
44 indice. The tool accepts a list of input periods and a list of lines to analyse, as well as 1D or 2D
45 spectra. To be more flexible with the instruments tested, the spectra have to be given in an array
46 format: for 2D spectra, the array should have dimensions of $(N_{\text{obs}}, N_{\text{spectrum axis}}, N_{\text{orders}}, N_{\text{pixels}})$,
47 where $N_{\text{spectrum axis}}$ refers for the wavelength, flux and flux error of the each spectrum. An
48 array with the dates of observations in BJD is also needed.

49 Besides the GLS periodograms, AMATERASU can also compute the Spearman ranked correla-
50 tion coefficient between all the central bandpasses and an input array, for example a known
51 activity indice, like the CCF FWHM of the observations.

52 AMATERASU has two modes: standard and full output. The standard output is the fastest,
53 as it only computes the EW time-series and correspondent GLS periodograms, prints whether
54 any of the periods is close to the input one and saves (if wanted) a `.csv` file with the matching
55 periods and respective central bandpasses and a `.csv` file with the correlations with the input
56 array per bandpass (if wanted). The full output allows to save all analysis data in a directory,
57 including:

- 58 ■ table with the EW values and errors for each central bandpass, as well as the BJD time
- 59 ■ plot of time-series of each EW computed and corresponding GLS periodogram
- 60 ■ separate periodogram and WF periodogram for each EW (for debugging purposes)
- 61 ■ table with the periods matching input
- 62 ■ table with all significant periods found
- 63 ■ table with all the information on the periodograms
- 64 ■ table with the Spearman correlations per bandpass with input array

65 The code is available and will be updated on [GitHub](#) and can be easily installed using `pip`.

66 References

- 67 Blind search of activity-sensitive lines in the NIR using HARPS and NIRPS observations of
68 Proxima and Gl 581. (2025). *Astronomy and Astrophysics*, 999, 999.
- 69 Gomes da Silva, João, Figueira, P., Santos, N., & Faria, J. (2018). ACTIN: A tool to
70 calculate stellar activity indices. *The Journal of Open Source Software*, 31, 667. <https://doi.org/10.21105/joss.00667>
- 72 Gomes da Silva, J., Santos, N. C., Adibekyan, V., Sousa, S. G., Campante, T. L., Figueira, P.,
73 Bossini, D., Delgado-Mena, E., Monteiro, M. J. P. F. G., de Laverny, P., Recio-Blanco, A.,
74 & Lovis, C. (2021). Stellar chromospheric activity of 1674 FGK stars from the AMBRE-
75 HARPS sample. I. A catalogue of homogeneous chromospheric activity. *Astronomy and*
76 *Astrophysics*, 646, a77. <https://doi.org/10.1051/0004-6361/202039765>
- 77 Zechmeister, M., & Kürster, M. (2009). The generalised Lomb-Scargle periodogram. A new
78 formalism for the floating-mean and Keplerian periodograms. *Astronomy and Astrophysics*,
79 496, 2. <https://doi.org/10.1051/0004-6361:200811296>