The Journal of Open Source Software

AMATERASU: a tool to track stellar activity

Telmo Monteiro [1,2], **João Gomes da Silva** [2], **Elisa Delgado-Mena** [3], **Nuno C. Santos** [1,2], and **Ana Soldevilla** [3]

1 Departamento de Física e Astronomia, Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre, 4169-007 Porto, Portugal **2** Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, 4150-762 Porto, Portugal **3** Centro de Astrobiología (CAB), CSIC-INTA, Camino Bajo del Castillo s/n, 28692, Villanueva de la Cañada (Madrid), Spain

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Open Journals

Reviewers:

- @openjournals

Submitted: 01 January 1970
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

AMATERASU (AutoMATic Equivalent-width Retrieval for Activity Signal Unveiling) is a Python tool to check for periods in spectral activity indices similar to an input period. This way, by running AMATERASU for a specific spectral line, the user can see if the input period may be correlated with activity. The aim of this program is to provide an easy and quick way to check for clues that the period observed in, for example, RV is of stellar origin. AMATERASU follows a methodology similar to João Gomes da Silva et al. (2025) and is inspired by ACTIN (J. Gomes da Silva et al., 2018, 2021).

Statement of need

Stellar variability can impact planetary signals detected via the radial velocity (RV) method. This is often addressed by tracking spectral lines sensitive to magnetic or/and temperature changes in the stellar atmosphere.

For this, different methods are currently used. For example, one can examine the variability in the cross-correlation function (CCF) parameters. The use of individual (or doublets and triplets) spectral lines as proxies for stellar activity is also popular, either in the optical or in the near infrared. Regarding these last ones, one can measure activity related line distortions through properties like the full width at half maximum (FWHM), depth or bisector, or through overall changes in the flux in the line, like equivalent-widths (EW) or simple flux ratios.

With the exoplanet hunters community pushing through to find Earth twins, understanding activity indicators behavior is crucial, as their sensitivity may vary with stellar properties, to detrend stellar activity from the signal from a orbiting planet.

In this context, we built the AutoMATic Equivalent-width Retrieval for Activity Signal Unveiling code (AMATERASU). This tool enables users to efficiently compute pseudo equivalent-widths (pEW) of spectral lines, search for periodicities in their time-series and identify whether those periods match an input. By combining instrument independent inputs, automatic adjustments and batch analysis of multiple lines and periods, AMATERASU provides a user-friendly and quick way to validate the stellar or planetary origin of RV signals.

State of the field

Some tools already exist to compute spectral indices, like ACTIN (J. Gomes da Silva et al., 2018, 2021) and iSTARMOD (Labarga & Montes, 2025, 2026). ACTIN computes the ratio between the flux inside a spectral line, delimited by a given central bandpass, and the flux in reference regions, usually the spectral continuum. iSTARMOD uses the spectral subtraction

technique, which involves subtracting the total flux emitted along a line by its photospheric contribution, obtained via synthesis of a quiescent version of the spectrum from a reference star with the same properties (spectral type, RV and rotational velocity). The final activity index is the EW of the residual profile.

While these tools are very useful for a variety of tasks, they only compute one version of a given activity index (with a fixed central bandpass), which may not be optimal to follow activity in different stars [J. Gomes da Silva et al. (2022); Monteiro et al., in prep]. Additionally, when testing a new index with ACTIN, the user needs to define the best reference regions, which is not straightforward when dealing with spectra affected by telluric lines (or their correction residuals) or lacking well defined continuum (often the case in M dwarves spectra).

Moreover, these tools do not provide a straightforward way to systematically test whether a given periodicity, such as a candidate planet signal, may be reproduced in activity indices.

AMATERASU fills this gap, allowing easy computation of spectral pEWs and search for periodicities in their time-series. In practice, AMATERASU tests if a given period can be obtained by adjusting different central bandpasses from a grid to a given target line, thereby finding, or not, if the period is also manifested through line deformations.

Software design

talk more about modularity of code

AMATERASU computes the equivalent-width (EW) of a spectral line in a normalization independent way, by using the 80th percentile of the flux in a given window as the continuum level. It then computes the EW for an array of bandpasses, going from 0.1 Å up to a user defined width. This way, the input includes the spectral line center, maximum bandpass width window and a window that includes both the line and some continuum flux. The flux is interpolated inside the window using the `specutils` package, with a step similar to the original spectrum's step, to align the spectrum portion with the bandpass edges.

The maximum bandpass and interpolation window can be given manually or automatically. In the automatic way, the spectra are coadded into a master spectrum, using the first spectrum as the common reference grid, which is then smoothed with a moving average. Then, the `find_peaks` function from the `scipy` package is used to find the spectral lines and then retrieves the FWHM of the closest line to be studied (with a threshold of 0.1 Å). The bandpass window is a multiple (rounded) of the FWHM retrieved (by default 5 times) and the interpolation window is 6 times that (30 times the FWHM). Having retrieved a time series of EWs measurements for a given bandpass, AMATERASU cleans the data by applying a 3-σ sequential clipping and binning the data by night.

Finally, AMATERASU runs Generalized Lomb-Scargle (GLS) periodograms (Zechmeister & Kürster, 2009) and extracts the significant peaks. These are the ones not close to peaks in the window function periodogram and with a false alarm probability (FAP) under the specified FAP threshold. If any of the significant peaks is close within the specified threshold to the input period, the program warns the user.

The user can choose one of the predefined indices in the `ind_table.csv` table or test a new indice. The tool accepts a list of input periods and a list of lines to analyse, as well as 1D or 2D spectra. To be more flexible with the instruments tested, the spectra have to be given in an array format: for 2D spectra, the array should have dimensions of $(N_{\text{obs}}, N_{\text{spectrum axis}}, N_{\text{orders}}, N_{\text{pixels}})$, where $N_{\text{spectrum axis}}$ refers for the wavelength, flux and flux error of each spectrum. An array with the dates of observations in BJD is also needed.

Besides the GLS periodograms, AMATERASU can also compute the Spearman ranked correlation coefficient between all the central bandpasses and an input array, for example a known activity indice, like the CCF FWHM of the observations.

AMATERASU has two modes: standard and full output. The standard output is the fastest, as it only computes the EW time-series and correspondent GLS periodograms, prints wether any of the periods is close to the input one and saves (if wanted) a `.csv` file with the matching periods and respective central bandpasses and a `.csv` file with the correlations with the input array per bandpass (if wanted). The full output allows to save all analysis data in a directory, including:

- table with the EW values and errors for each central bandpass, as well as the BJD time
- plot of time-series of each EW computed and corresponding GLS periodogram
- separate periodogram and WF periodogram for each EW (for debugging purposes)
- table with the periods matching input
- table with all significant periods found
- table with all the information on the periodograms
- table with the Spearman correlations per bandpass with input array

So far, we only tested AMATERASU with NIRPS data, so the predefined indices consist in NIR lines. Additionally, AMATERASU's automatic window definition only works for simmetrical absorption lines with high depth, so it will fail for lines with assimetrical profiles, with core emission or with blends.

The code is available and will be updated on GitHub and can be easily installed using pip.

Research impact statement

This tool could decrease false-positives arising from misidentified periodicities in RV and help improve planet search surveys.

something about helping RV surveys and stellar variability studies.

already being used in on-going works (Monteiro et al., in prep; Soldevilla et al., in prep.)

AI usage disclosure

Generative AI tools were used during development for code completion and debugging. All AI-generated content was reviewed and refined by the authors.

References

Gomes da Silva, J., Bensabat, A., Monteiro, T., & Santos, N. C. (2022). Optimising the Hα index for the identification of activity signals in FGK stars. Improvement of the correlation between Hα and Ca II H&K. *Astronomy and Astrophysics*, *668*, a174. https://doi.org/10.1051/0004-6361/202244595

Gomes da Silva, João, Delgado-Mena, E., Santos, N. C., Monteiro, T., Larue, P., Suárez Mascareño, A., Delfosse, X., Mignon, L., Artigau, É., Nari, N., Abreu, M., Aguiar, J. L. A., Al Moulla, K., Allain, G., Allart, R., Arial, T., Auger, H., Baron, F., Barros, S. C. C., … Zins, G. (2025). Blind search for activity-sensitive lines in the near-infrared using HARPS and NIRPS observations of Proxima and Gl 581. *700*, A177. https://doi.org/10.1051/0004-6361/202555013

Gomes da Silva, J., Figueira, P., Santos, N., & Faria, J. (2018). ACTIN: A tool to calculate stellar activity indices. *The Journal of Open Source Software*, *31*, 667. https://doi.org/10.21105/joss.00667

Gomes da Silva, J., Santos, N. C., Adibekyan, V., Sousa, S. G., Campante, T. L., Figueira, P., Bossini, D., Delgado-Mena, E., Monteiro, M. J. P. F. G., de Laverny, P., Recio-Blanco, A.,

DRAFT

& Lovis, C. (2021). Stellar chromospheric activity of 1674 FGK stars from the AMBRE-HARPS sample. I. A catalogue of homogeneous chromospheric activity. *Astronomy and Astrophysics*, *646*, a77. https://doi.org/10.1051/0004-6361/202039765

Labarga, F., & Montes, D. (2025). *iSTARMOD: Chromospheric activity in cool stars*. Astrophysics Source Code Library, record ascl:2512.002.

Labarga, F., & Montes, D. (2026). iSTARMOD: A Python Code to Quantify Chromospheric Activity by Using the Spectral Subtraction Technique. *171*(1), 15. https://doi.org/10.3847/1538-3881/ae173e

Zechmeister, M., & Kürster, M. (2009). The generalised Lomb-Scargle periodogram. A new formalism for the floating-mean and Keplerian periodograms. *Astronomy and Astrophysics*, *496*, 2. https://doi.org/10.1051/0004-6361:200811296

DRAFT