Kode Bibit

C0001

Tanggal

Fri Nov 20 2020 11:35:16 GMT+0700 (Indochina Time)

Longitude **164°45.25** E

latitude

70°30.5 S

Suspect Area Identification		
I. Does an identifiable circulation already exist at any level in the low or middle levels?		
II. Is there currently a deep convective disturbance which has persisted for at least 12 hours?	>	>
III. Does NWP indicate the development of any persistent low/mid-level circulation during the next 72 hours?		
Location of suspect area	Jakarta	Jakarta
Broadscale Environmental Conditions - Current		
A. Is the MJO active in our longitude?		
B. Are there any tropical waves currently propagating towards the suspect area?	~	>
C. Are there any strong low-level monsoon/trade flows or surges feeding into the suspect area now?		
D. Is there a low/mid-level trough in the suspect area now?		
E. Are the winds at 200/250 hPa over suspect the area <20 knots?		
F. Is there low (5-15 kt) 850-200 hPa vertical shear near the area and stronger (>15 kt) shear away (5-10°) from the suspect area?		
G. Is there a synoptic feature creating strong upper divergence above the suspect area? For example, an upper trough or retrogressing upper low within 25 degrees to the west, or a strong equatorial jet, with the ridge centre over the area.	\searrow	>
H. Is there evidence on satellite imagery or upper streamline charts of at least one outflow channel?		
I. Is there evidence of two outflow channels or fanning of cirrus (equatorward and eastward of CDO)? If so consider rapid development potential.		
J. Is the suspect area being fed by deep moist air in all sectors?		
Broadscale Environmental Conditions - 72 Hour Trend		
K. Are any tropical waves likely to move into the area within 72 hours?		
L. Are there any monsoon or trade surges likely to feed into the area during the next 72 hours?		
M. Will the 850-200 hPa shear conditions remain favourable / improve during the next 72 hours?	>	>
N. Will the upper divergence remain/become favourable for cyclogenesis during the next 72 hours?		
O. Is deep moisture infeed into the area likely to remain / improve during the next 72 hours?		
P. Do NWP cyclogenesis parameters support further development in the area during the 72 hours period?	>	>
Development of Circulation		
1. Does a vertically stacked circulation currently exist from surface to 500 hPa?		
2. Is there a low/mid-level circulation centre defined by cloud line or band curvature, or by cloud line mergence within an area of 2.5° latitude or less? (This is best seen on vis imagery.)		
3. Are there (disturbance-relative) wind speed observations of the order 15-25 kts (850 hPa) within 5° of the circulation centre?	~	>

4. Is there evidence of a significant (15-25 kt) middle-level circulation (850-500 hPa) present? (Vertically stacked and round.)			
5. Have there been pressure falls of greater than 2 hPa per day within 3° of the system centre?			
Development of Deep Convection			
6. Has there been an Early Convective Maximum (ECM) within the last 3 days? - i.e. was there deep convection (-65°C or colder/Dvorak black) focused in one area (within about a 2° radius) and persisting for 6-12 hours possibly against the diurnal cycle?			
7. Has there been a reduction in the area of deep convection after the occurrence of the ECM? (This quiet period may continue 1-3 days.)			
8. Is there currently persistent deep convection (at least 12 hours old) which is within 2° of a low level cloud system centre?		100	
Movement on water			
9. The disturbance remain over water or move over water			
10. The disturbance will not remain over water or not move over water		>	>
Checklist Result	Suggested Action		
ticks for 6 and 7, the broadscale environment is favourable for further development (many ticks for A-P) and the disturbance could remain or move over water (tick 9)	Zehr's Stage 1: TC development possible within 1-3 days - consider High-Key Standby (note: but The disturbance will not remain over water or not move over water)		