

THE PECULIAR TYPE II SUPERNOVA 1997D: A CASE FOR A VERY LOW ^{56}Ni MASS¹

M. TURATTO,^{2,3,4} P. A. MAZZALI,^{4,5,6} T. R. YOUNG,^{5,7} K. NOMOTO,^{4,7,8} K. IWAMOTO,^{4,7} S. BENETTI,² E. CAPPELLARO,^{3,4}
 I. J. DANZIGER,⁶ D. F. DE MELLO,⁹ M. M. PHILLIPS,¹⁰ N. B. SUNTZEFF,¹⁰ A. CLOCCHIATTI,^{10,11} A. PIEMONTE,²
 B. LEIBUNDGUT,¹² R. COVARRUBIAS,¹⁰ J. MAZA,¹³ AND J. SOLLERMAN¹⁴

Received 1998 January 14; accepted 1998 March 16; published 1998 April 17

ABSTRACT

SN 1997D in NGC 1536 is possibly the least luminous and energetic Type II supernova discovered to date. The entire light curve is subluminal, never reaching $M_V = -14.65$. The radioactive tail follows the ^{56}Co decay slope. In the case of a nearly complete trapping of the γ -rays, the ^{56}Ni mass derived from the tail brightness is extremely small, $\sim 0.002 M_\odot$. At discovery, the spectra showed a red continuum and line velocities on the order of 1000 km s^{-1} . The luminosity and the photospheric expansion velocity suggest that the explosion occurred about 50 days before discovery and that a plateau probably followed. Model light curves and spectra of the explosion of a $26 M_\odot$ star successfully fitted the observations. Low-mass models are inconsistent with the observations. The radius of the progenitor, constrained by the prediscussion upper limits, is $R_0 \lesssim 300 R_\odot$. A low explosion energy of $\sim 4 \times 10^{50}$ ergs is then required in the modeling. The strong Ba II lines in the photospheric spectra are reproduced with a solar abundance and low T_{eff} . A scenario in which the low ^{56}Ni mass observed in SN 1997D is due to fallback of material onto the collapsed remnant of the explosion of a $25\text{--}40 M_\odot$ star appears to be favored over the case of the explosion of an $8\text{--}10 M_\odot$ star with low ^{56}Ni production.

Subject headings: nuclear reactions, nucleosynthesis, abundances — stars: evolution — supernovae: general — supernovae: individual (SN 1997D)

1. INTRODUCTION

The Type II supernova SN 1997D was serendipitously discovered on January 14.15 UT (De Mello & Benetti 1997) in NGC 1536, a morphologically disturbed spiral galaxy belonging to a high-density group (Maia et al. 1994). Although NGC 1536 ($v_{\text{helio}} = 1296 \text{ km s}^{-1}$; de Vaucouleurs et al. 1991, hereafter RC3) is one of the galaxies patrolled visually by Rev. Evans for his SN search (R. Evans 1997, private communication), he missed the SN because its brightness apparently never exceeded his detection limit.

The first spectrum showed the main features of Type II SNe but also revealed peculiarities that make this object unique. The

extremely slow expansion velocities, the red color, and the strong Ba II lines were particularly noteworthy (Turatto et al. 1998). A campaign of photometric and spectroscopic observations was therefore promptly started at ESO and CTIO. The complete data set will be presented elsewhere.

2. PROPERTIES AND EVOLUTION

The SN was intrinsically faint compared with other SNe II. If we assume a distance modulus of $\mu = 30.64$ mag for NGC 1536 ($H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$; Tully 1988), and $A_B = 0.00$ (RC3), a set of closely spaced nondetections by Evans sets an upper limit to the SN luminosity of $M_V(\text{max}) \gtrsim -14.65$. This is about 2 mag fainter than regular SNe II at maximum (Patat et al. 1994).

In Figure 1, we compare the absolute V light curve of SN 1997D with those of other SNe II: the double-peaked SN 1987A, the prototypical SN IIb 1993J, and the linear SN 1979C. Even with the uncertainties on the epoch of maximum and in the distance modulus, SN 1997D was definitely fainter than all of these objects. The decline rates in the V , R , and I bands after JD 2,450,520 are close to the ^{56}Co decay rate.

The color of SN 1997D at discovery was unusually red. No sign of interstellar absorption lines, which are expected in the case of strong extinction, is present in the spectra. SN 1997D reddened rapidly over the first few weeks after discovery, reaching a maximum ($B - V$) = 2.5 in mid-February, and then turned gradually bluer. The analogy with the color curves of other SNe II, which reach the reddest point between 40 and 80 days after peak luminosity, indicates that maximum light should have occurred around JD 2,450,430 (± 20 days).

A bolometric light curve is needed for a proper analysis of the energetics. Unfortunately, most of our observations cover only the optical domain. Only at one phase are nearly simultaneous optical (January 31) and IR (January 27) spectra available. The overall luminosity between 0.35 and 2.3μ in the combined spectrum is $\log L_{\text{bol}} = 40.83 \text{ ergs s}^{-1}$. At that epoch,

¹ Based on observations collected at ESO–La Silla and Cerro Tololo Inter-American Observatory (CTIO) (Chile). CTIO, a part of the National Optical Astronomy Observatories, is operated by the Association of Universities for Research in Astronomy (AURA), Inc., under cooperative agreement with the National Science Foundation.

² European Southern Observatory (ESO), Alonso de Cordova 3107, Vitacura, Casilla 19001, Santiago 19, Chile.

³ Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, I-35122 Padova, Italy.

⁴ Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106-4030.

⁵ National Astronomical Observatory, Mitaka-shi, Tokyo 181, Japan.

⁶ Osservatorio Astronomico di Trieste, via G. B. Tiepolo 11, 34131 Trieste, Italy.

⁷ Department of Astronomy, University of Tokyo, Bunkyo-ku, Tokyo 113, Japan.

⁸ Research Center for the Early Universe, University of Tokyo, Bunkyo-ku, Tokyo 113, Japan.

⁹ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218.

¹⁰ Cerro Tololo Inter-American Observatory, Casilla 603, La Serena, Chile.

¹¹ Departamento de Astronomia, Pontificia Universidad Católica de Chile, Casilla 104, Santiago 22, Chile.

¹² European Southern Observatory, Karl-Schwarzschild-Strasse 2, 85748 Garching bei München, Germany.

¹³ Departamento de Astronomia, Universidad de Chile, Casilla 36-D, Santiago, Chile.

¹⁴ Stockholm Observatory, S-13336 Saltjöbaden, Sweden.




FIG. 1.—The comparison of the absolute V light curve of SN 1997D with those of SNe 1987A, 1993J, and 1979C. The epochs of maxima have been corrected to JD 2,450,042.0. Only galactic absorption has been considered. The upper limits from the negative observations of Rev. R. Evans (1997, private communication) have been also plotted. The insert shows the bolometric light curves of SN 1997D and SN 1987A (Catchpole et al. 1987, 1988). The star marks the optical + IR spectrophotometric observation of January 30.

the color is still sufficiently blue that Schmidt's (1998) empirical bolometric corrections to V can be used. One then obtains $\log L_{\text{bol}} = 40.96 \text{ ergs s}^{-1}$, in good agreement with the spectrophotometry. At later phases, the color of SN 1997D was so red that Schmidt's correction cannot be applied. However, the available spectra show that at all epochs, most of the energy was radiated in the V and R bands and that the color evolution is mostly due to variations in the B band, where the flux is much smaller. Therefore, during the photospheric phase, we assumed that the bolometric light curve can be described reasonably well by applying a constant bolometric correction to the V magnitudes. The bolometric light curve thus derived is shown in the insert of Figure 1. The last point has been obtained by adopting the same L_{bol}/L_{BVRI} ratio for SN 1997D as for SN 1987A in the nebular epoch (Schmidt 1998).

Although, in SN 1987A, the γ -rays from the ^{56}Co decay were not completely deposited (Pinto, Woosley, & Ensmann 1988), the late time decline of SN 1987A was close to the decay rate of ^{56}Co . The mass of ^{56}Ni ejected by SN 1987A was $\sim 0.075 M_{\odot}$ (Bouchet, Danziger, & Lucy 1991a; Bouchet et al. 1991b; Bouchet & Danziger 1993; Danziger et al. 1991; Nomoto et al. 1994). Assuming for SN 1997D the same γ -ray deposition as in SN 1987A, the ratio of the luminosities on the tail of the light curves can be used to determine the mass of ^{56}Ni synthesized in the explosion of SN 1997D. We then obtain a ratio $M_{^{56}\text{Ni}}(87A)/M_{^{56}\text{Ni}}(97D) \sim 40$. Taking into account the uncertainties in the determination of the epoch of maximum and of the bolometric luminosity, the mass of ^{56}Ni ejected by SN 1997D is very small, between 0.001 and 0.004 M_{\odot} .

A comparison with SN 1987A (Fig. 2) shows general similarities with respect to P Cygni lines of H I, Ca II, Na I, and Ba II and more subtle differences. The latter are emphasized by the strengths of the Ba II absorption lines and possibly those of other s -process elements. SN 1987A shows strong lines of Ba II, the analysis of which (Williams 1987; Lucy 1988; Höflich 1988; Mazzali & Chugai 1995) suggested an enhanced abundance of barium. SN 1997D shows even stronger lines of Ba II, with the identification of an array of multiplets, but, as




FIG. 2.—The comparison of the spectra of SN 1997D and SN 1987A. *Top*: the spectrum of SN 1997D at discovery is compared with that of 1987A 1.5 months past explosion. *Middle*: spectra taken 100 days later. The spectrum of SN 1997D is the result of the merging of two spectra with different resolutions and signal-to-noise ratios. *Bottom*: spectra taken 10 months past maximum. All spectra have been normalized to the peak of the $H\alpha$ emission and corrected for redshift.

we shall discuss later, an overabundance of barium is not required because a lower temperature prevails in the envelope.

Strong lines of Sr II, also an s -process element, were also observed in the IR spectrum taken on January 27 (Clocchiatti et al. 1998). In this spectrum, the He I 10830 Å line, which is the dominant feature in the IR spectra of other SNe II, is not clearly detected, although it might affect the blue wing of Sr II $\lambda 10914$.

Finally, one of the most striking peculiarities of SN 1997D is the low expansion velocity of the ejecta, approximately 1200 km s^{-1} , as measured from the line absorption minima. Only the strong Ba II lines show somewhat higher velocities, but it is well known that strong lines may form in outer parts of the envelope and that stratification effects may also be present. As we will discuss later (§ 3.2), the small velocity constrains the epoch of the first spectrum to be ~ 50 days.

3. MODEL LIGHT CURVE AND SPECTRA

The light curve of SN 1997D is unique because the luminosity is low, both at “maximum light” and on the radioactive tail. These properties, along with the narrowness of the lines and the redness of the continuum, can be used to deduce the physical characteristics of the progenitor and of the explosion.

The progenitor model used here is that of a 26 M_{\odot} star with an 8 M_{\odot} He core at the beginning of collapse (Nomoto & Hashimoto 1988) and an 18 M_{\odot} H-rich envelope. The stellar radius is $R_0 = 300 R_{\odot}$, which is 3–5 times smaller than typical red supergiants. The explosion was simulated using a one-dimensional Lagrangian hydrodynamical code (Young, Baron, & Branch 1995) with an energy of $E = 4 \times 10^{50}$ ergs and the ejection of 0.002 M_{\odot} of ^{56}Ni . In order to eject a small amount of ^{56}Ni , most of the material of the innermost region where Si is burned to Ni has to fall back onto the degenerate neutron core, where it is photodisintegrated before emitting γ -rays when forming the compact remnant. This is used to place the mass cut, resulting in a collapsing core of 1.8 M_{\odot} and an ejecta mass of 24.2 M_{\odot} .

els with a smaller mass give lines too shallow if $v_{\text{ph}} = 970 \text{ km s}^{-1}$.

Synthetic spectra show that the requirement that $t \sim 50$ days on January 17 is a rather strong one. If the explosion took place closer to discovery, i.e., $t < 50$ days, the observed expansion velocity implies a small radius. Together with the derived luminosity, this results in a higher T_{eff} than required. In this case, the Ba III/Ba II ratio increases, and the Ba II line intensities decrease. If, on the other hand, the explosion occurred earlier, so that $t > 50$ days, the radius is too large, and the temperature becomes too low. When the temperature drops below 5000 K, lines of Ti I and Fe I absorb most of the radiation below 4500 Å, and the synthetic spectrum changes dramatically.

We also computed a synthetic spectrum of our selected explosion model evolved to match the February 6 data. The fitting was satisfactory using an epoch consistent with $t = 50$ days on January 17.

4. DISCUSSION

SN 1997D is among the least luminous Type II supernovae known to date. The low luminosity persists in the tail also, where the decline rate is close to that of a ^{56}Co -powered tail. SN 1997D is also characterized by a low temperature and a very low expansion velocity. These features are well reproduced by an explosion model with a relatively small progenitor radius ($300 R_{\odot}$), massive ejecta ($\sim 24 M_{\odot}$), low explosion energy (4×10^{50} ergs), and small ejected ^{56}Ni mass ($\sim 0.002 M_{\odot}$).

A similarly small amount of ^{56}Ni has recently been proposed for another Type II supernova, SN 1994W (Sollerman, Cumming, & Lundqvist 1998), but this object was rather luminous, and its tail declined faster than the ^{56}Co slope.

The neutrino-heating mechanism of massive star explosions is not well understood, so the explosion energy and the amount of ^{56}Ni ejected as a function of progenitor mass are difficult to predict from the hydrodynamical models. According to pre-supernova model calculations, stars more massive than $\sim 25 M_{\odot}$ form an Fe core in a greater gravitational potential because of the significantly smaller C/O ratio and the consequently weak carbon shell burning (Nomoto et al. 1993; Woosley & Weaver 1995). Therefore, if the efficiency of neutrino heating does not change with exploding mass, more massive stars tend to produce lower explosion energies, and they suffer from more fallback of material onto the collapsed remnant. The mass cut is then placed farther out. However, the progenitor must be less massive than the Wolf-Rayet progenitors (i.e., $\lesssim 30\text{--}40 M_{\odot}$) because of the presence of an envelope with a mass of at least $15 M_{\odot}$ above the photosphere at day 50.

Stars in the range $\sim 10\text{--}25 M_{\odot}$ are likely to explode with typical explosion energies of $\sim 1 \times 10^{51}$ ergs and to produce $0.07\text{--}0.15 M_{\odot}$ of ^{56}Ni , as indicated by the brightness and light-curve shapes of SN 1993J, SN 1987A, and of Type Ibc supernovae (Nomoto, Iwamoto, & Suzuki 1995; Young et al. 1995). Stars of $8\text{--}10 M_{\odot}$ are predicted to produce little ^{56}Ni and other heavy elements (Nomoto 1984; Mayle & Wilson 1988). However, these stars are at the top of the asymptotic giant branch in the presupernova stages (Hashimoto, Iwamoto, & Nomoto 1993), which is inconsistent with the small radius of SN 1997D. Also, synthetic spectra obtained from the low-mass models have shallow absorption lines compared with the observations.

In conclusion, the progenitor of SN 1997D was probably as massive as $25\text{--}40 M_{\odot}$. Our $26 M_{\odot}$ model is in this mass range, thus being consistent with the low E/M and the small ^{56}Ni mass. The presupernova radius depends on many parameters, and the reason for its being rather small in SN 1997D is an open question. Spectral models appear to rule out the case of low metallicity and He enhancement. A possible scenario involves a close binary system in which the companion star spiraled in to make the envelope more massive (Podsiadlowski 1992).

The ejection of a small mass of ^{56}Ni provides a constraint on the mass of the collapsed remnant, M_{rem} (Nomoto et al. 1993; Thielemann, Nomoto, & Hashimoto 1996). In our $26 M_{\odot}$ model, $M_{\text{rem}} = 1.8 M_{\odot}$. If M_{rem} were assumed to be smaller, more ^{56}Ni would be ejected; for example, if $M_{\text{rem}} = 1.7 M_{\odot}$, then $M(^{56}\text{Ni}) = 0.1 M_{\odot}$. For more massive progenitor models, M_{rem} would be larger than $1.8 M_{\odot}$. If the equation of state of nuclear matter is relatively soft, the maximum (gravitational) mass of a cold neutron star is $\sim 1.5 M_{\odot}$ (Brown & Bethe 1994). If this is the case, the remnant of SN 1997D must be a small-mass black hole, and the low-luminosity tail of SN 1997D might be a signature of the formation of such an object.

We would like to thank Piero Rosati for observing SN 1997D and Rev. Robert Evans for providing his prediscovery limits. P. A. Mazzali acknowledges receipt of a Foreign Research Fellowship at NAO and is grateful to T. Kajino and the Department of Astronomy at the University of Tokyo for their hospitality. A. Piemonte acknowledges financial support from ESO during his stay in Chile. This work has been supported in part by the Grant-in-Aid for Scientific Research (05242102, 06233101) and COE research (07CE2002) of the Ministry of Education, Science, and Culture in Japan and by the National Science Foundation in the US under PHY 94-07194. Parts of the observations have been obtained at the ESO 1.5 m telescope operated under the agreement between ESO and Observatorio Nacional-Brasil.

REFERENCES

- Bouchet, P., & Danziger, I. J. 1993, *A&A*, 273, 451
 Bouchet, P., Danziger, I. J., & Lucy, L. B. 1991a, *AJ*, 102, 1135
 Bouchet, P., Phillips, M. M., Suntzeff, N. B., Gouffes, C., Hanuschik, R. W., & Wooden, D. H. 1991b, *A&A*, 245, 490
 Brown, G. E., & Bethe, H. A. 1994, *ApJ*, 423, 659
 Catchpole, R. M., et al. 1987, *MNRAS* 229, 15P
 ———. 1988, *MNRAS*, 231, 75P
 Clocciatti, A., Phillips, M. M., Spyromilio, J., & Leibundgut, B. 1998, in SN 1987A: Ten Years After, the Fifth CTIO/LCO Workshop, ed. N. B. Suntzeff & M. M. Phillips (San Francisco: ASP), in press
 Danziger, I. J., Lucy, L. B., Bouchet, P., & Gouffes, C. 1991, in *Supernovae*, ed. S. E. Woosley (New York: Springer), 69
 De Mello, D., & Benetti, S. 1997, *IAU Circ.* 6537
 de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H. G., Buta, R. J., Paturel, G., & Fouque, P. 1991, *Third Reference Catalogue of Bright Galaxies* (New York: Springer) (RC3)
 Hashimoto, M., Iwamoto, K., & Nomoto, K. 1993, *ApJ*, 414, L105
 Höflich, P. 1988, *Proc. Astron. Soc. Australia*, 7, 434
 Lucy, L. B. 1988, in *Proc. Fourth George Mason Conf., SN 1987A in the LMC*, ed. M. Kafatos (Cambridge: Cambridge Univ. Press), 323
 Maia, M. A. G., Pastoriza, M. G., Bica, E., & Dottori, H. 1994, *ApJS*, 93, 425
 Mayle, R., & Wilson, J. R. 1988, *ApJ*, 334, 909
 Mazzali, P. A., & Chugai, N. N. 1995, *A&A*, 303, 118
 Mazzali, P. A., & Lucy, L. B. 1993, *A&A*, 279, 447
 ———. 1998, in preparation

- Nomoto, K. 1984, *ApJ*, 277, 791
Nomoto, K., & Hashimoto, M. 1988, *Phys. Rep.*, 163, 13
Nomoto, K., Iwamoto, K., & Suzuki, T. 1995, *Phys. Rep.*, 256, 173
Nomoto, K., Iwamoto, K., Tsujimoto, T., & Hashimoto, M. 1993, in *Frontiers of Neutrino Astrophysics*, ed. Y. Suzuki (Tokyo: Universal Academy Press), 235
Nomoto, K., Shigeyama, T., Kumagai, S., Yamaoka, H., & Suzuki, T. 1994, in *Supernovae (Les Houches, Session LIV)*, ed. S. Bludman et al. (Amsterdam: Elsevier), 489
Patat, F., Barbon, R., Cappellaro, E., & Turatto, M. 1994, *A&A*, 282, 731
Pinto, P. A., Woosley, S. E., & Ensman, L. M. 1988, *ApJ*, 331, 101
Podsiadlowski, Ph. 1992, *PASP*, 104, 1
Schmidt, B. P. 1998, in *SN 1987A: Ten Years After, the Fifth CTIO/LCO Workshop*, ed. N. B. Suntzeff & M. M. Phillips (San Francisco: ASP), in press
Sollerman, J., Cumming, R. J., & Lundqvist, P. 1998, *ApJ*, 493, 933
Thielemann, F. K., Nomoto, K., & Hashimoto, M. 1996, *ApJ*, 460, 408
Tully, R. B. 1988, *Nearby Galaxy Catalog* (Cambridge: Cambridge Univ. Press)
Turatto, M., Benetti, S., Cappellaro, E., Danziger, I. J., & Mazzali, P. A. 1998, in *SN 1987A: Ten Years After, the Fifth CTIO/LCO Workshop*, ed. N. B. Suntzeff & M. M. Phillips (San Francisco: ASP), in press
Williams, R. 1987, *ApJ*, 320, L117
Woosley, S. E., & Weaver, T. A. 1995, *ApJS*, 101, 181
Young, T., Baron, E., & Branch, D. 1995, *ApJ*, 449, L51