

COMPUTATION OF DFT

DIRECT COMPUTATION OF DFT

$$X[k] = \sum_{n=0}^{N-1} x[n] W_N^{kn} \quad k = 0, 1, \dots, N-1$$

For each k ,

N complex multiplications (multiplication of two complex numbers)

$N - 1$ complex additions (addition of two complex numbers)

For all

N^2 complex multiplications

$N(N - 1)$ complex additions

Each complex multiplication = 4 real multiplications + 2 real additions

Each complex addition = 2 real additions

Therefore, for each k

$4N$ real multiplications

$4N - 2$ real additions $(2(N - 1) + 2N)$

In total

$4N^2$ real multiplications

$N(4N - 2)$ real additions $(N(4N - 2) \cong 4N^2 \text{ for } N \gg 1)$

If $x[n]$ is real, for each k

$2N$ real multiplications

$2(N - 1)$ real additions

In total

$2N^2$ real multiplications

$2N(N - 1)$ real additions $(2N(N - 1) \cong 2N^2 \text{ for } N \gg 1)$

FAST FOURIER TRANSFORM (FFT)

Periodicity and symmetry of W_N^q .

$$1) W_N^q = W_N^{q+N}$$

$$e^{-j\frac{2\pi}{N}(q+N)} = e^{-j\frac{2\pi}{N}q} \underbrace{e^{-j\frac{2\pi}{N}N}}_{=1}$$

$$\text{Ex: } W_{16}^4 = W_{16}^{20}$$

$$2) W_N^q = -W_N^{q+\frac{N}{2}}$$

$$e^{-j\frac{2\pi}{N}(q+\frac{N}{2})} = e^{-j\frac{2\pi}{N}q} \underbrace{e^{-j\pi}}_{=-1}$$

$$\text{Ex: } W_{16}^3 = -W_{16}^{11}$$

Let's have a close look at an 8-point DFT.

$$X[0] = x[0] + x[1] + x[2] + x[3] + x[4] + x[5] + x[6] + x[7]$$

$$X[1] = x[0] + x[1]W_8 + x[2]W_8^2 + x[3]W_8^3 + x[4]W_8^4 + x[5]W_8^5 + x[6]W_8^6 + x[7]W_8^7$$

$$X[2] = x[0] + x[1]W_8^2 + x[2]W_8^4 + x[3]W_8^6 + x[4]\underbrace{W_8^8}_{=1} + x[5]\underbrace{W_8^{10}}_{=W_8^2} + x[6]\underbrace{W_8^{12}}_{=W_8^4} + x[7]\underbrace{W_8^{14}}_{=W_8^6}$$

$$X[3] = x[0] + x[1]W_8^3 + x[2]W_8^6 + x[3]\underbrace{W_8^9}_{=W_8^1} + x[4]\underbrace{W_8^{12}}_{=W_8^4} + x[5]\underbrace{W_8^{15}}_{=W_8^7} + x[6]\underbrace{W_8^{18}}_{=W_8^2} + x[7]\underbrace{W_8^{21}}_{=W_8^5}$$

$$X[4] = x[0] + x[1]\underbrace{W_8^4}_{=-1} + x[2]\underbrace{W_8^8}_{=1} + x[3]\underbrace{W_8^{12}}_{=-1} + x[4]\underbrace{W_8^{16}}_{=1} + x[5]\underbrace{W_8^{20}}_{=-1} + x[6]\underbrace{W_8^{24}}_{=1} + x[7]\underbrace{W_8^{28}}_{=-1}$$

$$X[5] = x[0] + x[1]W_8^5 + x[2]\underbrace{W_8^{10}}_{=W_8^2} + x[3]\underbrace{W_8^{15}}_{=W_8^7} + x[4]\underbrace{W_8^{20}}_{=-1} + x[5]\underbrace{W_8^{25}}_{=W_8^1} + x[6]\underbrace{W_8^{30}}_{=W_8^6} + x[7]\underbrace{W_8^{35}}_{=W_8^3}$$

$$X[6] = x[0] + x[1]W_8^6 + x[2]\underbrace{W_8^{12}}_{=-1} + x[3]\underbrace{W_8^{18}}_{=W_8^2} + x[4]\underbrace{W_8^{24}}_{=1} + x[5]\underbrace{W_8^{30}}_{=W_8^6} + x[6]\underbrace{W_8^{36}}_{=-1} + x[7]\underbrace{W_8^{42}}_{=W_8^2}$$

$$X[7] = x[0] + x[1]W_8^7 + x[2]\underbrace{W_8^{14}}_{=W_8^6} + x[3]\underbrace{W_8^{21}}_{=W_8^5} + x[4]\underbrace{W_8^{28}}_{=-1} + x[5]\underbrace{W_8^{35}}_{=W_8^3} + x[6]\underbrace{W_8^{42}}_{=W_8^2} + x[7]\underbrace{W_8^{49}}_{=W_8^1}$$

Note that, since

$$W_8^4 = -W_8^0 \quad W_8^5 = -W_8^1 \quad W_8^6 = -W_8^2 \quad W_8^7 = -W_8^3$$

i.e.

$$W_N^q = -W_N^{q+\frac{N}{2}}$$

we effectively need only W_8^1, W_8^2, W_8^3 instead of $W_8^0, W_8^1, W_8^2, \dots, W_8^{49}$.

Rewriting

$$X[1] = (x[0] - x[4]) + (x[1] - x[5]) W_8^1 + (x[2] - x[6]) W_8^2 + (x[3] - x[7]) W_8^3$$

$$X[2] = (x[0] - x[2] + x[4] - x[6]) + (x[1] - x[3] + x[5] - x[7]) W_8^2$$

$$X[3] = (x[0] - x[4]) + (x[3] - x[7]) W_8^1 + (-x[2] + x[6]) W_8^2 + (x[1] - x[5]) W_8^3$$

$$X[4] = x[0] - x[1] + x[2] - x[3] + x[4] - x[5] + x[6] - x[7]$$

$$X[5] = (x[0] - x[4]) + (-x[1] + x[5]) W_8^1 + (x[2] - x[6]) W_8^2 + (-x[3] + x[7]) W_8^3$$

$$X[6] = x[0] - x[2] + x[4] - x[6] + (-x[1] + x[3] - x[5] + x[7]) W_8^2$$

$$X[7] = (x[0] - x[4]) + (-x[3] + x[7]) W_8^1 + (-x[2] + x[6]) W_8^2 + (-x[1] + x[5]) W_8^3$$

Therefore, the computation of the following expressions is sufficient to get all 8-point DFT values...

- | | |
|-----------------------------------------------------------|---------------------|
| 1) $x[0] + x[4], x[0] - x[4]$ | 2 adds |
| 2) $x[1] + x[5], x[1] - x[5]$ | 2 adds |
| 3) $x[2] + x[6], x[2] - x[6]$ | 2 adds |
| 4) $x[3] + x[7], x[3] - x[7]$ | 2 adds |
| 5) $(x[1] - x[5])W_8^1, (-x[1] + x[5])W_8^3$ | 4 multiplies |
| 6) $(x[3] - x[7])W_8^3, (-x[3] + x[7])W_8^1$ | 4 multiplies |
| 7) $(x[2] - x[6])W_8^2, (x[1] - x[3] + x[5] - x[7])W_8^2$ | 1 add, 4 multiplies |

plus “some” additions in the previous page.

We will study the two systematic methods

1) Decimation-in-time

2) Decimation-in-frequency

Decimation-in-Time

Assume that N is a power of two, i.e. $N = 2^m$.

$$X[k] = \sum_{n=0}^{N-1} x[n] W_N^{kn} \quad k = 0, 1, \dots, N-1$$

Decompose for even and odd indexed terms of $x[n]$:

$$\begin{aligned}
 X[k] &= \sum_{n:\text{even}}^{N-1} x[n] W_N^{kn} + \sum_{n:\text{odd}}^{N-1} x[n] W_N^{kn} \\
 &= \sum_{r=0}^{\frac{N}{2}-1} x[2r] W_N^{2kr} + \sum_{r=0}^{\frac{N}{2}-1} x[2r+1] W_N^{k(2r+1)} \\
 &= \underbrace{\sum_{r=0}^{\frac{N}{2}-1} x[2r] W_N^{\frac{kr}{2}}}_{G[k]} + W_N^k \underbrace{\sum_{r=0}^{\frac{N}{2}-1} x[2r+1] W_N^{\frac{kr}{2}}}_{H[k]}
 \end{aligned}$$


$G[k]: \frac{N}{2} - \text{point DFT of even indexed samples}$

$H[k]: \frac{N}{2} - \text{point DFT of odd indexed samples}$

Therefore

$$X[k] = G[k] + W_N^k H[k] \quad k = 0, 1, \dots, \underbrace{N-1}_{!!!}$$


Attention: $G[k]$ and $H[k]$ are $\frac{N}{2}$ —point DFTs.



$$2 \left(\frac{N}{2} \right)^2 + N = \frac{N^2}{2} + N \quad \text{complex multiplications required}$$

$\frac{N}{2}$ –point DFTs can be decomposed into $\frac{N}{4}$ –point DFTs.

Then the following computational flow arises.



$$4 \left(\frac{N}{4} \right)^2 + N + N = \frac{N^2}{4} + 2N$$

complex multiplications required

In general, the number of sections will be

$$\log_2 N = v$$

Therefore

$$\underbrace{\frac{N}{2} \left(\frac{N}{2} \right)^2}_{2N} + N + N + \dots + N = (v + 1)N$$

complex multiplications required

Considering the multiplications by -1 and 1, this value is approximately referred to as

$$N \log_2 N$$

This is also the approximate number of additions.

Ex:

$$N = 2^{10} = 1024$$


$$N^2 = 1\,048\,576 \approx 10^6$$

$$N \log_2 N = 10\,240 \approx 10^4$$


A reduction by a factor of 100 approximately.

“BUTTERFLY”s

Indeed, above diagrams contain many “butterfly” structures,



Such butterflies can be simplified as



DECIMATION-IN-FREQUENCY

Assume that N is a power of two, $N = 2^m$.

$$X[k] = \sum_{n=0}^{N-1} x[n] W_N^{kn} \quad k = 0, 1, \dots, N-1$$

Consider “even” and “odd” indexed elements of the DFT:

$$X[k] \begin{cases} \rightarrow X[2r] \\ \rightarrow X[2r + 1] \end{cases} \quad r = 0, 1, \dots, \frac{N}{2} - 1$$

Even indexed elements:

$$\begin{aligned}
 X[2r] &= \sum_{n=0}^{N-1} x[n] W_N^{2rn} \quad r = 0, 1, \dots, \frac{N}{2} - 1 \\
 &= \sum_{n=0}^{\frac{N}{2}-1} x[n] W_N^{2rn} + \sum_{n=\frac{N}{2}}^{N-1} x[n] W_N^{2rn} \\
 &= \sum_{n=0}^{\frac{N}{2}-1} x[n] W_N^{2rn} + \sum_{n=0}^{\frac{N}{2}-1} x\left[n + \frac{N}{2}\right] W_N^{2r\left(n + \frac{N}{2}\right)} \\
 &= \sum_{n=0}^{\frac{N}{2}-1} \left(x[n] + x\left[n + \frac{N}{2}\right] \right) W_{\frac{N}{2}}^{rn}
 \end{aligned}$$

This is an $\frac{N}{2}$ -point DFT of

$$\left(x[n] + x\left[n + \frac{N}{2}\right] \right) \quad n = 0, 1, \dots, \frac{N}{2} - 1.$$


Odd indexed elements: Similarly,

$$X[2r + 1] = \sum_{n=0}^{\frac{N}{2}-1} \left(x[n] - x \left[n + \frac{N}{2} \right] \right) W_N^n W_{\frac{N}{2}}^{rn}$$


This is a $\frac{N}{2}$ -point DFT of

$$\left(x[n] - x \left[n + \frac{N}{2} \right] \right) W_N^n \quad n = 0, 1, \dots, \frac{N}{2} - 1.$$

For $N = 8$



Decomposing $\frac{N}{2}$ -point DFT blocks down to 2-point DFTs we get the following
flow diagram



THE GOERTZEL ALGORITHM

The algorithm reduces the storage requirement

DFT expression:

$$X[k] = \sum_{n=0}^{N-1} x[n] W_N^{kn} \quad k = 0, 1, \dots, N - 1$$

can be written in a recursive form.

For example, let $N = 8$

$$X[k] = x[0] + x[1]W_N^k + x[2]W_N^{2k} + x[3]W_N^{3k} + x[4]W_N^{4k} + x[5]W_N^{5k} + x[6]W_N^{6k} + x[7]W_N^{7k}$$

$$= x[0] +$$

$$W_N^k(x[1] + x[2]W_N^k + x[3]W_N^{2k} + x[4]W_N^{3k} + x[5]W_N^{4k} +$$

$$x[6]W_N^{5k} + x[7]W_N^{6k})$$

\vdots

$$= x[0] +$$

$$W_N^k \left(x[1] +$$

$$W_N^k \left(x[2] +$$

$$W_N^k \left(x[3] + W_N^k \left(x[4] + W_N^k \left(x[5] + W_N^k (x[6] + x[7]W_N^k) \right) \right) \right) \right) \right)$$


OR

Since $W_N^{-kN} = 1$

$$\begin{aligned} X[k] &= W_N^{-kN} \sum_{n=0}^{N-1} x[n] W_N^{kn} \\ &= \sum_{n=0}^{N-1} x[n] W_N^{-k(N-n)} \end{aligned}$$

For $N = 8$

$$\begin{aligned} W_8^{-8k} X[k] &= \\ &\left(\left(\left(\left(\left(\left(\left(\left(x[0] W_8^{-k} + x[1] \right) W_8^{-k} + x[2] \right) W_8^{-k} + x[3] \right) W_8^{-k} + \right. \right. \right. \right. \right. \\ &\left. \left. \left. \left. \left. x[4] \right) W_8^{-k} + x[5] \right) W_8^{-k} + x[6] \right) W_8^{-k} + x[7] \right) \right) \end{aligned}$$



$$y_k[n] = W_N^{-k} y_k[n-1] + x[n] \quad n = 0, 1, \dots, N$$

$$y_k(-1) = 0 \quad x[N] = 0$$

Take $X[k] = y_k[N]$

Note that $X[k]$'s are computed sequentially.

Computational demand:

For each k , $y_k[1], y_k[2], \dots, y_k[N - 1]$ must be computed to find $y_k[N] = X[k]$.

This requires, (assuming a complex input) $4N$ multiplications and $4N$ additions.

This amount is almost the same as that of direct computation.

However, the storage requirement of W_N^{kn} values vanishes.

DOWNSAMPLING AND DECIMATION-IN-TIME

For the N -point DFT of $x[n]$ we obtained

$$X[k] = G[k] + W_N^k H[k]$$

where $G[k]$ and $H[k]$ are the $\frac{N}{2}$ -point DFTs, respectively, of even and odd indexed elements of the sequence $x[n]$, i.e.,

$$g[n] = x[2n]$$

$$h[n] = x_{-1}[2n] \quad x_{-1}[n] = x[n+1]$$

$$G(e^{j\omega}) = \frac{1}{2} \left(X(e^{j\frac{\omega}{2}}) + X(e^{j\frac{1}{2}(\omega-2\pi)}) \right)$$

$$H(e^{j\omega}) = \frac{1}{2} \left(e^{j\frac{\omega}{2}} X(e^{j\frac{\omega}{2}}) + e^{j\frac{1}{2}(\omega-2\pi)} X(e^{j\frac{1}{2}(\omega-2\pi)}) \right)$$

Therefore

$$X(e^{j\omega}) = G(e^{j2\omega}) + e^{-j\omega} H(e^{j2\omega})$$

$$X[k] = X(e^{j\omega}) \Big|_{\omega=k\frac{2\pi}{N}}$$

$$G(e^{j2\omega}) = \sum_{n=0}^{\frac{N}{2}-1} x[2n]e^{-j2\omega n}$$

$$G(e^{j2\omega}) \Big|_{\omega=k\frac{2\pi}{N}} = \sum_{n=0}^{\frac{N}{2}-1} x[2n]e^{-j2k\frac{2\pi}{N}n}$$

$$= \sum_{n=0}^{\frac{N}{2}-1} x[2n]W_{\frac{N}{2}}^{kn}$$

Likewise

$$H(e^{j2\omega}) \Big|_{\omega=k\frac{2\pi}{N}} = \sum_{n=0}^{\frac{N}{2}-1} x[2n+1]W_{\frac{N}{2}}^{kn}$$

RADIX 3 vs, RADIX 2

In place computations

Chirp transform 9.6