

2) a) $x[n] = \delta[n] + 3\delta[n-1] + \delta[n-2]$

$$X_3[k] = \sum_{n=0}^2 x[n] e^{-j\frac{2\pi}{3}kn} = 1 + 3e^{-j\frac{2\pi}{3}k} + e^{-j\frac{4\pi}{3}k}, \text{ for } k=0,1,2$$

0, elsewhere

$$X_5[k] = \sum_{n=0}^4 x[n] e^{-j\frac{2\pi}{5}kn} = 1 + 3e^{-j\frac{2\pi}{5}k} + e^{-j\frac{4\pi}{5}k}, \text{ for } k=0,1,2,3,4$$

0, elsewhere

b) $\tilde{X}_3[k] = \sum_{n=-\infty}^{\infty} X_3[k-3n] = X_3[(k)_3]$

$$\tilde{X}_5[k] = \sum_{n=-\infty}^{\infty} X_5[k-5n] = X_5[(k)_5]$$

c)
$$x[n] = \begin{cases} \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{j\frac{2\pi}{N}kn} & , n=0,1,2,\dots,N-1 \\ 0 & , \text{ else} \end{cases}$$

$$\Rightarrow x[n] = \frac{1}{3} \sum_{k=0}^2 X[k] e^{j\frac{2\pi}{3}kn}, \text{ for } n=0,1,2$$

0, else


$$\begin{aligned} x[0] &= \frac{1}{3} \sum_{k=0}^2 (1 + 3e^{-j\frac{2\pi}{3}k} + e^{-j\frac{4\pi}{3}k}) e^{j\frac{2\pi}{3}k \cdot 0} \\ &= \frac{1}{3} \sum_{k=0}^2 e^{j\frac{2\pi}{3}kn} + 3e^{j\frac{2\pi}{3}k(n-1)} + e^{j\frac{2\pi}{3}k(n-2)} \\ &= \frac{1}{3} [X_3[0] + X_3[1] e^{j\frac{2\pi}{3}n} + X_3[2] e^{j\frac{4\pi}{3}n}], \text{ for } n=0,1,2 \\ &\quad 0, \text{ else} \end{aligned}$$

in the same way:
$$x[n] = \frac{1}{5} [X_5[0] + X_5[1] e^{j\frac{2\pi}{5}n} + X_5[2] e^{j\frac{4\pi}{5}n} + X_5[3] e^{j\frac{6\pi}{5}n} + X_5[4] e^{j\frac{8\pi}{5}n}]$$

for $n=0,1,2,3,4$

d) Matlab

3) a)



b)

$$y[n] = x[n-2] \quad z[n] = x[(n+1)_5]$$

c)

$$y[k] = X[k] \cdot e^{-j\frac{4\pi}{5}k} \Rightarrow y[n] = x[n-2] = x[(n-2)_5] \text{ for this case}$$

$$z[k] = X[k] \cdot e^{j\frac{2\pi}{5}k}$$

$z[n]$ and $y[n]$ do not have 3-point DFTs since they are length of 5.

6)

$$x[n] = 3\delta[n] - 2\delta[n-1] + \delta[n-2] + \delta[n-3] - 2\delta[n-4] - \delta[n-5]$$

a)

$$W_3[k] = X(e^{j\omega}) \Big|_{\omega = \frac{2\pi k}{3}} \quad k=0,1,2 \Rightarrow W_3[k] \text{ is 3 point DFT of } x[n]$$

but $x[n]$ is of 6 in length.

\Rightarrow we find $w[n]$ by overlapping the remaining 3 value.

that is;

$$\left. \begin{aligned} w[0] &= x[0] + x[3] \\ w[1] &= x[1] + x[4] \\ w[2] &= x[2] + x[5] \\ w[n] &= 0 \text{ else} \end{aligned} \right\} w[n] = 4\delta[n] - 4\delta[n-1]$$

b)

in a similar way: $w[n] = 2\delta[n] - 2\delta[n-1] + \delta[n-2] + \delta[n-3] - 2\delta[n-4]$

c)

$$w[n] = x[n] = 3\delta[n] - 2\delta[n-1] + \delta[n-2] + \delta[n-3] - 2\delta[n-4] - \delta[n-5]$$

$$d) h[n] = 2\delta[n] - 8\delta[n-1]$$

$$y_3[k] = H_3[k]w_1[k] \Rightarrow y[n] = h[n] * w[n] \Rightarrow$$

$$\Rightarrow y[n] = 8\delta[n] - 12\delta[n-1] + 4\delta[n-2]$$

$$\begin{bmatrix} 4 & -4 & 0 \\ 2 & -1 & 0 \end{bmatrix} * \begin{bmatrix} 8 & -8 & 0 \\ 0 & -4 & 4 \\ 0 & 0 & 0 \end{bmatrix} \Rightarrow y[n] = \begin{bmatrix} 8 & -12 & 4 \end{bmatrix}$$

$$e) w[n] = [2 \ -2 \ 1 \ 1 \ -2] \\ h[n] = [2 \ -1 \ 0 \ 0 \ 0]$$

$$\begin{array}{rrrrr} 4 & -4 & 2 & 2 & -4 \\ + & 2 & -2 & 2 & -1 & -1 \\ \hline 6 & -6 & 4 & 1 & -5 \end{array}$$

$$\Rightarrow y[n] = 6\delta[n] - 6\delta[n-1] + 4\delta[n-2] + \delta[n-3] - 5\delta[n-4]$$

$$f) w[n] = [3 \ -2 \ 1 \ 1 \ -2 \ -1 \ 0 \ 0] \\ h[n] = [2 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0]$$

$$\begin{array}{rrrrrrrr} 6 & -4 & 2 & 2 & -4 & -2 & 0 & 0 \\ + & 0 & -3 & 2 & -1 & -1 & 2 & 1 & 0 \\ \hline 6 & -7 & 4 & 1 & -5 & 0 & 1 & 0 \end{array}$$

$$\Rightarrow y[n] = 6\delta[n] - 7\delta[n-1] + 4\delta[n-2] + \delta[n-3] - 5\delta[n-4] + \delta[n-5]$$

g) $x[n]$ is of length 6 and $h[n]$ is of length 2

\Rightarrow N -point circular convolution = linear convolution ; if $N \geq 7$

$$\Rightarrow y[n] = y_8[n] \neq y_3[n] \neq y_5[n]$$

$$7) a) X[k] = \sum_{n=0}^{N-1} x[n] W_N^{kn} ; k=0,1,2,\dots,N-1 \text{ where } W_N = e^{-j\frac{2\pi}{N}}$$

$$\begin{aligned} &= \sum_{n=\text{even}} x[n] W_N^{kn} + \sum_{n=\text{odd}} x[n] W_N^{kn} = \sum_{m=0}^{\frac{N}{2}-1} x[2m] W_N^{2mk} + \sum_{m=0}^{\frac{N}{2}-1} x[2m+1] W_N^{(2m+1)k} \\ &= \sum_{m=0}^{\frac{N}{2}-1} e[n] W_{\frac{N}{2}}^{mk} + W_N^k \sum_{m=0}^{\frac{N}{2}-1} o[n] W_{\frac{N}{2}}^{mk} \end{aligned}$$

here we used $W_N^{2km} = W_{\frac{N}{2}}^{km}$

$\frac{N}{2}$ point DFT of $e[n]$ $\frac{N}{2}$ point DFT of $o[n]$

$$\Rightarrow X[k] = E\left[\left(k\right)_{\frac{N}{2}}\right] + e^{-j\frac{2\pi}{N}k} O\left[\left(k\right)_{\frac{N}{2}}\right] ; \text{ for } k=0,1,2,\dots,N-1$$

b) i) $X[k] = \sum_{n=0}^{N-1} x[n] W_N^{kn} = \sum_{n=0}^{N-1} x[n] \left(\cos\left(\frac{2\pi}{N} kn\right) - j \sin\left(\frac{2\pi}{N} kn\right) \right)$ } here we are given that $x[n]$ is a real sequence

Direct computation involves $2N$ real multiplications and $2N-2$ real additions for each k .

\Rightarrow For all values of $X[k]$; direct computation involves $2N^2$ real multiplications and $2N^2-2N$ real additions.

ii) $X[k] = E\left[\left(k\right)_{\frac{N}{2}}\right] + W_N^k O\left[\left(k\right)_{\frac{N}{2}}\right]$ we know that $E[k] = E\left[k - \frac{N}{2}\right]$ } periodicity with $\frac{N}{2}$.
 $O[k] = O\left[k - \frac{N}{2}\right]$

and we also know that $W_N^{k-\frac{N}{2}} = -W_N^k$

\Rightarrow we can divide $X[k]$ into two as follows:

$X[k] = E[k] + W_N^k O[k]$ for $k=0, 1, 2, \dots, \frac{N}{2}-1$

$X\left[k + \frac{N}{2}\right] = E[k] - W_N^k O[k]$ for $k=0, 1, 2, \dots, \frac{N}{2}-1$

\Rightarrow We need to do $2\left(\frac{N}{2}\right)^2$ real multiplications and $2\left(\frac{N}{2}\right)^2 - 2\left(\frac{N}{2}\right)$ real additions for $E[k]$ and $O[k]$. Also $W_N^k \cdot O[k]$ involves $4\left(\frac{N}{2}\right)$ real multiplications and $2\left(\frac{N}{2}\right)$ real additions. To find $E[k] + W_N^k O[k]$, we also need to do $2\left(\frac{N}{2}\right)$ real additions. Method is called Split-Radix FFT algorithm

Computation involves total of $2\left(\frac{N}{2}\right)^2 + 2\left(\frac{N}{2}\right)^2 + 4\left(\frac{N}{2}\right) = N^2 + 4N$ real mults. and $N^2 + 2N$ real additions.

iii) Second computation method is better especially for large N 's (Includes less comps)

8) $h[n] = 2\delta[n] - \delta[n-1] + \delta[n-2]$ $x[n] = [1 \ 2 \ 3 \ 4 \ -1 \ -2 \ -3 \ -4 \ 1 \ 2 \ 3 \ 4]$ $0 \leq n < 11$

a) $P=3$ $L=4 \Rightarrow N=3+4-1=6$ -point DFT, we will use.

b) input is length of 12 \Rightarrow there are 3 input segments \Rightarrow $x_1 = [1 \ 2 \ 3 \ 4]$
 $x_2 = [-1 \ -2 \ -3 \ -4] = -x_1$
 $x_3 = [1 \ 2 \ 3 \ 4] = x_1$

c) $y_1 = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & -1 & 1 & 1 \end{bmatrix} \otimes = \begin{bmatrix} 2 & 3 & 5 & 7 & -1 & 4 \end{bmatrix}$
 $\Rightarrow y_2 = [-2 \ -3 \ -5 \ -7 \ 1 \ -4]$
 $y_3 = y_1$ (also verified with MATLAB)



d) $y = \begin{bmatrix} 2 & 3 & 5 & 7 & -1 & 4 \\ -2 & -3 & -5 & -7 & 1 & -4 \end{bmatrix}$
 $y[n] = [2 \ 3 \ 5 \ 7 \ -3 \ 1]$

APPENDIX:

2)d) MATLAB code for 3-point DFT:

```
Editor - D:\Lessons\4th Year\EE430\HW3\sor...
1 - x=[1 3 1];
2 - y=fft(x,3);
3 - ix=ifft(y);
4 - figure;
5 - stem(abs(y));
6 - figure;
7 - stem(phase(y)*180/pi);
8 -
9 -
```



Magnitude and phase(degree) of $X[k]$, respectively:



MATLAB code for 5-point DFT:

```
Editor - D:\Lessons\4th Year\EE430\HW3\sor...
1 - x=[1 3 1];
2 - y=fft(x,5);
3 - ix=ifft(y);
4 - figure;
5 - stem(abs(y));
6 - figure;
7 - stem(phase(y)*180/pi);
8 -
9 -
```

Magnitude and phase(degree) of $X[k]$, respectively:



8)c) MATLAB code for obtaining the outputs corresponding to each input segment:

```
Editor - D:\Lessons\4th Year\EE430\HW3\soru2d.m
1 - x1=[1 2 3 4];
2 - x2=[-1 -2 -3 -4];
3 - x3=x1;
4 - h=[2 -1 1];
5 - X1=fft(x1,6);
6 - X2=fft(x2,6);
7 - X3=fft(x3,6);
8 - H=fft(h,6);
9 - Y1=X1.*H;
10 - Y2=X2.*H;
11 - Y3=X3.*H;
12 - y1=ifft(Y1);
13 - y2=ifft(Y2);
14 - y3=ifft(Y3);
15 - stem(y1,'color','red');
16 - hold on;
17 - stem(y2,'color','blue');
18 - hold on;
19 - stem(y3,'color','green');
20
21
```

Outputs, y1 is red, y2 is green and y3 is blue: (since $y_1=y_3$ red line is not seen)

