

Formalization of Weingarten Calculus

Melody Lee

Yanting Teng

June 12, 2026

Chapter 1

Schur Weyl Duality

1.1 Operator Space

Suppose $\mathcal{X} = \mathbb{C}^{[d]}$ and $\mathcal{Y} = \mathbb{C}^{[d]}$ where $d \in \mathbb{N}$. Let the set of linear operators be $\mathcal{L}(\mathcal{X}, \mathcal{Y})$. When $\mathcal{Y} = \mathcal{X}$, let the set of linear operators be $\mathcal{L}(\mathcal{X})$.

We briefly restate some relevant definitions and properties of linear operators that we will use in the subsequent sections.

Remark 1.1.1. For any pair of operators $X, Y \in \mathcal{L}(\mathcal{X})$, the Lie bracket $[X, Y] \in \mathcal{L}(\mathcal{X})$ is defined as

$$[X, Y] = XY - YX, \quad (1.1)$$

such that $[X, Y] = 0$ if and only if X and Y commute.

Remark 1.1.2. Let $\mathcal{A} \subseteq \mathcal{L}(\mathcal{X})$ be a subalgebra of $\mathcal{L}(\mathcal{X})$. The commutant of $\mathcal{A}$ is given by

$$\text{Comm}(\mathcal{A}) = \{Y \in \mathcal{L}(\mathcal{X}) : [X, Y] = 0 \text{ for all } X \in \mathcal{A}\}. \quad (1.2)$$

Definition 1.1.3 (Diagonal operators). An operator $X \in \mathcal{L}(\mathcal{X})$ is a diagonal operator if $X_{a,b} = 0$ for all $a, b \in [d]$ where $a \neq b$. For a given vector $u \in \mathcal{X}$, we can define a diagonal operator $\text{Diag}(u) \in \mathcal{L}(\mathcal{X})$ to denote the diagonal operator whose diagonal entries are given by the entries of u . That is, we have

$$\text{Diag}(u)_{a,b} = \begin{cases} u_a & \text{if } a = b \\ 0 & \text{if } a \neq b \end{cases} \quad (1.3)$$

Remark 1.1.4 (Projection operators). Let Π denote the project operator onto a subspace $\mathcal{V}$. Then Π is a linear operator that satisfies the following properties:

- $\Pi^2 = \Pi$, the idempotent property; and
- $\Pi^* = \Pi$, the self-adjoint property.

Let the collection of all projection operators be denoted $\text{Proj}(\mathcal{X})$. There is a uniquely defined projection operator $\Pi \in \text{Proj}(\mathcal{X})$ such that its image $\text{Im}(\Pi) = \mathcal{V}$. When necessary, we denote this projection operator as $\Pi_{\mathcal{V}}$ to indicate the subspace $\mathcal{V}$ onto which it projects.

Remark 1.1.5 (Isometries). An operator $\mathcal{A} \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$ is an isometry if it satisfies the condition

$$\mathcal{A}^* \mathcal{A} = I_{\mathcal{X}}, \quad (1.4)$$

which is equivalent to preserving the Euclidean norm of vectors in $\mathcal{X}$, i.e., $\|\mathcal{A}|\psi\rangle\| = \| |\psi\rangle \|$ for all $|\psi\rangle \in \mathcal{X}$.

Remark 1.1.6 (Unitary operators). Denote the set of isometries that map $\mathcal{X}$ to itself as the set of unitary operators $U(\mathcal{X})$. Every unitary operator $U \in U(\mathcal{X})$ is necessarily invertible and satisfies the normal condition

$$UU^* = U^*U = I_{\mathcal{X}}. \quad (1.5)$$

1.2 Symmetric subspaces

Definition 1.2.1 (Symmetric subspace of linear operators). The symmetric subspace of linear operators $\mathcal{L}(\mathcal{X})^{\otimes k}$ is the set of all operators $X \in \mathcal{L}(\mathcal{X}^{\otimes k})$ that remain unchanged (invariant) under the permutation W_{π} . That is,

$$\mathcal{L}(\mathcal{X})^{\otimes k} = \{X \in \mathcal{L}(\mathcal{X}^{\otimes k}) : X = W_{\pi} X W_{\pi}^* \text{ for all } \pi \in S_k\}. \quad (1.6)$$

Remark 1.2.2. Let S_k denote the symmetric group. Then given $\pi \in S_k$, the permutation matrix (operator) $V_d(\pi)$ is the unitary matrix that satisfies

$$W_{\pi} |\psi_1\rangle \otimes \dots \otimes |\psi_k\rangle = |\psi_{\pi^{-1}(1)}\rangle \otimes \dots \otimes |\psi_{\pi^{-1}(k)}\rangle \quad (1.7)$$

for all $|\psi_1\rangle, \dots, |\psi_k\rangle \in \mathbb{C}^d$.

Definition 1.2.3 (Standard computational basis of linear operators). The standard computational basis of linear operators $\{E_{i,j}\}_{i,j \in [d]}$ is defined as the set of d^2 operators in $\mathcal{L}(\mathcal{X})$ such that

$$E_{i,j} = |i\rangle\langle j| \quad (1.8)$$

for all $i, j \in [d]$, where $\{|i\rangle\}_{i \in [d]}$ is the standard computational basis of $\mathcal{X}$.

Definition 1.2.4 (Operator-vector correspondence). The operator-vector correspondence is a bijective mapping between the set of linear operators $\mathcal{L}(\mathcal{X})$ and the tensor product space $\mathcal{X} \otimes \mathcal{X}$ defined as

$$\text{vec} : \mathcal{L}(\mathcal{X}) \rightarrow \mathcal{X} \otimes \mathcal{X}, \quad E_{i,j} \mapsto |i\rangle \otimes |j\rangle \text{ for all } i, j \in [d]. \quad (1.9)$$

One relevant identity we use is that for $A_0, A_1, B \in \mathcal{L}(\mathcal{X})$,

$$(A_0 \otimes A_1) \text{vec}(B) = \text{vec}(A_0 B A_1^T). \quad (1.10)$$

1.3 Permutation Invariance and Properties of Unitarily Invariant Measures

Theorem 1.3.1 (Tensor power span theorem). Let k be a positive integer, and let $X \in \mathbb{C}^{[d]}$. For any set $\mathcal{A} \subseteq \mathbb{C}$ satisfying $|\mathcal{A}| \geq k + 1$, it holds that

$$\text{Span}\{v^{\otimes k} : v \in \mathcal{A}^{[d]}\} = \mathcal{X}^{\otimes k}. \quad (1.11)$$

Proposition 1.3.2 (Symmetric subspace and span of tensor product space). Let $\mathcal{X}$ be a complex Euclidean space and k be a positive integer. Then the following statements are equivalent:

1. $X \in \mathcal{L}(\mathcal{X})^{\otimes k}$.

2. For $V \in U(\mathcal{X}^{\otimes k} \otimes \mathcal{X}^{\otimes k}, (\mathcal{X} \otimes \mathcal{X})^{\otimes k})$ being the isometry defined by

$$V \text{vec}(Y_1 \otimes \cdots \otimes Y_k) = \text{vec}(Y_1) \otimes \cdots \otimes \text{vec}(Y_k) \quad (1.12)$$

holding for all $Y_1, \dots, Y_k \in \mathcal{L}(\mathcal{X})$, one has that

$$V \text{vec}(X) \in (\mathcal{X} \otimes \mathcal{X})^{\otimes k}. \quad (1.13)$$

3. $X \in \text{Span}\{Y^{\otimes k} : Y \in \mathcal{L}(\mathcal{X})\}$

Theorem 1.3.3 (Equivalence of symmetric subspace and unitary tensor span). *Let $\mathcal{X}$ be a complex Euclidean space and k be a positive integer. Then the symmetric subspace of linear operators is equivalent to the span of the unitary tensor products, i.e.,*

$$\mathcal{L}(\mathcal{X})^{\otimes k} = \text{Span}\{U^{\otimes k} : U \in \mathcal{U}(\mathcal{X})\}. \quad (1.14)$$

Lemma 1.3.4 (Invariant subspace lemma). *Let $\mathcal{X}$ be a complex Euclidean space, let $\mathcal{V} \subseteq \mathcal{X}$ be a subspace, and let $A \in \mathcal{L}(\mathcal{X})$ be an operator. The following statements are equivalent:*

1. It holds that both $A\mathcal{V} \subseteq \mathcal{V}$ and $A^*\mathcal{V} \subseteq \mathcal{V}$.
2. It holds that $[A, \Pi_{\mathcal{V}}] = 0$.

Theorem 1.3.5 (Double Commutant Theorem). *Let $\mathcal{A}$ be a self-adjoint, unital subalgebra of $\mathcal{L}(\mathcal{X})$. Then the double commutant of $\mathcal{A}$ is equal to $\mathcal{A}$, i.e.,*

$$\text{Comm}(\text{Comm}(\mathcal{A})) = \mathcal{A}. \quad (1.15)$$

Proof. For all $X \in \text{Comm}(\mathcal{A})$ and any $A \in \mathcal{A}$, we have

$$[X, A] = AX - XA = 0. \quad (1.16)$$

Therefore, $A \in \text{Comm}(\text{Comm}(\mathcal{A}))$, so $\mathcal{A} \subseteq \text{Comm}(\text{Comm}(\mathcal{A}))$.

To demonstrate reverse inclusion, consider the subalgebra $\mathcal{B} \subseteq \mathcal{L}(\mathcal{X} \otimes \mathcal{X})$ defined as

$$\mathcal{B} = \{X \otimes \mathbb{I} : X \in \mathcal{A}\}. \quad (1.17)$$

For all $Y \in \mathcal{L}(\mathcal{X} \otimes \mathcal{X})$, it can be written in terms of the standard computational basis

$$Y = \sum_{i,j \in [d]} \sum_{a,b \in [d]} c_{i,j,a,b} |i\rangle\langle j| \otimes |a\rangle\langle b|. \quad (1.18)$$

Using the definition of the standard computational basis of operators, we can rearrange the right-hand side into the form

$$Y = \sum_{a,b \in [d]} Y_{a,b} \otimes E_{a,b} \quad (1.19)$$

where $Y_{a,b} = \sum_{i,j \in [d]} c_{i,j,a,b} |i\rangle\langle j|$.

Y commutes with $\mathcal{B}$ if and only if $Y(X \otimes \mathbb{I}) = (X \otimes \mathbb{I})Y$ for all $X \in \mathcal{A}$. Substituting Equation ?? and simplifying shows that Y commutes with $\mathcal{B}$ if and only if $[Y_{a,b}, X] = 0$. For any $X \in \text{Comm}(\text{Comm}(\mathcal{A}))$, X must commute with $\text{Comm}(\mathcal{A})$, and therefore

$$X \otimes \mathbb{I} \in \text{Comm}(\text{Comm}(\mathcal{B})). \quad (1.20)$$

It remains to show that every element in $\text{Comm}(\text{Comm}(\mathcal{B}))$ is of the form $X \otimes \mathbb{I}$. Define a subspace $\mathcal{V} \subseteq X \otimes X$, where

$$\mathcal{V} = \left\{ \text{vec}(X) = \sum_{a,b \in [d]} c_{a,b} |a\rangle \otimes |b\rangle : X \in \mathcal{A} \right\}. \quad (1.21)$$

For any arbitrary $X \in \mathcal{A}$, we find that

$$\begin{aligned} (X \otimes \mathbb{I})\mathcal{V} &= \{(X \otimes \mathbb{I})V : V \in \mathcal{V}\} \\ &= \left\{ (X \otimes \mathbb{I}) \left(\sum_{a,b \in [d]} c_{a,b} |a\rangle \otimes |b\rangle \right) : \sum_{a,b \in [d]} c_{a,b} |a\rangle \langle b| \in \mathcal{A} \right\} \\ &= \left\{ \sum_{a,b \in [d]} c_{a,b} X|a\rangle \otimes |b\rangle : \sum_{a,b \in [d]} c_{a,b} |a\rangle \langle b| \in \mathcal{A} \right\} \\ &\subseteq \mathcal{V}, \end{aligned} \quad (1.22)$$

the inclusion of which follows from the fact that $\mathcal{A}$ is an algebra so $X|a\rangle = Xe_a \in \mathcal{A}$. Similarly, since $\mathcal{A}$ is self-adjoint,

$$(X^* \otimes \mathbb{I})\mathcal{V} \subseteq \mathcal{V}. \quad (1.23)$$

By Lemma ??, this implies $[X \otimes \mathbb{I}, \Pi_{\mathcal{V}}] = 0$, and hence

$$\Pi_{\mathcal{V}} \in \text{Comm}(\mathcal{B}). \quad (1.24)$$

Therefore, for any arbitrary $X \in \text{Comm}(\text{Comm}(\mathcal{A}))$, Equation ?? implies

$$X \otimes \mathbb{I} \in \text{Comm}(\text{Comm}(\mathcal{B})). \quad (1.25)$$

It follows from Equation ?? that

$$[X \otimes \mathbb{I}, \Pi_{\mathcal{V}}] = 0. \quad (1.26)$$

By Lemma ??, this yields

$$(X \otimes \mathbb{I})\mathcal{V} \subseteq \mathcal{V}. \quad (1.27)$$

Since the subalgebra $\mathcal{A}$ is unital, $\mathbb{I} \in \mathcal{A}$ and thus $\text{vec}(\mathbb{I}) \in \mathcal{V}$. Therefore,

$$\begin{aligned} (X \otimes \mathbb{I})\text{vec}(\mathbb{I}) &= (X \otimes \mathbb{I}) \sum_{a \in [d]} |a\rangle \otimes |a\rangle \\ &= \sum_{a \in [d]} X|a\rangle \otimes |a\rangle = \sum_{a \in [d]} \left(\sum_{b,c \in [d]} X_{b,c} |b\rangle \langle c| \right) |a\rangle \otimes |a\rangle \\ &= \sum_{a \in [d]} \sum_{b \in [d]} X_{b,a} |b\rangle \otimes |a\rangle \\ &= \sum_{b,a \in [d]} X_{b,a} |b\rangle \otimes |a\rangle = \text{vec}(X) \in \mathcal{V}. \end{aligned} \quad (1.28)$$

By the bijectivity of vec , $X \in \mathcal{A}$, so $\text{Comm}(\text{Comm}(\mathcal{A})) \subseteq \mathcal{A}$. □

Theorem 1.3.6 (Schur-Weyl duality). *Let $\mathcal{X}$ be a complex Euclidean space, k be a positive integer, and $X \in \mathcal{L}(\mathcal{X}^{\otimes k})$ be an operator. Then the set of all commutators of the unitary group is equivalent to the span of the permutation operators, i.e.,*

$$\text{Comm}(\mathcal{U}(\mathcal{X}), k) = \text{Span}\{W_{\pi} : \pi \in S_k\} \quad (1.29)$$

Proof. Let $X \in \text{Comm}(\mathcal{U}(\mathcal{X}), k)$. then by the bilinearity of the Lie bracket,

$$X \in \text{Comm}(\text{Span}\{U^{\otimes k} : U \in \mathcal{U}(\mathcal{X})\}) \quad (1.30)$$

Proposition ?? implies that $X \in \text{Comm}(\mathcal{L}(\mathcal{X})^{\otimes k})$, so

$$\text{Comm}(\text{Span}\{U^{\otimes k} : U \in \mathcal{U}(\mathcal{X})\}) = \text{Comm}(\mathcal{L}(\mathcal{X})^{\otimes k}). \quad (1.31)$$

By Definition ??, $Z \in \mathcal{L}(\mathcal{X})^{\otimes k}$ if and only if there exists some $X \in \mathcal{L}(\mathcal{X})$ such that

$$Z = X = W_{\pi} X W_{\pi}^* \text{ for all } \pi \in S_k. \quad (1.32)$$

Right-multiplying and rearranging by W_{π} yields $X W_{\pi} - W_{\pi} X = 0$, and so by the bilinearity of the Lie bracket,

$$Z = X \in \text{Comm}(\text{Span}\{W_{\pi} : \pi \in S_k\}). \quad (1.33)$$

Hence

$$\mathcal{L}(\mathcal{X})^{\otimes k} = \text{Comm}(\text{Span}\{W_{\pi} : \pi \in S_k\}). \quad (1.34)$$

By Theorem ??, taking the commutant of both sides yields

$$\text{Comm}(\mathcal{L}(\mathcal{X})^{\otimes k}) = \text{Span}\{W_{\pi} : \pi \in S_k\}. \quad (1.35)$$

By substitution of Eq. ??, we have shown

$$\text{Comm}(\mathcal{U}(\mathcal{X}), k) = \text{Span}\{W_{\pi} : \pi \in S_k\}. \quad (1.36)$$

□

1.4 Examples

We illustrate some simple examples of Haar measure computations. In particular, we are interested in first-order computations on discrete and continuous groups. We approach this from the perspective of group averaging and twirling.

Definition 1.4.1 (Group average). Let G be a group of order N and R a representation of the group. Define the group average to be

$$\langle X \rangle_G = \frac{1}{N} \sum_g R(g) X R(g)^{\dagger}. \quad (1.37)$$

Remark 1.4.2 (Assumptions). Consider a n -qubit system. Let $d = 2^n$, and suppose we seek to compute moments of order k . We assume for now that $d > k$ for the examples below.

1.4.1 Pauli twirling

The end goal of this section is to prove the Irrep Group Averaging Corollary

$$\langle X \rangle_G = \frac{1}{d} \text{Tr}[X] I, \quad (1.38)$$

where $n_a = d$ is the dimension of the vector space of the representation. To illustrate this, we first consider the group average of the single-qubit Pauli group on some arbitrary initial state ρ .

Remark 1.4.3 (Pauli groups). We denote the single-qubit Pauli group as

$$G = \{\pm(i)\sigma_x, \pm(i)\sigma_y, \pm(i)\sigma_z, \pm(i)I\}, \quad (1.39)$$

where the Pauli matrices are given by

$$\sigma_0 = I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \sigma_x = X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \sigma_y = Y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \sigma_z = Z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \quad (1.40)$$

We note the following useful expressions.

Remark 1.4.4 (Bloch vectors). For any single-qubit quantum state ρ , we can write it in terms of its Bloch vector,

$$\rho = \frac{1}{2}(I + r \cdot \sigma), \quad (1.41)$$

where $r = (r_x, r_y, r_z) \in \mathbb{R}^3$ is the coordinate of the quantum state on the Bloch sphere and $\sigma = (\sigma_x, \sigma_y, \sigma_z)$ is a three-tuple of the Pauli operators.

Remark 1.4.5 (Pauli group identities). For all non-identity σ_i, σ_j in the Pauli group where $i \neq j$, the following three statements are true.

1. $\sigma_i \sigma_j \sigma_i = -\sigma_j$,
2. $\sigma_j^3 = \sigma_j$, and
3. $\sigma_i^2 = \sigma_i$, since σ_i is represented by a self-adjoint matrix.

We want to show the following simpler equivalence using matrix computations.

Theorem 1.4.6 (Pauli twirling of a single qubit). *For an arbitrary single-qubit state ρ ,*

$$\langle \rho \rangle_G = \frac{1}{N} \sum_g R(g) X R(g)^\dagger = \frac{I}{2}, \quad (1.42)$$

the maximally mixed state (since it is proportional to the identity matrix).

Proof. Note that G has order $N = 4$. For any Pauli operator $\pm(i)X \in G$, X is self-adjoint. Thus, we have that

$$\pm(i)X \rho (\pm(i)X)^\dagger = (\pm(i)(\mp i)X) \rho X = X \rho X. \quad (1.43)$$

By Def. ?? and Eq. ??,

$$\langle \rho \rangle_G = \frac{1}{4} (\sigma_x \rho \sigma_x + \sigma_y \rho \sigma_y + \sigma_z \rho \sigma_z + I \rho I). \quad (1.44)$$

By Remark ?? and Remark ??, we can rewrite ρ such that

$$\begin{aligned} \langle \rho \rangle_G &= \frac{1}{4} \left(\sigma_x \left(\frac{1}{2} (I + r \cdot \sigma) \right) \sigma_x + \sigma_y \left(\frac{1}{2} (I + r \cdot \sigma) \right) \sigma_y + \sigma_z \left(\frac{1}{2} (I + r \cdot \sigma) \right) \sigma_z + I \left(\frac{1}{2} (I + r \cdot \sigma) \right) I \right) \\ &= \frac{1}{4} \left(\frac{1}{2} [\sigma_x I \sigma_x + r_x \sigma_x^3 + r_y \sigma_x \sigma_y \sigma_x + r_z \sigma_x \sigma_z \sigma_x] + \frac{1}{2} [\sigma_y I \sigma_y + r_x \sigma_y \sigma_x \sigma_y + r_y \sigma_y^3 + r_z \sigma_y \sigma_z \sigma_y] \right. \\ &\quad \left. + \frac{1}{2} [\sigma_z I \sigma_z + r_x \sigma_z \sigma_x \sigma_z + r_y \sigma_z \sigma_y \sigma_z + r_z \sigma_z^3] + \frac{1}{2} [I^3 + r_x I \sigma_x I + r_y I \sigma_y I + r_z I \sigma_z I] \right) \\ &= \frac{1}{8} ([I + r_x \sigma_x - r_y \sigma_y - r_z \sigma_z] + [I - r_x \sigma_x + r_y \sigma_y - r_z \sigma_z] \\ &\quad + [I - r_x \sigma_x - r_y \sigma_y + r_z \sigma_z] + [I + r_x \sigma_x + r_y \sigma_y + r_z \sigma_z]) \\ &= \frac{1}{8} (4I) = \frac{I}{2}. \end{aligned} \quad (1.45)$$

□

1.5 Explicit Calculation of single qubit first moment

We consider systems of n qubits, represented by vectors of dimension $d = 2^n$. Consider the calculation of the first moment of a system with $n = 1$. Due to the small size of the system, we can explicitly define the Haar measure and parameterize the unitary in our calculations, which we do in the example below.

Remark 1.5.1 (Explicit construction of 2×2 unitaries). Every unitary U in the unitary group of size 2×2 , denoted $U(2)$, can be parameterized in terms of $\phi, \theta, \omega \in \mathbb{R}$, where

$$U = \begin{bmatrix} e^{-i(\phi+\omega)/2} \cos \frac{\theta}{2} & -e^{i(\phi-\omega)/2} \sin \frac{\theta}{2} \\ e^{-i(\phi-\omega)/2} \sin \frac{\theta}{2} & e^{i(\phi+\omega)/2} \cos \frac{\theta}{2} \end{bmatrix}. \quad (1.46)$$

The conjugate transpose $U^\dagger$ is given by

$$U^\dagger = \begin{bmatrix} e^{i(\phi+\omega)/2} \cos \frac{\theta}{2} & e^{i(\phi-\omega)/2} \sin \frac{\theta}{2} \\ -e^{-i(\phi-\omega)/2} \sin \frac{\theta}{2} & e^{-i(\phi+\omega)/2} \cos \frac{\theta}{2} \end{bmatrix}. \quad (1.47)$$

Let μ_H denote the generalized Haar measure on any n -qubit system. On a single qubit, we can write the moment calculation in terms of the expectation value, which in turn can be written as an integral under μ_H . The first moment of the operator of an n -qubit system O , where $O \in \mathcal{L}(\mathbb{C}^d)$, is

$$\mathbb{E}_{U \sim \mu_H}[UOU^\dagger] = \int_{U \sim \mu_H} UOU^\dagger d\mu_H. \quad (1.48)$$

We now define a Haar measure specific to the case $n = 1$.

Definition 1.5.2 (Parameterized single-qubit Haar measure). Let μ_2 denote the Haar measure on a single qubit, represented with vectors in $\mathbb{C}^2$. Define the single-qubit Haar measure in terms of the parameters given in Remark ??, where

$$d\mu_2 = \sin \theta d\theta \cdot d\omega \cdot d\phi \quad (1.49)$$

is the unnormalized Haar measure. The normalized computation of moments using the Haar measure is given by

$$\mathbb{E}_{U \sim \mu_2}[UOU^\dagger] = \frac{1}{8\pi^2} \int_{\phi=0}^{2\pi} \int_{\omega=0}^{2\pi} \int_{\theta=0}^{\pi} UOU^\dagger \sin \theta d\theta d\omega d\phi, \quad (1.50)$$

where the factor of $\frac{1}{8\pi^2}$ can be found by computing the integral and setting $UOU^\dagger = 1$.

Theorem 1.5.3 (Example: First moment of Pauli-Z operator). Consider $O = \sigma_z$, the Pauli-Z operator. We show that

$$\mathbb{E}_{U \sim \mu_2}[U\sigma_z U^\dagger] = \frac{\text{Tr}(\sigma_z)}{2} I. \quad (1.51)$$

Proof. The Pauli-Z operator is given by

$$\sigma_z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \quad (1.52)$$

Hence

$$\frac{\text{Tr}(\sigma_z)}{2} I = \frac{0}{2} I = \mathbf{0}, \quad (1.53)$$