Galantamine (Razadyne®)



Dementia Drug Education Handout

Why is this drug used?	Treatment of mild to moderate stages of Alzheimer's disease.
Typical dosing:	 Tablet or Solution, immediate release: Begin with 4 mg twice daily. May increase the dose in 4 week intervals if no significant side effects. 4 mg → 8 mg → 12 mg, twice daily (4-weeks = →) Capsule, extended release: Begin with 8 mg daily. May increase the dose in 4 week intervals if no significant side effects. 8 mg → 16 mg → 24 mg, once daily (4-weeks = →) If your drug is stopped for 3 or more days, restart at the lowest dose and increase dose every 4 weeks to current dose.
Available Formulations & Estimated Cost:	1. Tablet, immediate release (brand & generic): Strengths: 4 mg, 8 mg, 12 mg Cost (30 day supply): Generic \$190; Brand \$320 2. Capsule, extended release (brand & generic): Strengths: 8 mg, 16 mg, 24 mg Cost (30 day supply): Generic \$190; Brand \$320 3. Solution, immediate release (brand & generic): Strengths: 4 mg/mL Cost (30 day supply): Generic \$290; Brand \$350
Contraindications & Precautions for this drug:	Hypersensitivity to galantamine. May include: swelling, rash, hives.
Common Side Effects:	Dizziness Vomiting Shakiness Headache Diarrhea Slow heartbeat Upset stomach Weight loss

1. Take this drug at the same time each date	ay.
--	-----

2. Take it with food may minimize side effects.

Managing this medication:

3. Inform your health-care provider RIGHT AWAY if you experience: very bad dizziness, slow heartbeat, black or bloody stools, throwing up blood or throw up that looks like coffee grounds, seizures, very upset stomach, a big weight loss, shakiness, diarrhea or other side effects.



References:

- 1. Lexicomp galantamine, Razadyne®
- 2. Red Book Average Wholesale Price of Drugs 2013
- 3. Kirby Lee, PharmD, MA, MAS and Ron Finley, RPh, UCSF Memory and Aging Center