

Monads are monadic and locally presentable: a double perspective

T. Tsantilas

joint work with V. Aravantinos-Sotiropoulos and C. Vasilakopoulou

School of Applied Mathematical and Physical Sciences,
National Technical University of Athens

This work was implemented in the framework of H.F.R.I call "3rd Call for H.F.R.I.'s Research Projects to Support Faculty Members & Researchers" (H.F.R.I. Project Number: 23249)



H.F.R.I.
Hellenic Foundation for
Research & Innovation

Fibrant double categories

Definition

A **pseudo double category** $\mathbb{D}$ is a pseudo category object in $\mathbf{Cat}$.

Convention. All double categories will be pseudo.

Definition

A **pseudo double category** $\mathbb{D}$ is a pseudo category object in $\mathbf{Cat}$.

Convention. All double categories will be pseudo.

Notation. $\mathbb{D}$ consists of a *category* $\mathbb{D}_0$ of objects and tight arrows, a *category* $\mathbb{D}_1$ of loose arrows and cells

$$\begin{array}{ccccc}
 X & \xrightarrow{A} & Y & & X & \xrightarrow{A} & Y & & X & \xrightarrow{1_X} & X \\
 f \downarrow & f \alpha^g & \downarrow g & & \text{id}_X \downarrow & 1_A & \downarrow \text{id}_Y & & f \downarrow & 1_f & \downarrow f \\
 Z & \xrightarrow{B} & W, & & X & \xrightarrow{A} & Y, & & Y & \xrightarrow{1_Y} & Y
 \end{array}$$

source, target, identity and composition *functors*

$$\begin{array}{c}
 \xleftarrow{s} \\
 \mathbb{D}_0 \xrightarrow{1} \mathbb{D}_1 \xleftarrow{\odot} \mathbb{D}_1 \times_{\mathbb{D}_0} \mathbb{D}_1, \\
 \xleftarrow{t}
 \end{array}$$

and natural isomorphisms α, ℓ, r (associator, left and right unitors) subject to the usual axioms and coherence conditions.

Definition

A double category $\mathbb{D}$ is called **fibrant** if $\langle s, t \rangle: \mathbb{D}_1 \rightarrow \mathbb{D}_0 \times \mathbb{D}_0$ is a fibration.

Definition

A double category $\mathbb{D}$ is called **fibrant** if $\langle s, t \rangle : \mathbb{D}_1 \rightarrow \mathbb{D}_0 \times \mathbb{D}_0$ is a fibration.

In a fibrant double category, we can turn tight arrows to loose arrows (their liftings through the fibration) thus obtaining a rich interplay between $\mathbb{D}_0$ and $\mathbb{D}_1$.

Definition

A double category $\mathbb{D}$ is called **fibrant** if $\langle s, t \rangle: \mathbb{D}_1 \rightarrow \mathbb{D}_0 \times \mathbb{D}_0$ is a fibration.

In a fibrant double category, we can turn tight arrows to loose arrows (their liftings through the fibration) thus obtaining a rich interplay between $\mathbb{D}_0$ and $\mathbb{D}_1$.

Notation. For a tight arrow $f: X \rightarrow Y$ we denote by $\hat{f}: X \rightarrowtail Y$ and $\check{f}: Y \rightarrowtail X$ its **companion** and **conjoint** respectively with

$$\begin{array}{ccccc}
 X & \xrightarrow{\hat{f}} & Y & X & \xlongequal{\quad} & X & Y & \xrightarrow{\check{f}} & X & X & \xlongequal{\quad} & X \\
 f \downarrow & p_1 & \parallel & \parallel & p_2 & \downarrow f & \parallel & q_1 & \downarrow f & f \downarrow & q_2 & \parallel \\
 Y & \xlongequal{\quad} & Y & X & \xrightarrow[\hat{f}]{} & Y & Y & \xlongequal{\quad} & Y & Y & \xrightarrow[\check{f}]{} & X
 \end{array}$$

as the appropriate cells.

Examples. 1. The double category $\text{Span}(\mathcal{C})$ of spans in a category $\mathcal{C}$ with pullbacks. $\text{Span}(\mathcal{C})$ consists of the objects, arrows, spans and span homomorphisms of $\mathcal{C}$. Composition is given by pullbacks. For a tight $f : X \rightarrow Y$, $\hat{f}$ is $X \xrightarrow{\text{id}_X} X \xrightarrow{f} Y$ and $\check{f}$ is $Y \xleftarrow{f} X \xrightarrow{\text{id}_X} X$.

Examples. 1. The double category $\text{Span}(\mathcal{C})$ of spans in a category $\mathcal{C}$ with pullbacks. $\text{Span}(\mathcal{C})$ consists of the objects, arrows, spans and span homomorphisms of $\mathcal{C}$. Composition is given by pullbacks. For a tight $f: X \rightarrow Y$, $\hat{f}$ is $X \xrightarrow{\text{id}_X} X \xrightarrow{f} Y$ and $\check{f}$ is $Y \xleftarrow{f} X \xrightarrow{\text{id}_X} X$.

2. The double category $\mathbb{R}\text{el}(\text{Set})$ of sets, functions, relations and inclusion of relations. Composition is composition of relations. For a function $f: X \rightarrow Y$, $\hat{f}$ is the graph of f and $\check{f}$ is the opposite relation f^o .

Examples. 1. The double category $\text{Span}(\mathcal{C})$ of spans in a category $\mathcal{C}$ with pullbacks. $\text{Span}(\mathcal{C})$ consists of the objects, arrows, spans and span homomorphisms of $\mathcal{C}$. Composition is given by pullbacks. For a tight $f: X \rightarrow Y$, $\hat{f}$ is $X \xrightarrow{\text{id}_X} X \xrightarrow{f} Y$ and $\check{f}$ is $Y \xleftarrow{f} X \xrightarrow{\text{id}_X} X$.

2. The double category $\mathbb{R}\text{el}(\text{Set})$ of sets, functions, relations and inclusion of relations. Composition is composition of relations. For a function $f: X \rightarrow Y$, $\hat{f}$ is the graph of f and $\check{f}$ is the opposite relation f^o .

3. The double category $\mathbb{R}\text{ing}$ of rings, ring homomorphisms, bimodules and bimodule homomorphisms. Composition is given by the tensor product. For a ring homomorphism $f: R \rightarrow S$, $\hat{f}$ is ${}_R S_S$ and $\check{f}$ is ${}_S S_R$.

Examples. 1. The double category $\text{Span}(\mathcal{C})$ of spans in a category $\mathcal{C}$ with pullbacks. $\text{Span}(\mathcal{C})$ consists of the objects, arrows, spans and span homomorphisms of $\mathcal{C}$. Composition is given by pullbacks. For a tight $f : X \rightarrow Y$, $\hat{f}$ is $X \xrightarrow{\text{id}_X} X \xrightarrow{f} Y$ and $\check{f}$ is $Y \xleftarrow{f} X \xrightarrow{\text{id}_X} X$.

2. The double category $\text{Rel}(\text{Set})$ of sets, functions, relations and inclusion of relations. Composition is composition of relations. For a function $f : X \rightarrow Y$, $\hat{f}$ is the graph of f and $\check{f}$ is the opposite relation f° .

3. The double category Ring of rings, ring homomorphisms, bimodules and bimodule homomorphisms. Composition is given by the tensor product. For a ring homomorphism $f : R \rightarrow S$, $\hat{f}$ is ${}_R S_S$ and $\check{f}$ is ${}_S S_R$.

4. For a monoidal $\mathcal{V}$ with small coproducts preserved by $\otimes$ in each variable, $\mathcal{V}\text{-Mat}$ is the double category of sets, functions, profunctors $Y^{\text{op}} \times X \rightarrow \mathcal{V}$ with X, Y sets (called $\mathcal{V}$ -**matrices**) and natural transformations. Composition of $T : Y \rightarrowtail Z$ and $S : X \rightarrowtail Y$ is given by $\left\{ \sum_{y \in Y} T_{z,y} \otimes S_{y,x} \right\}_{z,x}$. For any function $f : X \rightarrow Y$, $\hat{f}$ and $\check{f}$ are the family $\{I \text{ if } f(x) = y \text{ and } 0 \text{ otherwise}\}_{y,x}$.

Definition [ASV]

A *fibrant* double category will be called **locally presentable** if

- $\mathbb{D}_0, \mathbb{D}_1$ are locally presentable,
- s, t are cocontinuous right adjoints and
- for every loose arrow A , the functors $- \odot A$ and $A \odot -$ are accessible.

Definition [ASV]

A *fibrant* double category will be called **locally presentable** if

- $\mathbb{D}_0, \mathbb{D}_1$ are locally presentable,
- $\mathfrak{s}, \mathfrak{t}$ are cocontinuous right adjoints and
- for every loose arrow A , the functors $- \odot A$ and $A \odot -$ are accessible.

Examples [ASV]

1. $\text{Span}(\mathcal{C})$ is locally presentable when $\mathcal{C}$ is locally presentable.

Definition [ASV]

A *fibrant* double category will be called **locally presentable** if

- $\mathbb{D}_0, \mathbb{D}_1$ are locally presentable,
- $\mathfrak{s}, \mathfrak{t}$ are cocontinuous right adjoints and
- for every loose arrow A , the functors $- \odot A$ and $A \odot -$ are accessible.

Examples [ASV]

1. $\text{Span}(\mathcal{C})$ is locally presentable when $\mathcal{C}$ is locally presentable.
2. $\text{Rel}(\text{Set})$ is locally presentable.

Definition [ASV]

A *fibrant* double category will be called **locally presentable** if

- $\mathbb{D}_0, \mathbb{D}_1$ are locally presentable,
- $\mathfrak{s}, \mathfrak{t}$ are cocontinuous right adjoints and
- for every loose arrow A , the functors $- \odot A$ and $A \odot -$ are accessible.

Examples [ASV]

1. $\text{Span}(\mathcal{C})$ is locally presentable when $\mathcal{C}$ is locally presentable.
2. $\text{Rel}(\text{Set})$ is locally presentable.
3. $\mathcal{V}\text{-Mat}$ is locally presentable when $\mathcal{V}$ is locally presentable and $\otimes$ preserves filtered colimits in each variable.

(co)Monads

Fix a double category $\mathbb{D}$. A **monad** in $\mathbb{D}$ is an loose endo-arrow $A_X : X \multimap X$ equipped with *globular multiplication* and **unit** cells

$$\begin{array}{ccc}
 X & \xrightarrow{A} & X & \xrightarrow{A} & X & & X & \xrightarrow{1_X} & X \\
 \parallel & & & & \parallel & & \parallel & & \parallel \\
 & & \Downarrow \mu_A & & & & \Downarrow \eta_A & & \\
 X & \xrightarrow{\quad A \quad} & X & & X & \xrightarrow{\quad A \quad} & X
 \end{array}$$

satisfying the usual associativity and unit laws.

Fix a double category $\mathbb{D}$. A **monad** in $\mathbb{D}$ is an loose endo-arrow $A_X : X \multimap X$ equipped with *globular multiplication* and **unit** cells

$$\begin{array}{ccc} X & \xrightarrow{A} X & \xrightarrow{A} X \\ \parallel & \Downarrow \mu_A & \parallel \\ X & \xrightarrow{A} X & \end{array} \quad \begin{array}{ccc} X & \xrightarrow{1_X} X & \\ \parallel & \Downarrow \eta_A & \parallel \\ X & \xrightarrow{A} X & \end{array}$$

satisfying the usual associativity and unit laws. Given two monads A_X, B_Y , a **monad morphism** is a cell ${}^f\alpha^f \equiv \alpha_f : A_X \Rightarrow B_Y$ that respects the multiplication and the units:

$$\begin{array}{ccccc} \begin{array}{ccccc} X & \xrightarrow{A} X & \xrightarrow{A} X \\ f \downarrow & \alpha & f \downarrow & \alpha & \downarrow f \\ Y & \xrightarrow{B} Y & \xrightarrow{B} Y \\ \parallel & \Downarrow \mu_B & \parallel \\ Y & \xrightarrow{B} Y & \end{array} & = & \begin{array}{ccccc} X & \xrightarrow{A} X & \xrightarrow{A} X \\ \parallel & \Downarrow \mu_A & \parallel \\ X & \xrightarrow{A} X & \\ f \downarrow & \alpha & \downarrow f \\ Y & \xrightarrow{B} Y & \end{array} & & \begin{array}{ccccc} X & \xrightarrow{1_X} X & \\ \parallel & \Downarrow \eta_A & \parallel \\ X & \xrightarrow{A} X & \\ f \downarrow & \alpha & \downarrow f \\ Y & \xrightarrow{B} Y & \end{array} & = & \begin{array}{ccccc} X & \xrightarrow{1_X} X & \\ f \downarrow & 1_f & \downarrow f \\ Y & \xrightarrow{1_Y} Y & \\ \parallel & \Downarrow \eta_B & \parallel \\ Y & \xrightarrow{B} Y & \end{array} \end{array}$$

Fix a double category $\mathbb{D}$. A **monad** in $\mathbb{D}$ is an loose endo-arrow $A_X : X \multimap X$ equipped with *globular multiplication* and **unit** cells

$$\begin{array}{ccc} X & \xrightarrow{A} X & \xrightarrow{A} X \\ \parallel & \Downarrow \mu_A & \parallel \\ X & \xrightarrow{A} X & \end{array} \quad \begin{array}{ccc} X & \xrightarrow{1_X} X & \\ \parallel & \Downarrow \eta_A & \parallel \\ X & \xrightarrow{A} X & \end{array}$$

satisfying the usual associativity and unit laws. Given two monads A_X, B_Y , a **monad morphism** is a cell ${}^f\alpha^f \equiv \alpha_f : A_X \Rightarrow B_Y$ that respects the multiplication and the units:

$$\begin{array}{ccccc} \begin{array}{ccccc} X & \xrightarrow{A} X & \xrightarrow{A} X \\ f \downarrow & \alpha & f \downarrow & \alpha & \downarrow f \\ Y & \xrightarrow{B} Y & \xrightarrow{B} Y \\ \parallel & \Downarrow \mu_B & \parallel \\ Y & \xrightarrow{B} Y \end{array} & = & \begin{array}{ccccc} X & \xrightarrow{A} X & \xrightarrow{A} X \\ \parallel & \Downarrow \mu_A & \parallel \\ X & \xrightarrow{A} X \\ f \downarrow & \alpha & \downarrow f \\ Y & \xrightarrow{B} Y \end{array} & & \begin{array}{ccccc} X & \xrightarrow{1_X} X & \\ \parallel & \Downarrow \eta_A & \parallel \\ X & \xrightarrow{A} X \\ f \downarrow & \alpha & \downarrow f \\ Y & \xrightarrow{B} Y \end{array} & = & \begin{array}{ccccc} X & \xrightarrow{1_X} X & \\ f \downarrow & 1_f & \downarrow f \\ Y & \xrightarrow{1_Y} Y \\ \parallel & \Downarrow \eta_B & \parallel \\ Y & \xrightarrow{B} Y \end{array} \end{array}$$

Dually, we can define comonads and comonad morphisms.

Monads and monad morphisms assemble into a category, $\mathbf{Mnd}(\mathbb{D})$. Dually, comonads and comonad morphisms assemble into the category $\mathbf{Cmd}(\mathbb{D})$.

There are obvious forgetful functors

$$\mathbf{Mnd}(\mathbb{D}) \rightarrow \mathbf{End}(\mathbb{D}) \quad \text{and} \quad \mathbf{Cmd}(\mathbb{D}) \rightarrow \mathbf{End}(\mathbb{D}).$$

Then, under certain assumptions on $\mathbb{D}$,

- there is a (co)free (co)monad,
- (co)monads are (co)monadic over the loose endo-arrows as soon as the free (co)monad exists, and
- the category of (co)monads is locally presentable.

Examples. 1. $\mathbf{Mnd}(\mathbf{Span}(\mathcal{C})) = \mathbf{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0 \quad \rightsquigarrow \quad \text{source and target } X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\Leftrightarrow$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\Leftrightarrow$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$

Examples. 1. $\mathbf{Mnd}(\mathbf{Span}(\mathcal{C})) = \mathbf{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightarrowtail X_0$	$\Leftrightarrow$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\Leftrightarrow$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\Leftrightarrow$	identity $X_0 \rightarrow X_1$

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\Leftrightarrow$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\Leftrightarrow$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\Leftrightarrow$	identity $X_0 \rightarrow X_1$
monad morphism	$\Leftrightarrow$	internal functor

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\Leftrightarrow$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\Leftrightarrow$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\Leftrightarrow$	identity $X_0 \rightarrow X_1$
monad morphism	$\Leftrightarrow$	internal functor

On the other hand, $\text{Cmd}(\text{Span}(\mathcal{C}))$ are just arrows in $\mathcal{C}$.

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\Leftrightarrow$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\Leftrightarrow$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\Leftrightarrow$	identity $X_0 \rightarrow X_1$
monad morphism	$\Leftrightarrow$	internal functor

On the other hand, $\text{Cmd}(\text{Span}(\mathcal{C}))$ are just arrows in $\mathcal{C}$.

$$\text{loose endo-arrow } A_X : X \rightharpoonup X \quad \Leftrightarrow \quad \text{span } X \xleftarrow{f} Y \xrightarrow{g} X$$

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\iff$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\iff$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\iff$	identity $X_0 \rightarrow X_1$
monad morphism	$\iff$	internal functor

On the other hand, $\text{Cmd}(\text{Span}(\mathcal{C}))$ are just arrows in $\mathcal{C}$.

loose endo-arrow $A_X : X \rightharpoonup X$	$\iff$	span $X \xleftarrow{f} Y \xrightarrow{g} X$
counit $\varepsilon_A : A \Rightarrow 1_A$	$\iff$	$f = g = \varepsilon_A$

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\iff$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\iff$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\iff$	identity $X_0 \rightarrow X_1$
monad morphism	$\iff$	internal functor

On the other hand, $\text{Cmd}(\text{Span}(\mathcal{C}))$ are just arrows in $\mathcal{C}$.

loose endo-arrow $A_X : X \rightharpoonup X$	$\iff$	span $X \xleftarrow{f} Y \xrightarrow{g} X$
counit $\varepsilon_A : A \Rightarrow 1_A$	$\iff$	$f = g = \varepsilon_A$

2. $\text{Mnd}(\text{Rel}(\text{Set})) = \text{Preord}(\text{Set})$, the category of reflexive and transitive relations.

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\iff$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\iff$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\iff$	identity $X_0 \rightarrow X_1$
monad morphism	$\iff$	internal functor

On the other hand, $\text{Cmd}(\text{Span}(\mathcal{C}))$ are just arrows in $\mathcal{C}$.

loose endo-arrow $A_X : X \rightharpoonup X$	$\iff$	span $X \xleftarrow{f} Y \xrightarrow{g} X$
counit $\varepsilon_A : A \Rightarrow 1_A$	$\iff$	$f = g = \varepsilon_A$

2. $\text{Mnd}(\text{Rel}(\text{Set})) = \text{Preord}(\text{Set})$, the category of reflexive and transitive relations.

loose endo-arrow $R_X : X \rightharpoonup X$	$\iff$	relation $R \subseteq X \times X$
--	--------	-----------------------------------

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\iff$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\iff$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\iff$	identity $X_0 \rightarrow X_1$
monad morphism	$\iff$	internal functor

On the other hand, $\text{Cmd}(\text{Span}(\mathcal{C}))$ are just arrows in $\mathcal{C}$.

loose endo-arrow $A_X : X \rightharpoonup X$	$\iff$	span $X \xleftarrow{f} Y \xrightarrow{g} X$
counit $\varepsilon_A : A \Rightarrow 1_A$	$\iff$	$f = g = \varepsilon_A$

2. $\text{Mnd}(\text{Rel}(\text{Set})) = \text{Preord}(\text{Set})$, the category of reflexive and transitive relations.

loose endo-arrow $R_X : X \rightharpoonup X$	$\iff$	relation $R \subseteq X \times X$
multiplication $\mu_R : R \odot R \Rightarrow R$	$\iff$	$RR \subseteq R \iff R$ is transitive

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\Leftrightarrow$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\Leftrightarrow$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\Leftrightarrow$	identity $X_0 \rightarrow X_1$
monad morphism	$\Leftrightarrow$	internal functor

On the other hand, $\text{Cmd}(\text{Span}(\mathcal{C}))$ are just arrows in $\mathcal{C}$.

loose endo-arrow $A_X : X \rightharpoonup X$	$\Leftrightarrow$	span $X \xleftarrow{f} Y \xrightarrow{g} X$
counit $\varepsilon_A : A \Rightarrow 1_A$	$\Leftrightarrow$	$f = g = \varepsilon_A$

2. $\text{Mnd}(\text{Rel}(\text{Set})) = \text{Preord}(\text{Set})$, the category of reflexive and transitive relations.

loose endo-arrow $R_X : X \rightharpoonup X$	$\Leftrightarrow$	relation $R \subseteq X \times X$
multiplication $\mu_R : R \odot R \Rightarrow R$	$\Leftrightarrow$	$RR \subseteq R \Leftrightarrow R$ is transitive
unit $\eta_R : 1_R \Rightarrow R$	$\Leftrightarrow$	$\Delta_X \subseteq R \Leftrightarrow R$ is reflexive

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\iff$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\iff$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\iff$	identity $X_0 \rightarrow X_1$
monad morphism	$\iff$	internal functor

On the other hand, $\text{Cmd}(\text{Span}(\mathcal{C}))$ are just arrows in $\mathcal{C}$.

loose endo-arrow $A_X : X \rightharpoonup X$	$\iff$	span $X \xleftarrow{f} Y \xrightarrow{g} X$
counit $\varepsilon_A : A \Rightarrow 1_A$	$\iff$	$f = g = \varepsilon_A$

2. $\text{Mnd}(\text{Rel}(\text{Set})) = \text{Preord}(\text{Set})$, the category of reflexive and transitive relations.

loose endo-arrow $R_X : X \rightharpoonup X$	$\iff$	relation $R \subseteq X \times X$
multiplication $\mu_R : R \odot R \Rightarrow R$	$\iff$	$RR \subseteq R \iff R$ is transitive
unit $\eta_R : 1_R \Rightarrow R$	$\iff$	$\Delta_X \subseteq R \iff R$ is reflexive
monad morphism	$\iff$	order preserving morphism

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\Leftrightarrow$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\Leftrightarrow$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\Leftrightarrow$	identity $X_0 \rightarrow X_1$
monad morphism	$\Leftrightarrow$	internal functor

On the other hand, $\text{Cmd}(\text{Span}(\mathcal{C}))$ are just arrows in $\mathcal{C}$.

loose endo-arrow $A_X : X \rightharpoonup X$	$\Leftrightarrow$	span $X \xleftarrow{f} Y \xrightarrow{g} X$
counit $\varepsilon_A : A \Rightarrow 1_A$	$\Leftrightarrow$	$f = g = \varepsilon_A$

2. $\text{Mnd}(\text{Rel}(\text{Set})) = \text{Preord}(\text{Set})$, the category of reflexive and transitive relations.

loose endo-arrow $R_X : X \rightharpoonup X$	$\Leftrightarrow$	relation $R \subseteq X \times X$
multiplication $\mu_R : R \odot R \Rightarrow R$	$\Leftrightarrow$	$RR \subseteq R \Leftrightarrow R$ is transitive
unit $\eta_R : 1_R \Rightarrow R$	$\Leftrightarrow$	$\Delta_X \subseteq R \Leftrightarrow R$ is reflexive
monad morphism	$\Leftrightarrow$	order preserving morphism

A comonad $C_X : X \rightarrow X$ in $\text{Cmd}(\text{Rel}(\text{Set}))$ is just a subobject of X .

Examples. 1. $\text{Mnd}(\text{Span}(\mathcal{C})) = \text{Cat}(\mathcal{C})$, the category of categories internal to $\mathcal{C}$.

loose endo-arrow $A_{X_0} : X_0 \rightharpoonup X_0$	$\iff$	source and target $X_0 \xleftarrow{s} X_1 \xrightarrow{t} X_0$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\iff$	composition $X_1 \times_{X_0} X_1 \rightarrow X_1$
unit $\eta_A : 1_A \Rightarrow A$	$\iff$	identity $X_0 \rightarrow X_1$
monad morphism	$\iff$	internal functor

On the other hand, $\text{Cmd}(\text{Span}(\mathcal{C}))$ are just arrows in $\mathcal{C}$.

loose endo-arrow $A_X : X \rightharpoonup X$	$\iff$	span $X \xleftarrow{f} Y \xrightarrow{g} X$
counit $\varepsilon_A : A \Rightarrow 1_A$	$\iff$	$f = g = \varepsilon_A$

2. $\text{Mnd}(\text{Rel}(\text{Set})) = \text{Preord}(\text{Set})$, the category of reflexive and transitive relations.

loose endo-arrow $R_X : X \rightharpoonup X$	$\iff$	relation $R \subseteq X \times X$
multiplication $\mu_R : R \odot R \Rightarrow R$	$\iff$	$RR \subseteq R \iff R$ is transitive
unit $\eta_R : 1_R \Rightarrow R$	$\iff$	$\Delta_X \subseteq R \iff R$ is reflexive
monad morphism	$\iff$	order preserving morphism

A comonad $C_X : X \rightarrow X$ in $\text{Cmd}(\text{Rel}(\text{Set}))$ is just a subobject of X .

$$\text{counit } \varepsilon_C : C \Rightarrow 1_C \iff C \subseteq \Delta_X$$

3. $\text{Mnd}(\mathbb{R}\text{ing}) = \text{Alg}$, the global category of algebras over rings.

3. $\text{Mnd}(\mathbb{R}\text{ing}) = \text{Alg}$, the global category of algebras over rings.

loose endo-arrow $M_R : R \rightarrowtail R \quad \Leftrightarrow \quad (R, R)\text{-bimodule } M$

3. $\mathbf{Mnd}(\mathbf{Ring}) = \mathbf{Alg}$, the global category of algebras over rings.

loose endo-arrow $M_R : R \rightarrowtail R$	$\Leftrightarrow$	(R, R) -bimodule M
multiplication $\mu_M : M \odot M \Rightarrow M$	$\Leftrightarrow$	multiplication $M \otimes_R M \rightarrow M$

3. $\mathbf{Mnd}(\mathbf{Ring}) = \mathbf{Alg}$, the global category of algebras over rings.

loose endo-arrow $M_R : R \rightarrowtail R$	$\Leftrightarrow$	(R, R) -bimodule M
multiplication $\mu_M : M \odot M \Rightarrow M$	$\Leftrightarrow$	multiplication $M \otimes_R M \rightarrow M$
unit $\eta_M : 1_M \Rightarrow M$	$\Leftrightarrow$	unit $R \rightarrow M$

3. $\mathbf{Mnd}(\mathbf{Ring}) = \mathbf{Alg}$, the global category of algebras over rings.

loose endo-arrow $M_R : R \rightarrowtail R$	$\Leftrightarrow$	(R, R) -bimodule M
multiplication $\mu_M : M \odot M \Rightarrow M$	$\Leftrightarrow$	multiplication $M \otimes_R M \rightarrow M$
unit $\eta_M : 1_M \Rightarrow M$	$\Leftrightarrow$	unit $R \rightarrow M$
monad morphism	$\Leftrightarrow$	algebra homomorphism

3. $\mathbf{Mnd}(\mathbf{Ring}) = \mathbf{Alg}$, the global category of algebras over rings.

loose endo-arrow $M_R : R \rightarrowtail R$	$\Leftrightarrow$	(R, R) -bimodule M
multiplication $\mu_M : M \odot M \Rightarrow M$	$\Leftrightarrow$	multiplication $M \otimes_R M \rightarrow M$
unit $\eta_M : 1_M \Rightarrow M$	$\Leftrightarrow$	unit $R \rightarrow M$
monad morphism	$\Leftrightarrow$	algebra homomorphism

Dually, $\mathbf{Cmd}(\mathbf{Ring})$ is the global category $\mathbf{Coalg}$ of coalgebras over rings.

3. $\mathbf{Mnd}(\mathbf{Ring}) = \mathbf{Alg}$, the global category of algebras over rings.

loose endo-arrow $M_R : R \rightharpoonup R$	$\Leftrightarrow$	(R, R) -bimodule M
multiplication $\mu_M : M \odot M \Rightarrow M$	$\Leftrightarrow$	multiplication $M \otimes_R M \rightarrow M$
unit $\eta_M : 1_M \Rightarrow M$	$\Leftrightarrow$	unit $R \rightarrow M$
monad morphism	$\Leftrightarrow$	algebra homomorphism

Dually, $\mathbf{Cmd}(\mathbf{Ring})$ is the global category $\mathbf{Coalg}$ of coalgebras over rings.

4. $\mathbf{Mnd}(\mathcal{V}\text{-}\mathbf{Mat}) = \mathcal{V}\text{-}\mathbf{Cat}$.

3. $\text{Mnd}(\mathbb{R}\text{ing}) = \text{Alg}$, the global category of algebras over rings.

loose endo-arrow $M_R : R \rightarrowtail R$	$\Leftrightarrow$	(R, R) -bimodule M
multiplication $\mu_M : M \odot M \Rightarrow M$	$\Leftrightarrow$	multiplication $M \otimes_R M \rightarrow M$
unit $\eta_M : 1_M \Rightarrow M$	$\Leftrightarrow$	unit $R \rightarrow M$
monad morphism	$\Leftrightarrow$	algebra homomorphism

Dually, $\text{Cmd}(\mathbb{R}\text{ing})$ is the global category Coalg of coalgebras over rings.

4. $\text{Mnd}(\mathcal{V}\text{-Mat}) = \mathcal{V}\text{-Cat}$.

loose endo-arrow $A_X : X \rightarrowtail X$	$\Leftrightarrow$	family of objects $A_{x,x}$ in $\mathcal{V}$
--	-------------------	--

3. $\text{Mnd}(\mathbb{R}\text{ing}) = \text{Alg}$, the global category of algebras over rings.

loose endo-arrow $M_R : R \rightarrowtail R$	$\Leftrightarrow$	(R, R) -bimodule M
multiplication $\mu_M : M \odot M \Rightarrow M$	$\Leftrightarrow$	multiplication $M \otimes_R M \rightarrow M$
unit $\eta_M : 1_M \Rightarrow M$	$\Leftrightarrow$	unit $R \rightarrow M$
monad morphism	$\Leftrightarrow$	algebra homomorphism

Dually, $\text{Cmd}(\mathbb{R}\text{ing})$ is the global category Coalg of coalgebras over rings.

4. $\text{Mnd}(\mathcal{V}\text{-Mat}) = \mathcal{V}\text{-Cat}$.

loose endo-arrow $A_X : X \rightarrowtail X$	$\Leftrightarrow$	family of objects $A_{x,x}$ in $\mathcal{V}$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\Leftrightarrow$	morphisms $A_{x',x''} \otimes A_{x,x'} \rightarrow A_{x,x''}$

3. $\mathbf{Mnd}(\mathbf{Ring}) = \mathbf{Alg}$, the global category of algebras over rings.

loose endo-arrow $M_R : R \rightarrowtail R$	$\iff$	(R, R) -bimodule M
multiplication $\mu_M : M \odot M \Rightarrow M$	$\iff$	multiplication $M \otimes_R M \rightarrow M$
unit $\eta_M : 1_M \Rightarrow M$	$\iff$	unit $R \rightarrow M$
monad morphism	$\iff$	algebra homomorphism

Dually, $\mathbf{Cmd}(\mathbf{Ring})$ is the global category $\mathbf{Coalg}$ of coalgebras over rings.

4. $\mathbf{Mnd}(\mathcal{V}\text{-}\mathbf{Mat}) = \mathcal{V}\text{-}\mathbf{Cat}$.

loose endo-arrow $A_X : X \rightarrowtail X$	$\iff$	family of objects $A_{x,x}$ in $\mathcal{V}$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\iff$	morphisms $A_{x',x''} \otimes A_{x,x'} \rightarrow A_{x,x''}$
unit $\eta_A : 1_A \Rightarrow A$	$\iff$	$I \rightarrow A_{x,x'}$

3. $\text{Mnd}(\mathbb{R}\text{ing}) = \text{Alg}$, the global category of algebras over rings.

loose endo-arrow $M_R : R \rightarrowtail R$	$\Leftrightarrow$	(R, R) -bimodule M
multiplication $\mu_M : M \odot M \Rightarrow M$	$\Leftrightarrow$	multiplication $M \otimes_R M \rightarrow M$
unit $\eta_M : 1_M \Rightarrow M$	$\Leftrightarrow$	unit $R \rightarrow M$
monad morphism	$\Leftrightarrow$	algebra homomorphism

Dually, $\text{Cmd}(\mathbb{R}\text{ing})$ is the global category Coalg of coalgebras over rings.

4. $\text{Mnd}(\mathcal{V}\text{-Mat}) = \mathcal{V}\text{-Cat}$.

loose endo-arrow $A_X : X \rightarrowtail X$	$\Leftrightarrow$	family of objects $A_{x,x}$ in $\mathcal{V}$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\Leftrightarrow$	morphisms $A_{x',x''} \otimes A_{x,x'} \rightarrow A_{x,x''}$
unit $\eta_A : 1_A \Rightarrow A$	$\Leftrightarrow$	$I \rightarrow A_{x,x'}$
monad morphism	$\Leftrightarrow$	$\mathcal{V}$ -functors

3. $\text{Mnd}(\mathbb{R}\text{ing}) = \text{Alg}$, the global category of algebras over rings.

loose endo-arrow $M_R : R \rightarrowtail R$	$\Leftrightarrow$	(R, R) -bimodule M
multiplication $\mu_M : M \odot M \Rightarrow M$	$\Leftrightarrow$	multiplication $M \otimes_R M \rightarrow M$
unit $\eta_M : 1_M \Rightarrow M$	$\Leftrightarrow$	unit $R \rightarrow M$
monad morphism	$\Leftrightarrow$	algebra homomorphism

Dually, $\text{Cmd}(\mathbb{R}\text{ing})$ is the global category Coalg of coalgebras over rings.

4. $\text{Mnd}(\mathcal{V}\text{-Mat}) = \mathcal{V}\text{-Cat}$.

loose endo-arrow $A_X : X \rightarrowtail X$	$\Leftrightarrow$	family of objects $A_{x,x}$ in $\mathcal{V}$
multiplication $\mu_A : A \odot A \Rightarrow A$	$\Leftrightarrow$	morphisms $A_{x',x''} \otimes A_{x,x'} \rightarrow A_{x,x''}$
unit $\eta_A : 1_A \Rightarrow A$	$\Leftrightarrow$	$I \rightarrow A_{x,x'}$
monad morphism	$\Leftrightarrow$	$\mathcal{V}$ -functors

Dually, $\text{Cmd}(\mathcal{V}\text{-Mat})$ are $\mathcal{V}$ -cocategories and $\mathcal{V}$ -cofunctors.

(co)Free (co)monads

Theorem (free monads, [ASTV])

Suppose that $\mathbb{D}$ is a fibrant double category. If either of the following holds

- 1. $\mathbb{D}_0$ and $\mathcal{H}(\mathbb{D})(X, Y)$ have coproducts which are preserved by $A \odot -$ and $- \odot A$ for any $X, Y \in \mathbb{D}_0$ and $A \in \mathbb{D}_1$.*

Theorem (free monads, [ASTV])

Suppose that $\mathbb{D}$ is a fibrant double category. If either of the following holds

- 1. $\mathbb{D}_0$ and $\mathcal{H}(\mathbb{D})(X, Y)$ have coproducts which are preserved by $A \odot -$ and $- \odot A$ for any $X, Y \in \mathbb{D}_0$ and $A \in \mathbb{D}_1$.*
- 2. $\mathbb{D}$ is locally presentable*

Theorem (free monads, [ASTV])

Suppose that $\mathbb{D}$ is a fibrant double category. If either of the following holds

- 1. $\mathbb{D}_0$ and $\mathcal{H}(\mathbb{D})(X, Y)$ have coproducts which are preserved by $A \odot -$ and $- \odot A$ for any $X, Y \in \mathbb{D}_0$ and $A \in \mathbb{D}_1$.*
- 2. $\mathbb{D}$ is locally presentable*

then the forgetful functor $U: \text{Mnd}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$ has a left adjoint.

Theorem (free monads, [ASTV])

Suppose that $\mathbb{D}$ is a fibrant double category. If either of the following holds

- 1. $\mathbb{D}_0$ and $\mathcal{H}(\mathbb{D})(X, Y)$ have coproducts which are preserved by $A \odot -$ and $- \odot A$ for any $X, Y \in \mathbb{D}_0$ and $A \in \mathbb{D}_1$.*
- 2. $\mathbb{D}$ is locally presentable*

then the forgetful functor $U: \text{Mnd}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$ has a left adjoint.

Proof. Both $\text{Mnd}(\mathbb{D})$ and $\text{End}(\mathbb{D})$ are fibered over $\mathbb{D}_0$ [ASV]

Theorem (free monads, [ASTV])

Suppose that $\mathbb{D}$ is a fibrant double category. If either of the following holds

1. $\mathbb{D}_0$ and $\mathcal{H}(\mathbb{D})(X, Y)$ have coproducts which are preserved by $A \odot -$ and $- \odot A$ for any $X, Y \in \mathbb{D}_0$ and $A \in \mathbb{D}_1$.
2. $\mathbb{D}$ is locally presentable

then the forgetful functor $U: \text{Mnd}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$ has a left adjoint.

Proof. Both $\text{Mnd}(\mathbb{D})$ and $\text{End}(\mathbb{D})$ are fibered over $\mathbb{D}_0$ [ASV] and U is a fibered functor between these fibrations

$$\begin{array}{ccc} \text{Mnd}(\mathbb{D}) & \xrightarrow{U} & \text{End}(\mathbb{D}) \\ & \searrow & \swarrow \\ & \mathbb{D}_0 & \end{array}$$

Theorem (free monads, [ASTV])

Suppose that $\mathbb{D}$ is a fibrant double category. If either of the following holds

1. $\mathbb{D}_0$ and $\mathcal{H}(\mathbb{D})(X, Y)$ have coproducts which are preserved by $A \odot -$ and $- \odot A$ for any $X, Y \in \mathbb{D}_0$ and $A \in \mathbb{D}_1$.
2. $\mathbb{D}$ is locally presentable

then the forgetful functor $U: \text{Mnd}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$ has a left adjoint.

Proof. Both $\text{Mnd}(\mathbb{D})$ and $\text{End}(\mathbb{D})$ are fibered over $\mathbb{D}_0$ [ASV] and U is a fibered functor between these fibrations

$$\begin{array}{ccc} \text{Mnd}(\mathbb{D}) & \xrightarrow{U} & \text{End}(\mathbb{D}) \\ & \searrow & \swarrow \\ & \mathbb{D}_0 & \end{array}$$

So U has a left adjoint as long as each restriction on the fibers $U_X: \text{Mnd}(\mathbb{D})_X \rightarrow \text{End}(\mathbb{D})_X$ has a left adjoint. $\square$

Point (2) can be dualized.

(co)Monadicity

Theorem (monadicity of monads, [ASTV])

If $\mathbb{D}$ is a fibrant double category such that the forgetful functor $U : \text{Mnd}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$ has a left adjoint

Theorem (monadicity of monads, [ASTV])

If $\mathbb{D}$ is a fibrant double category such that the forgetful functor $U : \text{Mnd}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$ has a left adjoint then $\text{Mnd}(\mathbb{D})$ is monadic over $\text{End}(\mathbb{D})$.

Proof. By Beck's Monadicity Theorem.

Theorem (monadicity of monads, [ASTV])

If $\mathbb{D}$ is a fibrant double category such that the forgetful functor $U : \text{Mnd}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$ has a left adjoint then $\text{Mnd}(\mathbb{D})$ is monadic over $\text{End}(\mathbb{D})$.

Proof. By Beck's Monadicity Theorem. U has a left adjoint by hypothesis

Theorem (monadicity of monads, [ASTV])

If $\mathbb{D}$ is a fibrant double category such that the forgetful functor $U : \text{Mnd}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$ has a left adjoint then $\text{Mnd}(\mathbb{D})$ is monadic over $\text{End}(\mathbb{D})$.

Proof. By Beck's Monadicity Theorem. U has a left adjoint by hypothesis and reflects isos.

Theorem (monadicity of monads, [ASTV])

If $\mathbb{D}$ is a fibrant double category such that the forgetful functor $U : \text{Mnd}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$ has a left adjoint then $\text{Mnd}(\mathbb{D})$ is monadic over $\text{End}(\mathbb{D})$.

Proof. By Beck's Monadicity Theorem. U has a left adjoint by hypothesis and reflects isos. Lastly, If a pair of parallel *monad* maps

$$\begin{array}{ccc} X & \xrightarrow{M} & X \\ f \downarrow & \alpha_f, \beta_g & \downarrow g \\ Y & \xrightarrow{N} & Y \end{array}$$

have a *split* coequalizer in $\text{End}(\mathbb{D})$

$$\begin{array}{ccccc} & T_t & & S_s & \\ & \curvearrowright & & \curvearrowright & \\ M_X & \xrightarrow{\alpha_f} & N_Y & \xrightarrow{\gamma_h} & C_Z \\ & \xleftarrow{\beta_g} & & & \end{array}$$

Theorem (monadicity of monads, [ASTV])

If $\mathbb{D}$ is a fibrant double category such that the forgetful functor $U : \mathbf{Mnd}(\mathbb{D}) \rightarrow \mathbf{End}(\mathbb{D})$ has a left adjoint then $\mathbf{Mnd}(\mathbb{D})$ is monadic over $\mathbf{End}(\mathbb{D})$.

Proof. By Beck's Monadicity Theorem. U has a left adjoint by hypothesis and reflects isos. Lastly, If a pair of parallel *monad* maps

$$\begin{array}{ccc}
 X & \xrightarrow{M} & X \\
 f \downarrow & \alpha_f, \beta_g & \downarrow g \\
 Y & \xrightarrow{N} & Y
 \end{array}$$

have a *split* coequalizer in $\mathbf{End}(\mathbb{D})$

$$\begin{array}{ccccc}
 & T_t & & S_s & \\
 & \curvearrowright & & \curvearrowright & \\
 M_X & \xrightarrow{\alpha_f} & N_Y & \xrightarrow{\gamma_h} & C_Z \\
 & \xrightarrow{\beta_g} & & &
 \end{array}$$

then C_Z can be given the structure of a monad such that the above diagram is a coequalizer in $\mathbf{Mnd}(\mathbb{D})$.

For the multiplication apply $-\odot-$ to the previous diagram to obtain another coequalizer and use the multiplication on N .

$$\begin{array}{ccccc}
 M_X \odot M_X & \xrightleftharpoons[\beta_g \odot \beta_g]{\alpha_f \odot \alpha_f} & N_Y \odot N_Y & \xrightarrow{\gamma_h \odot \gamma_h} & C_Z \odot C_Z \\
 \mu_M \downarrow & & \mu_N \downarrow & & \downarrow \exists! \mu_C \\
 M_X & \xrightleftharpoons[\beta_g]{\alpha_f} & N_Y & \xrightarrow{\gamma_h} & C_Z
 \end{array}$$

For the multiplication apply $-\odot-$ to the previous diagram to obtain another coequalizer and use the multiplication on N .

$$\begin{array}{ccccc}
 M_X \odot M_X & \xrightarrow[\beta_g \odot \beta_g]{\alpha_f \odot \alpha_f} & N_Y \odot N_Y & \xrightarrow{\gamma_h \odot \gamma_h} & C_Z \odot C_Z \\
 \downarrow \mu_M & & \downarrow \mu_N & & \downarrow \exists! \mu_C \\
 M_X & \xrightarrow[\beta_g]{\alpha_f} & N_Y & \xrightarrow{\gamma_h} & C_Z
 \end{array}$$

For the unit apply $1_{(-)} \circ \mathfrak{s}$

$$\begin{array}{ccccc}
 1_X & \xrightarrow[1_g]{1_f} & 1_Y & \xrightarrow{1_h} & 1_Z \\
 \downarrow \eta_M & & \downarrow \eta_N & & \downarrow \exists! \eta_C \\
 M_X & \xrightarrow[\beta_g]{\alpha_f} & N_Y & \xrightarrow{\gamma_h} & C_Z
 \end{array}$$

and use similar arguments. □

For the multiplication apply $-\odot-$ to the previous diagram to obtain another coequalizer and use the multiplication on N .

$$\begin{array}{ccccc}
 M_X \odot M_X & \xrightarrow[\beta_g \odot \beta_g]{\alpha_f \odot \alpha_f} & N_Y \odot N_Y & \xrightarrow{\gamma_h \odot \gamma_h} & C_Z \odot C_Z \\
 \downarrow \mu_M & & \downarrow \mu_N & & \downarrow \exists! \mu_C \\
 M_X & \xrightarrow[\beta_g]{\alpha_f} & N_Y & \xrightarrow{\gamma_h} & C_Z
 \end{array}$$

For the unit apply $1_{(-)} \circ \mathfrak{s}$

$$\begin{array}{ccccc}
 1_X & \xrightarrow[1_g]{1_f} & 1_Y & \xrightarrow{1_h} & 1_Z \\
 \downarrow \eta_M & & \downarrow \eta_N & & \downarrow \exists! \eta_C \\
 M_X & \xrightarrow[\beta_g]{\alpha_f} & N_Y & \xrightarrow{\gamma_h} & C_Z
 \end{array}$$

and use similar arguments. □

Everything can be *dualized*, giving comonadicity of comonads over $\text{End}(\mathbb{D})$.

Local presentability

Theorem ($\mathbf{Mnd}(\mathbb{D})$ is locally presentable, [ASTV])

If $\mathbb{D}$ is locally presentable and the functors $-\odot-$ and $1_{(-)}$ preserve filtered colimits then $\mathbf{Mnd}(\mathbb{D})$ is locally presentable.

Theorem ($\mathbf{Mnd}(\mathbb{D})$ is locally presentable, [ASTV])

If $\mathbb{D}$ is locally presentable and the functors $- \odot -$ and $1_{(-)}$ preserve filtered colimits then $\mathbf{Mnd}(\mathbb{D})$ is locally presentable.

Proof. Consider the endofunctor

$$F : \mathbf{End}(\mathbb{D}) \rightarrow \mathbf{End}(\mathbb{D})$$

$$M_X \mapsto (M_X \odot M_X) + M_X.$$

Theorem ($\mathbf{Mnd}(\mathbb{D})$ is locally presentable, [ASTV])

If $\mathbb{D}$ is locally presentable and the functors $-\odot-$ and $1_{(-)}$ preserve filtered colimits then $\mathbf{Mnd}(\mathbb{D})$ is locally presentable.

Proof. Consider the endofunctor

$$F: \mathbf{End}(\mathbb{D}) \rightarrow \mathbf{End}(\mathbb{D})$$

$$M_X \mapsto (M_X \odot M_X) + M_X.$$

F is finitary and $\mathbf{End}(\mathbb{D})$ is locally presentable since $\mathbb{D}$ is [ASV], therefore $\mathbf{Alg}(F)$ is locally presentable.

Theorem ($\mathbf{Mnd}(\mathbb{D})$ is locally presentable, [ASTV])

If $\mathbb{D}$ is locally presentable and the functors $-\odot-$ and $1_{(-)}$ preserve filtered colimits then $\mathbf{Mnd}(\mathbb{D})$ is locally presentable.

Proof. Consider the endofunctor

$$F : \mathbf{End}(\mathbb{D}) \rightarrow \mathbf{End}(\mathbb{D})$$

$$M_X \mapsto (M_X \odot M_X) + M_X.$$

F is finitary and $\mathbf{End}(\mathbb{D})$ is locally presentable since $\mathbb{D}$ is [ASV], therefore $\mathbf{Alg}(F)$ is locally presentable.

$\mathbf{Mnd}(\mathbb{D})$ is a full subcategory of $\mathbf{Alg}(F)$ closed under limits and filtered colimits. Thus, by the Reflection Theorem [AR], $\mathbf{Mnd}(\mathbb{D})$ is also locally presentable. $\square$

Theorem ($\mathbf{Mnd}(\mathbb{D})$ is locally presentable, [ASTV])

If $\mathbb{D}$ is locally presentable and the functors $-\odot-$ and $1_{(-)}$ preserve filtered colimits then $\mathbf{Mnd}(\mathbb{D})$ is locally presentable.

Proof. Consider the endofunctor

$$F: \mathbf{End}(\mathbb{D}) \rightarrow \mathbf{End}(\mathbb{D})$$

$$M_X \mapsto (M_X \odot M_X) + M_X.$$

F is finitary and $\mathbf{End}(\mathbb{D})$ is locally presentable since $\mathbb{D}$ is [ASV], therefore $\mathbf{Alg}(F)$ is locally presentable.

$\mathbf{Mnd}(\mathbb{D})$ is a full subcategory of $\mathbf{Alg}(F)$ closed under limits and filtered colimits. Thus, by the Reflection Theorem [AR], $\mathbf{Mnd}(\mathbb{D})$ is also locally presentable. $\square$

The above theorem *cannot* be dualized.

Theorem ($\text{Cmd}(\mathbb{D})$ is locally presentable, [ASTV])

Suppose $\mathbb{D}$ is locally presentable and the functors $- \odot -$ and $1_{(-)}$ preserve filtered colimits. If furthermore, the categories $\mathcal{H}(\mathbb{D})(X, Y)$ have initial objects preserved by $A \odot -$ and $- \odot A$ for any A , then $\text{Cmd}(\mathbb{D})$ is locally presentable.

Theorem ($\text{Cmd}(\mathbb{D})$ is locally presentable, [ASTV])

Suppose $\mathbb{D}$ is locally presentable and the functors $- \odot -$ and $1_{(-)}$ preserve filtered colimits. If furthermore, the categories $\mathcal{H}(\mathbb{D})(X, Y)$ have initial objects preserved by $A \odot -$ and $- \odot A$ for any A , then $\text{Cmd}(\mathbb{D})$ is locally presentable.

Proof. Consider the endofunctor

$$F : \text{End}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$$

$$C_X \mapsto (C_X \odot C_X) \times 1_X.$$

Theorem ($\text{Cmd}(\mathbb{D})$ is locally presentable, [ASTV])

Suppose $\mathbb{D}$ is locally presentable and the functors $- \odot -$ and $1_{(-)}$ preserve filtered colimits. If furthermore, the categories $\mathcal{H}(\mathbb{D})(X, Y)$ have initial objects preserved by $A \odot -$ and $- \odot A$ for any A , then $\text{Cmd}(\mathbb{D})$ is locally presentable.

Proof. Consider the endofunctor

$$F : \text{End}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$$

$$C_X \mapsto (C_X \odot C_X) \times 1_X.$$

Again, F is finitary and $\text{End}(\mathbb{D})$ is locally presentable [ASV], therefore $\text{Coalg}(F)$ is also locally presentable.

Theorem ($\text{Cmd}(\mathbb{D})$ is locally presentable, [ASTV])

Suppose $\mathbb{D}$ is locally presentable and the functors $- \odot -$ and $1_{(-)}$ preserve filtered colimits. If furthermore, the categories $\mathcal{H}(\mathbb{D})(X, Y)$ have initial objects preserved by $A \odot -$ and $- \odot A$ for any A , then $\text{Cmd}(\mathbb{D})$ is locally presentable.

Proof. Consider the endofunctor

$$F : \text{End}(\mathbb{D}) \rightarrow \text{End}(\mathbb{D})$$

$$C_X \mapsto (C_X \odot C_X) \times 1_X.$$

Again, F is finitary and $\text{End}(\mathbb{D})$ is locally presentable [ASV], therefore $\text{Coalg}(F)$ is also locally presentable.

$\text{Cmd}(\mathbb{D})$ is contained in $\text{Coalg}(F)$ but for F -coalgebras, their structure maps are neither globular nor do they need to satisfy any coassociativity and counitality conditions.

Consider first the full subcategory $\overline{\text{Coalg}(F)}$ of $\text{Coalg}(F)$ spanned by F -coalgebras with *globular* structure maps.

Consider first the full subcategory $\overline{\text{Coalg}(F)}$ of $\text{Coalg}(F)$ spanned by F -coalgebras with *globular* structure maps.

$\overline{\text{Coalg}(F)}$ is closed under all colimits in $\text{Coalg}(F)$, thus cocomplete.

Consider first the full subcategory $\overline{\text{Coalg}(F)}$ of $\text{Coalg}(F)$ spanned by F -coalgebras with *globular* structure maps.

$\overline{\text{Coalg}(F)}$ is closed under all colimits in $\text{Coalg}(F)$, thus cocomplete.


Furthermore, we can prove that

Lemma [ASTV]

Under the above assumptions if $\alpha_f : C_X \Rightarrow D_Y$ is a monomorphism in $\text{Coalg}(F)$ then its source/target $f : X \rightarrow Y$ is a monomorphism in $\mathbb{D}_0$.

and thus conclude that $\overline{\text{Coalg}(F)}$ is closed under all subobjects in $\text{Coalg}(F)$ hence accessible [AR]. So $\overline{\text{Coalg}(F)}$ is locally presentable.


$\text{Cmd}(\mathbb{D})$ is now the equifier of



in $\overline{\text{Coalg}(F)}$, that is the subcategory spanned of those objects C for which $(\phi_i)_C = (\psi_i)_C$ for $i = 1, 2, 3$ where

$$\left\{ \begin{array}{l} (\phi_1)_C = (\delta_C * \text{id}_C) \circ \delta_C : C \rightarrow (C \odot C) \odot C \\ (\psi_1)_C = (\text{id}_C * \delta_C) \circ \delta_C : C \rightarrow C \odot (C \odot C) \\ (\phi_2)_C = (\epsilon_C * \text{id}_C) \circ \delta : C \rightarrow 1_{s(C)} \odot C \\ (\psi_2)_C = \ell_C : C \rightarrow 1_{s(C)} \odot C \\ (\phi_3)_C = (\text{id}_C * \epsilon_C) \circ \delta : C \rightarrow C \odot 1_{t(C)} \\ (\psi_3)_C = r_C : C \rightarrow C \odot 1_{t(C)} \end{array} \right.$$

$\text{Cmd}(\mathbb{D})$ is now the equifier of



in $\overline{\text{Coalg}(F)}$, that is the subcategory spanned of those objects C for which $(\phi_i)_C = (\psi_i)_C$ for $i = 1, 2, 3$ where

$$\left\{ \begin{array}{l} (\phi_1)_C = (\delta_C * \text{id}_C) \circ \delta_C : C \rightarrow (C \odot C) \odot C \\ (\psi_1)_C = (\text{id}_C * \delta_C) \circ \delta_C : C \rightarrow C \odot (C \odot C) \\ (\phi_2)_C = (\epsilon_C * \text{id}_C) \circ \delta : C \rightarrow 1_{s(C)} \odot C \\ (\psi_2)_C = \ell_C : C \rightarrow 1_{s(C)} \odot C \\ (\phi_3)_C = (\text{id}_C * \epsilon_C) \circ \delta : C \rightarrow C \odot 1_{t(C)} \\ (\psi_3)_C = r_C : C \rightarrow C \odot 1_{t(C)} \end{array} \right.$$

All functors are accessible, therefore $\text{Cmd}(\mathbb{D})$ is accessible. Furthermore it is cocomplete [ASV], hence locally presentable.

- [AR] J. Adámek, J. Rosický, *Locally presentable and Accessible Categories*, London Mathematical Society Lecture Note Series 189, Cambridge University Press.
- [ASV] V. Aravantinos-Sotiropoulos, C. Vasilakopoulou, *Sweedler Theory for double categories*. arXiv:2408.03180.
- [ASTV] V. Aravantinos-Sotiropoulos, T.T., C. Vasilakopoulou, *On categories of monads and comonads in double categories*. arXiv:2606.26755.

Thank you for your attention!