

1 Vector Calculus

Scalar Fields

$$f: \mathbb{R}^n \rightarrow \mathbb{R}$$

Vector Fields

$$\underline{v}: \mathbb{R}^n \rightarrow \mathbb{R}^m$$

Differentiation of Vector Fields

Let $\underline{a}, \underline{b}, \underline{c}$ be vector fields and φ be a scalar field, then

$$\begin{aligned}\frac{d}{dt}(\varphi \underline{a}) &= \frac{d\varphi}{dt} \underline{a} + \varphi \frac{d\underline{a}}{dt} \\ \frac{d}{dt}(\underline{a} \cdot \underline{b}) &= \frac{d\underline{a}}{dt} \cdot \underline{b} + \underline{a} \cdot \frac{d\underline{b}}{dt} \\ \frac{d}{dt}(\underline{a} \times \underline{b}) &= \frac{d\underline{a}}{dt} \times \underline{b} + \underline{a} \times \frac{d\underline{b}}{dt} \\ \frac{d}{dt}(\underline{a}(\varphi(t))) &= \frac{d\underline{a}}{d\varphi} \frac{d\varphi}{dt}\end{aligned}$$

Differential Operators

For a scalar field $f: \mathbb{R}^n \rightarrow \mathbb{R}, \underline{x} \mapsto f(\underline{x})$ the gradient is

$$\text{grad } f(\underline{x}) = \begin{pmatrix} \frac{\partial f}{\partial x_1} \\ \vdots \\ \frac{\partial f}{\partial x_n} \end{pmatrix}$$

and gives the direction of [steepest ascent](#)

For a vector field $\underline{v}: \mathbb{R}^n \rightarrow \mathbb{R}^n$ the divergence is defined as

$$\text{div } \underline{v}(\underline{x}) = \sum_{j=1}^n \frac{\partial v_j}{\partial x_j}(\underline{x})$$

and gives the [source density](#) at a point

For a vector field $\underline{v}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ the curl is defined as

$$\text{curl}(\underline{v}(x, y, z)) = \begin{pmatrix} \frac{\partial v_3}{\partial y} - \frac{\partial v_2}{\partial z} \\ \frac{\partial v_1}{\partial z} - \frac{\partial v_3}{\partial x} \\ \frac{\partial v_2}{\partial x} - \frac{\partial v_1}{\partial y} \end{pmatrix}$$

and gives the [vortex strength](#)

The Laplace operator for a scalar field $f: \mathbb{R}^n \rightarrow \mathbb{R}, \underline{x} \mapsto f(\underline{x})$ gives

$$\Delta f(\underline{x}) = \nabla \cdot \nabla f(\underline{x}) = \sum_{j=1}^n \frac{\partial^2 f}{\partial^2 x_j}(\underline{x})$$

Expressed with the ∇ operator we get the representations

$$\begin{aligned}\text{grad } f &= \nabla f \\ \text{div } \underline{v} &= \nabla \cdot \underline{v} \\ \text{curl } \underline{v} &= \nabla \times \underline{v} \\ \Delta &= \nabla \cdot \nabla\end{aligned}$$

Computation Rules for Differential Operators

The three operators grad, div and curl are linear.

Rules for one active operator:

$$\begin{aligned}\text{grad}(f_1 f_2) &= f_1 \text{ grad } f_2 + f_2 \text{ grad } f_1 \\ \Leftrightarrow \nabla(f_1 f_2) &= f_1 \nabla f_2 + f_2 \nabla f_1\end{aligned}$$

$$\begin{aligned}\text{grad } F(f) &= F'(f) \text{ grad } f \\ \Leftrightarrow \nabla F(f) &= F'(f) \nabla f\end{aligned}$$

$$\begin{aligned}\text{div}(\underline{v}_1 \times \underline{v}_2) &= \underline{v}_2 \cdot \text{curl } \underline{v}_1 - \underline{v}_1 \cdot \text{curl } \underline{v}_2 \\ \Leftrightarrow \nabla \cdot (\underline{v}_1 \times \underline{v}_2) &= \underline{v}_2 \cdot (\nabla \times \underline{v}_1) - \underline{v}_1 \cdot (\nabla \times \underline{v}_2)\end{aligned}$$

Rules for the interactions between vector- and scalarfields:

$$\begin{aligned}\text{div } f \underline{v} &= \underline{v} \text{ grad } f + f \text{ div } \underline{v} \\ \Leftrightarrow \nabla \cdot (f \underline{v}) &= \underline{v} \nabla f + f(\nabla \cdot \underline{v})\end{aligned}$$

$$\begin{aligned}\text{curl } f \underline{v} &= f \text{ curl } \underline{v} - \underline{v} \times \text{grad } f \\ \Leftrightarrow \nabla \times (f \underline{v}) &= f(\nabla \times \underline{v}) - \underline{v} \times \nabla f\end{aligned}$$

Rules for the concatenation of the operators:

$$\begin{aligned}\text{div curl } \underline{v} &= 0 \\ \Leftrightarrow \nabla \cdot (\nabla \times \underline{v}) &= 0\end{aligned}$$

$$\begin{aligned}\text{curl grad } f &= \underline{0} \\ \Leftrightarrow \nabla \times \nabla f &= \underline{0}\end{aligned}$$

$$\begin{aligned}\text{div grad } f &= \Delta f \\ \Leftrightarrow \nabla \cdot \nabla f &= \Delta f\end{aligned}$$

$$\begin{aligned}\text{curl curl } \underline{v} &= \text{grad div } \underline{v} - \Delta \underline{v} \\ \Leftrightarrow \nabla \times (\nabla \times \underline{v}) &= \nabla(\nabla \cdot \underline{v}) - \Delta \underline{v}\end{aligned}$$

Flux (Φ)

Q: Let be \underline{v} the velocity field of a flow; how much fluid flows through a surface S per unit time in direction \underline{n} ?

A: Split S into infinitesimal area elements dS . Because they are infinitesimal, the dS are flat and \underline{v} is homogeneous within a single dS . The total flow through a single dS is

$$d\Phi = \underline{v} \cdot \underline{n} dS$$

The flux is the “sum” over all infinitesimal dS and thus given by the integral

$$\Phi = \int_S \underline{v} \cdot \underline{n} dS$$

2 Modeling Spatial Effects

Reynolds transport theorem

$$\begin{aligned}\frac{D\varphi}{Dt} &= \frac{D}{Dt} \int_{V(t)} f dV(t) \\ &= \int_{V(t)} \frac{Df}{Dt} dV(t) + \int_{V(t)} f \underbrace{\frac{D}{Dt}[dV(t)]}_{\frac{\partial dV(t)}{\partial t} + (\underline{v} \cdot \nabla) dV(t)} \\ &= \int_{V(t)} \left[\frac{Df}{Dt} + f(\underline{v} \cdot \nabla) \right] dV(t) \\ &= \int_{V(t)} \left[\frac{\partial f}{\partial t} + \underbrace{\underline{v} \cdot (\nabla f) + f(\nabla \cdot \underline{v})}_{= \nabla \cdot (f \underline{v}): \text{ see compute rules}} \right] dV(t) \\ &= \int_{V(t)} \left[\frac{\partial f}{\partial t} + \nabla \cdot (f \underline{v}) \right] dV(t).\end{aligned}$$

Vector Fields

$$\underline{v}: \mathbb{R}^n \rightarrow \mathbb{R}^m$$