

Probability Theory and Mathematical Statistics Notes

Contents

1	Chapter 1: Basic Probability	3
1.1	Set Operations and Logic	3
1.2	Probability Properties and Inclusion-Exclusion	3
1.3	Classical Probability (Sampling)	3
1.4	The Matching Problem	3
1.5	Conditional Probability and Independence	4
1.6	Calculations with Set Operations	4
1.7	Sampling Example (Urn Problem)	4
1.8	Two-Urn Transfer Problem (Total Probability & Bayes)	5
2	Chapter 2: Random Variables and Distributions	5
2.1	Discrete Random Variables and Common Distributions	5
2.2	Distribution Functions (CDF) and Probability Density Functions (PDF)	5
2.2.1	Cumulative Distribution Function (CDF)	5
2.2.2	Probability Density Function (PDF)	6
2.3	Continuous Random Variables and Common Distributions	6
2.4	Selected Examples	6
2.4.1	Discrete Probability Calculation	6
2.4.2	From CDF to Probabilities	6
2.4.3	Finding Constants in CDF	7
2.4.4	Quadratic Equation Probability	7
2.4.5	Normal Distribution Symmetry	7
2.5	Functions of Random Variables	7
2.5.1	Discrete Case	7
2.5.2	Continuous Case	8
3	Chapter 3: Multidimensional Random Variables	8
3.1	Definitions and Basic Properties	8
3.1.1	Joint Distribution Function (CDF)	8
3.1.2	Marginal Distribution Functions	8
3.1.3	Conditional Distribution (Discrete Case)	8
3.1.4	Joint Probability Density Function (PDF)	8
3.1.5	Marginal PDFs	8
3.1.6	Conditional PDFs	8
3.1.7	Independence	9
3.2	Discrete Example: Determination of Parameters	9
3.3	Discrete Example: Balls in Boxes	9
3.4	Continuous Example: Triangular Region	10
3.5	Example: Finding the PDF of the Difference of Two Random Variables	11
4	Chapter 4: Numerical Characteristics of Random Variables	12
4.1	Expectation (Mean)	12
4.1.1	Properties of Expectation	12
4.2	Variance	12
4.2.1	Properties of Variance	12
4.3	Covariance and Correlation Coefficient	12
4.3.1	Covariance	12

4.3.2	Correlation Coefficient	12
4.4	Expectation and Variance of Common Distributions	12
4.5	Examples and Applications	12
4.5.1	Example 1: Variance of a Linear Combination	12
4.5.2	Example 2: Finding $E(X^2)$ for a Binomial Variable	13
4.5.3	Example 3: Properties of Covariance	13
4.5.4	Example 4: Correlation for Joint PDF	13
4.6	Chebyshev's Inequality	13
4.6.1	Example: Applying Chebyshev's Inequality	13
5	Chapter 5: Limit Theorems	14
5.1	Chebyshev's Inequality	14
5.2	Laws of Large Numbers (LLN)	14
5.2.1	Khinchine's Law of Large Numbers	14
5.3	Central Limit Theorem (CLT)	14
5.4	Applications of the Central Limit Theorem	14
5.4.1	Example 1: Sum of Exponential Lifetimes	14
5.4.2	Example 2: Probability of Defects (Normal Approximation to Binomial)	15
6	Chapter 6: Samples and Sampling Distributions	15
6.1	Sample Statistics	15
6.1.1	Definitions of Common Sample Statistics	15
6.1.2	Estimation of Population Characteristics	16
6.2	Important Sampling Distributions	16
6.2.1	Distribution of the Sample Mean	16
6.2.2	Chi-Square (χ^2) Distribution	16
6.2.3	t-Distribution (Student's t-Distribution)	16
6.2.4	F-Distribution	16
7	Chapter 7: Parameter Estimation	16
7.1	Point Estimation	16
7.1.1	Method of Moments	16
7.1.2	Maximum Likelihood Estimation (MLE)	17
7.2	Properties of Estimators	17
7.2.1	Unbiasedness and Efficiency	17
7.3	Interval Estimation (Confidence Intervals)	18
7.3.1	Confidence Interval for the Population Mean μ	18
7.3.2	Confidence Interval for the Population Variance σ^2	18
8	Chapter 8: Hypothesis Testing	19
8.1	Concepts and Structure of Hypothesis Testing	19
8.2	Inference for the Population Mean (μ)	19
8.3	Inference for the Population Variance (σ^2)	19
8.4	Examples	20
8.4.1	Example: Mean μ Test (Variance Known)	20
8.4.2	Example: Variance σ^2 Test	20

1 Chapter 1: Basic Probability

1.1 Set Operations and Logic

- **Complement/Inverse:** $A \cup \bar{A} = S$ and $A\bar{A} = \emptyset$.
- **Circuit Systems:**
 - Series connection corresponds to intersection: $\bigcap A_i$.
 - Parallel connection corresponds to union: $\bigcup A_i$.
- **Difference Formulas:**
 - If $A \subset B$, then $P(B - A) = P(B) - P(A)$.
 - In general, $P(B - A) = P(B) - P(AB)$.

1.2 Probability Properties and Inclusion-Exclusion

Example: Persons A, B, and C arrive. The probability of each arriving is 0.4. The probability that at least 2 arrive is 0.3. The probability that all arrive is 0.05. Find the probability that at least 1 arrives.

Solution: Given:

$$P(A) = P(B) = P(C) = 0.4$$

$$P(AB \cup BC \cup AC) = 0.3$$

$$P(ABC) = 0.05$$

Using the relationship for the union of intersections:

$$P(AB \cup BC \cup AC) = P(AB) + P(BC) + P(AC) - 2P(ABC)$$

$$0.3 = \sum P(AB) - 2(0.05) \implies \sum P(AB) = 0.4$$

To find the probability of at least 1 person arriving ($P(A \cup B \cup C)$):

$$P(A \cup B \cup C) = \sum P(A) - \sum P(AB) + P(ABC)$$

$$P(A \cup B \cup C) = (0.4 + 0.4 + 0.4) - 0.4 + 0.05$$

$$P(A \cup B \cup C) = 1.2 - 0.4 + 0.05 = 0.85$$

1.3 Classical Probability (Sampling)

Example: There are 8 balls: 3 red and 5 yellow. Draw 2 balls without replacement. What is the probability of drawing 1 red and 1 yellow?

Solution: Using Combinations:

$$P = \frac{C_1^3 C_1^5}{C_2^8} = \frac{3 \times 5}{28} = \frac{15}{28}$$

Alternatively, using the Multiplication Rule (considering order then summing):

$$P = 2 \times \frac{3}{8} \times \frac{5}{7} = \frac{15}{28}$$

1.4 The Matching Problem

Example: There are n people and n numbers. What is the probability that no person matches their number (derangement)?

Formula:

$$P = \sum_{k=0}^n \frac{(-1)^k}{k!}$$

1.5 Conditional Probability and Independence

Definition:

$$P(B|A) = \frac{P(AB)}{P(A)}$$

Total Probability Theorem:

$$P(A) = \sum P(AB_j) = \sum P(B_j)P(A|B_j)$$

Bayes' Theorem:

$$P(B_i|A) = \frac{P(B_i)P(A|B_i)}{\sum_j P(B_j)P(A|B_j)}$$

Example: 10 items total, 8 good, 2 defective. Let A_i be the event that the i -th draw is good.

- If sampling without replacement: $P(A_2|A_1) = \frac{7}{9}$.
- Note: The joint probability is $\frac{8}{10} \times \frac{7}{9}$, but since condition A_1 is assumed, we only look at the second stage.

Independence/Chain Rule:

$$P(A_{i_1} \dots A_{i_k}) = \prod_{j=1}^k P(A_{i_j})$$

if mutually independent.

1.6 Calculations with Set Operations

Example: Given $P(B) = 0.4$, $P(A \cup B) = 0.5$. Find $P(A|\bar{B})$.

Solution:

$$P(A|\bar{B}) = \frac{P(A \cap \bar{B})}{P(\bar{B})} = \frac{P(A) - P(AB)}{1 - P(B)}$$

First, find $P(A) - P(AB)$ using the addition rule:

$$P(A \cup B) = P(A) + P(B) - P(AB)$$

$$0.5 = (P(A) - P(AB)) + 0.4$$

$$P(A) - P(AB) = 0.1$$

Substituting back:

$$P(A|\bar{B}) = \frac{0.1}{1 - 0.4} = \frac{0.1}{0.6} = \frac{1}{6}$$

Note: If A and B were independent, then $P(\bar{B}|A) = P(\bar{B}) = 0.6$.

1.7 Sampling Example (Urn Problem)

Example: An urn contains 5 White and 6 Black balls (Total 11). Draw 2 balls.

1. Find the probability of 1 White and 1 Black.
2. Find the probability of at least 1 Black.

Solution: 1. One of each:

$$\frac{C_1^5 C_1^6}{C_2^{11}}$$

2. At least one black (1 minus probability of all white):

$$1 - \frac{C_2^5}{C_2^{11}}$$

1.8 Two-Urn Transfer Problem (Total Probability & Bayes)

Example:

- Urn A: 5 Red, 4 White.
- Urn B: 4 Red, 5 White.
- **Process:** Draw 1 ball from A, put it into B. Then draw 1 ball from B.

Questions: 1. What is the probability that the ball drawn from B is White? 2. If the ball drawn from B is White, what is the probability that the ball transferred from A was White?

Solution:

1. Let W_B be the event B draws White. Let W_A be transfer White, R_A be transfer Red.

$$P(W_B) = P(R_A)P(W_B|R_A) + P(W_A)P(W_B|W_A)$$

$$P(W_B) = \frac{5}{9} \times \frac{5}{10} + \frac{4}{9} \times \frac{6}{10}$$

(Note: Denominator 10 implies B has 9 balls initially + 1 transferred).

2. Bayes' Theorem: Find $P(W_A|W_B)$.

$$P(W_A|W_B) = \frac{P(W_A \cap W_B)}{P(W_B)} = \frac{\frac{4}{9} \times \frac{6}{10}}{\text{Result from (1)}}$$

2 Chapter 2: Random Variables and Distributions

2.1 Discrete Random Variables and Common Distributions

1. **0-1 Distribution (Bernoulli Distribution):** $X \sim B(1, p)$ or $X \sim 0 - 1(p)$.

$$P(X = k) = p^k(1 - p)^{1-k}, \quad k = 0, 1$$

2. **Binomial Distribution:** $X \sim B(n, p)$.

$$P(X = k) = C_k^n p^k (1 - p)^{n-k}, \quad k = 0, 1, \dots, n$$

3. **Poisson Distribution:** $X \sim P(\lambda)$.

$$P(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}, \quad k = 0, 1, 2, \dots, \quad (\lambda > 0)$$

Poisson Approximation: When n is large (> 10) and p is small (< 0.1), the Binomial distribution can be approximated by Poisson with $\lambda = np$:

$$C_k^n p^k (1 - p)^{n-k} \approx \frac{\lambda^k e^{-\lambda}}{k!}$$

4. **Geometric Distribution:** $X \sim G(p)$. Represents the number of trials to get the first success.

$$P(X = k) = p(1 - p)^{k-1}, \quad k = 1, 2, 3, \dots$$

5. **Pascal Distribution (Negative Binomial):** Waiting for the r -th success.

$$P(X = k) = C_{r-1}^{k-1} p^r (1 - p)^{k-r}, \quad k = r, r + 1, \dots$$

2.2 Distribution Functions (CDF) and Probability Density Functions (PDF)

2.2.1 Cumulative Distribution Function (CDF)

Defined as $F(x) = P(X \leq x)$.

- Property: $0 \leq F(x) \leq 1$.
- For discrete variables: $F(x) = \sum_{x_k \leq x} P(X = x_k)$.

2.2.2 Probability Density Function (PDF)

For continuous variables, $F(x) = \int_{-\infty}^x f(t)dt$.

Example: Given a PDF:

$$f(x) = \begin{cases} 1/3 & 0 < x < 1 \\ 2/9 & 3 < x < 6 \\ 0 & \text{others} \end{cases}$$

Calculate $F(x)$:

$$F(x) = \begin{cases} 0 & x < 0 \\ \int_0^x \frac{1}{3} dt = \frac{x}{3} & 0 \leq x < 1 \\ 1/3 & 1 \leq x < 3 \quad (\text{Value holds constant in gap}) \\ 1/3 + \int_3^x \frac{2}{9} dt & 3 \leq x < 6 \\ 1 & x \geq 6 \end{cases}$$

2.3 Continuous Random Variables and Common Distributions

1. **Uniform Distribution:** $X \sim U(a, b)$.

$$f(x) = \begin{cases} \frac{1}{b-a} & a < x < b \\ 0 & \text{others} \end{cases}$$

2. **Exponential Distribution:** $X \sim E(\lambda)$.

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & x > 0 \\ 0 & x \leq 0 \end{cases}$$

3. **Normal (Gaussian) Distribution:** $X \sim N(\mu, \sigma^2)$.

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad -\infty < x < +\infty$$

Curve is symmetric about $x = \mu$.

2.4 Selected Examples

2.4.1 Discrete Probability Calculation

Example: Roll a dice twice. Let X be the smaller number of points. Find the distribution law of X .

Solution: Total outcomes = 36.

- $X = 1$: (1 with 1..6) + (2..6 with 1). Total 11. $P = 11/36$.
- $X = 2$: (2 with 2..6) + (3..6 with 2). Total 9. $P = 9/36$.
- Pattern continues ($2n - 1$ logic decreasing).

Distribution Table:

X	1	2	3	4	5	6
P	11/36	9/36	7/36	5/36	3/36	1/36

2.4.2 From CDF to Probabilities

Example: Given X has CDF:

$$F(x) = \begin{cases} 0 & x < 1 \\ \ln x & 1 \leq x < e \\ 1 & x \geq e \end{cases}$$

1. Find $f(x)$. 2. Find $P(X < 2)$ and $P(0 < X \leq 3)$.

Solution: 1. $f(x) = F'(x) = \frac{1}{x}$ for $1 \leq x < e$, and 0 otherwise. 2. $P(X < 2) = F(2) = \ln 2$. $P(0 < X \leq 3) = F(3) - F(0)$. Note that since max range is $e \approx 2.718$, $F(3) = 1$. $P(0 < X \leq 3) = 1 - 0 = 1$. Alternatively via integration: $\int_1^e \frac{1}{x} dx = \ln e - \ln 1 = 1$.

2.4.3 Finding Constants in CDF

Example: $f(x) = 2e^{-2x}$ for $x \geq 0$. Find $F(x)$.

$$F(x) = \int_0^x 2e^{-2t} dt = -e^{-2t} \Big|_0^x = 1 - e^{-2x}$$

Using boundary conditions $F(+\infty) = 1$, the integration constant resolves.

Example: Continuity of CDF.

$$f(x) = \begin{cases} 1 & 0 < x < 1 \\ 0 & \text{others} \end{cases}$$

$$F(x) = \begin{cases} C_1 & x \leq 0 \\ x + C_2 & 0 < x < 1 \\ C_3 & x > 1 \end{cases}$$

Since F must be a valid CDF:

- $x \leq 0 \implies F(x) = 0 \implies C_1 = 0$.
- $x > 1 \implies F(x) = 1 \implies C_3 = 1$.
- Continuity at $x = 0$: $\lim_{x \rightarrow 0^-} 0 = \lim_{x \rightarrow 0^+} (0 + C_2) \implies C_2 = 0$.

2.4.4 Quadratic Equation Probability

Example: $X \sim U(0, 5)$ (implied). What is the probability that the equation $4t^2 + 4Xt + (X + 2) = 0$ has real solutions for t ?

Solution: Discriminant $\Delta \geq 0$:

$$(4X)^2 - 4(4)(X + 2) \geq 0$$

$$16X^2 - 16(X + 2) \geq 0 \implies X^2 - X - 2 \geq 0$$

$$(X - 2)(X + 1) \geq 0$$

Solution implies $X \geq 2$ or $X \leq -1$. Given domain $X \in (0, 5)$: We look for $P(X \geq 2)$. For Uniform distribution $U(0, 5)$, $f(x) = 1/5$.

$$P(X \geq 2) = \frac{5 - 2}{5 - 0} = \frac{3}{5}$$

2.4.5 Normal Distribution Symmetry

Example: $X \sim N(2, \sigma^2)$. Given $P(2 < X < 4) = 0.3$. Find $P(X < 0)$.

Solution: Due to symmetry about $\mu = 2$: $P(0 < X < 2) = P(2 < X < 4) = 0.3$. Total probability is 1, so half probability is 0.5.

$$P(X < 2) = 0.5$$

$$P(X < 0) + P(0 < X < 2) = 0.5$$

$$P(X < 0) + 0.3 = 0.5 \implies P(X < 0) = 0.2$$

2.5 Functions of Random Variables

2.5.1 Discrete Case

Example: X has distribution:

X	-2	-1	0	1	3
P	1/5	1/6	1/5	1/15	11/30

Find distribution of $Y = X^2$. Possible values for Y : 0, 1, 4.

- $P(Y = 0) = P(X = 0) = 1/5$.
- $P(Y = 1) = P(X = -1) + P(X = 1) = 1/6 + 1/15 = 5/30 + 2/30 = 7/30$.
- $P(Y = 4) = P(X = -2) = 1/5$.
- $P(Y = 9) = P(X = 3) = 11/30$.

2.5.2 Continuous Case

Example: $X \sim U(0, 2)$. Find the PDF of $Y = X^3$. $f_X(x) = 1/2$ for $0 < x < 2$. Range of Y : $(0, 8)$.
CDF Method:

$$F_Y(y) = P(Y \leq y) = P(X^3 \leq y) = P(X \leq y^{1/3})$$

For $y < 0$:

$$F_Y(y) = \int_0^{y^{1/3}} f(x)dx = 0$$

For $0 < y < 8$:

$$F_Y(y) = \int_0^{y^{1/3}} \frac{1}{2}dx = \frac{1}{2}y^{1/3}$$

For $8 < y$:

$$F_Y(y) = \int_0^2 \frac{1}{2}dx = 1$$

Differentiating to find PDF:

$$f_Y(y) = F'_Y(y) = \frac{1}{2} \cdot \frac{1}{3}y^{-2/3} = \frac{1}{6\sqrt[3]{y^2}}, \quad 0 < y < 8$$

3 Chapter 3: Multidimensional Random Variables

3.1 Definitions and Basic Properties

3.1.1 Joint Distribution Function (CDF)

The joint cumulative distribution function for random variables X and Y is defined as:

$$F(x, y) = P(X \leq x, Y \leq y)$$

3.1.2 Marginal Distribution Functions

$$F_X(x) = F(x, +\infty) = \lim_{y \rightarrow \infty} F(x, y)$$

$$F_Y(y) = F(+\infty, y) = \lim_{x \rightarrow \infty} F(x, y)$$

3.1.3 Conditional Distribution (Discrete Case)

$$P(X = x_i | Y = y_j) = \frac{P(X = x_i, Y = y_j)}{P(Y = y_j)}$$

3.1.4 Joint Probability Density Function (PDF)

For continuous variables, the joint PDF $f(x, y)$ satisfies:

$$\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} f(x, y)dx dy = 1$$

3.1.5 Marginal PDFs

$$f_X(x) = \int_{-\infty}^{+\infty} f(x, y)dy$$

$$f_Y(y) = \int_{-\infty}^{+\infty} f(x, y)dx$$

3.1.6 Conditional PDFs

$$f_{X|Y}(x|y) = \frac{f(x, y)}{f_Y(y)}$$

3.1.7 Independence

X and Y are independent if and only if:

$$f(x, y) = f_X(x)f_Y(y)$$

(Or for discrete: $P(X = x, Y = y) = P(X = x)P(Y = y)$ for all pairs).

3.2 Discrete Example: Determination of Parameters

Example Given the joint probability mass function (PMF) of discrete random variables X and Y :

$$P(Y \leq 0 \mid X < 2) = 0.5$$

The joint PMF is presented in the table below:

$X \setminus Y$	-1	0	1
1	a	0.2	0.2
2	0.1	0.1	b

Solution: The total probability must sum to 1:

$$P(X = 1, Y = -1) + P(X = 1, Y = 0) + \dots + P(X = 2, Y = 1) = 1$$

$$a + 0.2 + 0.2 + 0.1 + 0.1 + b = 1 \implies a + b + 0.6 = 1 \implies a + b = 0.4 \quad (\text{Mistake in image: } a + b = 1)$$

Assuming the formula in the image is correct: $a + b = 1$ implies a different set of probabilities must be in the table.

Let's use the given conditional probability $P(Y \leq 0 \mid X < 2) = 0.5$.

- **Event B :** $X < 2$ $P(B) = P(X = 1) = P(X = 1, Y = -1) + P(X = 1, Y = 0) + P(X = 1, Y = 1) = a + 0.2 + 0.2 = a + 0.4$
- **Event $A \cap B$:** $Y \leq 0$ and $X < 2$ $P(A \cap B) = P(X = 1, Y \leq 0) = P(X = 1, Y = -1) + P(X = 1, Y = 0) = a + 0.2$

Using the conditional probability formula:

$$P(Y \leq 0 \mid X < 2) = \frac{P(Y \leq 0 \cap X < 2)}{P(X < 2)} = \frac{a + 0.2}{a + 0.4}$$

We are given that this value is 0.5. Thus, we have the equation:

$$\frac{a + 0.2}{a + 0.4} = 0.5$$

Solving for a :

$$a + 0.2 = 0.5(a + 0.4) \implies a + 0.2 = 0.5a + 0.2 \implies 0.5a = 0 \implies a = 0$$

Since $a = 0$ and the sum of all probabilities must be 1:

$$a + 0.2 + 0.2 + 0.1 + 0.1 + b = 1 \implies 0.6 + b = 1 \implies b = 0.4$$

Therefore, the values are $a = 0$ and $b = 0.4$.

3.3 Discrete Example: Balls in Boxes

Problem: Place 2 balls randomly into 4 boxes (numbered 1, 2, 3, 4). Let X be the number of balls in box 1. Let Y be the number of balls in box 2.

1. Joint Distribution Law

$X \setminus Y$	0	1	2
0	1/4	4/16	1/16
1	4/16	2/16	0
2	1/16	0	0

2. Marginal Distributions

X	0	1	2	Y	0	1	2
P	$\frac{9}{16}$	$\frac{6}{16}$	$\frac{1}{16}$	P	$\frac{9}{16}$	$\frac{6}{16}$	$\frac{1}{16}$

3. Conditional Probability

X	0	1	2
$P(X Y = 1)$	$\frac{1/4}{3/8}$	$\frac{1/8}{3/8}$	$\frac{0}{3/8}$

4. Independence Check if $P_{ij} = P_i \cdot P_j$. For example: $P(X = 2) = 1/16$, $P(Y = 2) = 1/16$. $P(X = 2)P(Y = 2) = 1/256$. But $P(X = 2, Y = 2) = 0$. Since $1/256 \neq 0$, X and Y are **not independent**.

3.4 Continuous Example: Triangular Region

Problem: Given joint PDF:

$$f(x, y) = \begin{cases} kx & 0 \leq x \leq y \leq 1 \\ 0 & \text{others} \end{cases}$$

1. Find Constant k

$$\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} f(x, y) dx dy = 1$$

$$\int_0^1 dy \int_0^y kx dx \quad \text{OR} \quad \int_0^1 dx \int_x^1 kx dy$$

Using the second order (dy first):

$$\begin{aligned} \int_0^1 kx \left(\int_x^1 dy \right) dx &= \int_0^1 kx(1-x) dx = k \int_0^1 (x-x^2) dx \\ &= k \left[\frac{1}{2}x^2 - \frac{1}{3}x^3 \right]_0^1 = k \left(\frac{1}{2} - \frac{1}{3} \right) = \frac{k}{6} \\ \frac{k}{6} &= 1 \implies k = 6 \end{aligned}$$

2. Probability Calculation Find $P(X + Y \leq 1)$. Intersection of region $0 \leq x \leq y$ and $x + y \leq 1$. The line $y = 1 - x$ intersects $y = x$ at $x = 1/2$.

$$\begin{aligned} P &= \int_0^{1/2} dx \int_x^{1-x} 6x dy \\ &= \int_0^{1/2} 6x(1-x-x) dx = \int_0^{1/2} (6x - 12x^2) dx \\ &= [3x^2 - 4x^3]_0^{1/2} = 3(1/4) - 4(1/8) = \frac{3}{4} - \frac{1}{2} = \frac{1}{4} \end{aligned}$$

3. Marginal PDFs

$$\begin{aligned} f_X(x) &= \int_x^1 6x dy = 6x(1-x), \quad 0 < x < 1 \\ f_Y(y) &= \int_0^y 6x dx = 6 \left[\frac{x^2}{2} \right]_0^y = 3y^2, \quad 0 < y < 1 \end{aligned}$$

4. Conditional PDF

$$f_{Y|X}(y|x) = \frac{f(x, y)}{f_X(x)} = \frac{6x}{6x(1-x)} = \frac{1}{1-x}, \quad x < y < 1$$

5. Independence Since $f(x, y) \neq f_X(x)f_Y(y)$ (or simply because the domain is coupled), they are **not independent**.

3.5 Example: Finding the PDF of the Difference of Two Random Variables

Given the joint probability density function (PDF) of the random variables (X, Y) :

$$f(x, y) = \begin{cases} \frac{1}{2}e^{-y/2}, & 0 < x < 1, y > 0 \\ 0, & \text{otherwise} \end{cases}$$

Find the cumulative distribution function (CDF) $F_Z(z)$ and the probability density function (PDF) $f_Z(z)$ of $Z = X - Y$.

1. Finding the Cumulative Distribution Function $F_Z(z)$

The CDF is $F_Z(z) = P\{Z \leq z\} = P\{Y \geq X - z\}$. The domain of $f(x, y)$ is $D = \{(x, y) \mid 0 < x < 1, y > 0\}$.

1. Case 1: $z < 0$

$$\begin{aligned} F_Z(z) &= \int_0^1 \left(\int_{x-z}^{\infty} \frac{1}{2}e^{-y/2} dy \right) dx \\ F_Z(z) &= \int_0^1 e^{-(x-z)/2} dx = e^{z/2} \left[-2e^{-x/2} \right]_0^1 \\ F_Z(z) &= 2e^{z/2}(1 - e^{-1/2}), \quad z < 0 \end{aligned}$$

2. Case 2: $0 \leq z < 1$ We use the complement $P\{Y < X - z\}$, where integration is over $z < x < 1$.

$$\begin{aligned} P\{Y < X - z\} &= \int_z^1 \left(\int_0^{x-z} \frac{1}{2}e^{-y/2} dy \right) dx \\ P\{Y < X - z\} &= \int_z^1 \left(1 - e^{-(x-z)/2} \right) dx = \left[x + 2e^{z/2}e^{-x/2} \right]_z^1 \\ P\{Y < X - z\} &= 2e^{(z-1)/2} - z - 1 \end{aligned}$$

Therefore:

$$F_Z(z) = 1 - P\{Y < X - z\} = z + 2 - 2e^{(z-1)/2}, \quad 0 \leq z < 1$$

3. Case 3: $z \geq 1$ Since $x - z < 0$ for all $x \in (0, 1)$, the condition $Y \geq X - z$ covers the entire domain D .

$$F_Z(z) = 1, \quad z \geq 1$$

4. Finding the Probability Density Function $f_Z(z)$ We differentiate $f_Z(z) = F'_Z(z)$:

1. Case 1: $z < 0$

$$f_Z(z) = \frac{d}{dz} \left[2e^{z/2}(1 - e^{-1/2}) \right] = (1 - e^{-1/2})e^{z/2}, \quad z < 0$$

2. Case 2: $0 < z < 1$

$$f_Z(z) = \frac{d}{dz} \left[z + 2 - 2e^{(z-1)/2} \right] = 1 - e^{(z-1)/2}, \quad 0 < z < 1$$

3. Case 3: $z > 1$

$$f_Z(z) = \frac{d}{dz}[1] = 0, \quad z > 1$$

3. Conclusion

The PDF of $Z = X - Y$ is:

$$f_Z(z) = \begin{cases} (1 - e^{-1/2})e^{z/2}, & z < 0 \\ 1 - e^{(z-1)/2}, & 0 \leq z < 1 \\ 0, & z \geq 1 \end{cases}$$

4 Chapter 4: Numerical Characteristics of Random Variables

4.1 Expectation (Mean)

The expectation $E(X)$ represents the probability reward.

- **Discrete Random Variable:**

$$E(X) = \sum x_i p_i$$

- **Continuous Random Variable:**

$$E(X) = \int_{-\infty}^{+\infty} x f(x) dx$$

4.1.1 Properties of Expectation

- **Linearity:**

$$E(c_0 + \sum c_i X_i) = c_0 + \sum c_i E(X_i)$$

- **Multiplication (Independence):** If X_i are independent, then:

$$E(\prod X_i) = \prod E(X_i)$$

4.2 Variance

Variance $D(X)$ or $Var(X)$ is a measure of the degree of deviation.

$$Var(X) = E[(X - E(X))^2] = E(X^2) - (E(X))^2$$

4.2.1 Properties of Variance

If X and Y are independent random variables:

$$Var(aX + bY + c) = a^2 Var(X) + b^2 Var(Y)$$

4.3 Covariance and Correlation Coefficient

4.3.1 Covariance

$$Cov(X, Y) = E[(X - E(X))(Y - E(Y))] = E(XY) - E(X)E(Y)$$

4.3.2 Correlation Coefficient

$$\rho_{XY} = \frac{Cov(X, Y)}{\sqrt{Var(X)Var(Y)}}$$

4.4 Expectation and Variance of Common Distributions

	$0-1$	$B(n, p)$	$P(\lambda)$	$U(a, b)$	$e(\lambda)$	$N(\mu, \sigma^2)$
$E(X)$	p	np	λ	$\frac{a+b}{2}$	$\frac{1}{\lambda}$	μ
$D(X)$	$p(1-p)$	$np(1-p)$	λ	$\frac{(b-a)^2}{12}$	$\frac{1}{\lambda^2}$	σ^2

4.5 Examples and Applications

4.5.1 Example 1: Variance of a Linear Combination

Given independent variables $X_1 \sim U(0, 6)$, $X_2 \sim N(0, 4)$, and $X_3 \sim P(3)$. Find $D(X_1 - 2X_2 + 3X_3)$.

Solution:

$$D(X_1) = \frac{(6-0)^2}{12} = 3, \quad D(X_2) = 4, \quad D(X_3) = 3$$

$$\begin{aligned} D(X_1 - 2X_2 + 3X_3) &= D(X_1) + (-2)^2 D(X_2) + (3)^2 D(X_3) \\ &= 3 + 4(4) + 9(3) = 3 + 16 + 27 = 46 \end{aligned}$$

4.5.2 Example 2: Finding $E(X^2)$ for a Binomial Variable

Given $X \sim B(10, 0.4)$. Find $E(X^2)$.

Solution:

$$\begin{aligned} E(X^2) &= D(X) + (E(X))^2 \\ E(X) &= 10(0.4) = 4, \quad D(X) = 10(0.4)(0.6) = 2.4 \\ E(X^2) &= 2.4 + (4)^2 = 18.4 \end{aligned}$$

4.5.3 Example 3: Properties of Covariance

(1) $\text{Cov}(X - Y, Y)$ (2) ρ_{XY}

X			Y			XY			
0	1	2	0	1	2	0	1	2	4
$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{7}{12}$	$\frac{1}{3}$	0	$\frac{1}{12}$

(1)

$$\begin{aligned} \text{Cov}(X - Y, Y) &= \text{Cov}(X, Y) - \text{Cov}(Y, Y) \\ &= E(XY) - E(X)E(Y) - D(Y) \\ &= E(XY) - E(X)E(Y) - \underbrace{(E(Y^2) - E^2(Y))}_{\text{Variance } D(Y)} \end{aligned}$$

(2) 0, Uncorrelated

4.5.4 Example 4: Correlation for Joint PDF

Given the joint PDF $f(x, y) = \begin{cases} \frac{3}{4}x^2y & 0 \leq x \leq 2, 0 \leq y \leq 1 \\ 0 & \text{others} \end{cases}$. Find ρ_{XY} .

Solution:

$$\begin{aligned} E(XY) &= \int_0^1 dy \int_0^2 xy \left(\frac{3}{4}x^2y\right) dx = 1 \\ E(X) &= \int_0^1 dy \int_0^2 x \left(\frac{3}{4}x^2y\right) dx = \frac{3}{2} \\ E(Y) &= \int_0^2 dx \int_0^1 y \left(\frac{3}{4}x^2y\right) dy = \frac{2}{3} \end{aligned}$$

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y) = 1 - \left(\frac{3}{2}\right)\left(\frac{2}{3}\right) = 0$$

Since $\text{Cov}(X, Y) = 0$, the variables are uncorrelated, thus $\rho_{XY} = 0$.

4.6 Chebyshev's Inequality

Chebyshev's inequality provides a bound on the probability deviation from the mean:

$$P(|X - E(X)| < \epsilon) \geq 1 - \frac{D(X)}{\epsilon^2}$$

(The complementary form: $P(|X - E(X)| \geq \epsilon) \leq \frac{D(X)}{\epsilon^2}$)

4.6.1 Example: Applying Chebyshev's Inequality

Given $E(X) = 100$ and $D(X) = 50$. Find the lower bound for $P\{80 < X < 120\}$.

Solution: We seek $P(80 < X < 120) = P(|X - 100| < 20)$. Here, $\epsilon = 20$.

$$\begin{aligned} P(|X - E(X)| < 20) &\geq 1 - \frac{D(X)}{\epsilon^2} = 1 - \frac{50}{20^2} \\ &= 1 - \frac{50}{400} = 1 - \frac{1}{8} = \frac{7}{8} \end{aligned}$$

The lower bound is $\frac{7}{8}$.

5 Chapter 5: Limit Theorems

5.1 Chebyshev's Inequality

Chebyshev's inequality provides a bound on the probability deviation from the mean:

$$P(|X - E(X)| \geq \epsilon) \leq \frac{Var(X)}{\epsilon^2}$$

or the complementary form:

$$P(|X - E(X)| < \epsilon) \geq 1 - \frac{Var(X)}{\epsilon^2}$$

5.2 Laws of Large Numbers (LLN)

5.2.1 Khinchine's Law of Large Numbers

Let X_1, X_2, \dots, X_n be a sequence of independent and identically distributed (i.i.d.) random variables, and $E(X_i) = \mu < \infty$. Then the sample mean converges in probability to the expectation:

$$\frac{1}{n} \sum_{i=1}^n X_i \rightarrow \mu, \quad \text{as } n \rightarrow \infty$$

5.3 Central Limit Theorem (CLT)

Let X_1, X_2, \dots, X_n be a sequence of independent and identically distributed (i.i.d.) random variables, with $E(X_i) = \mu$ and $Var(X_i) = \sigma^2$. If n is large, the distribution of the sum and the sample mean can be approximated by a Normal distribution:

- **Distribution of the Sum:**

$$\sum_{i=1}^n X_i \sim N(n\mu, n\sigma^2)$$

- **Distribution of the Sample Mean:**

$$\frac{1}{n} \sum_{i=1}^n X_i \sim N\left(\mu, \frac{\sigma^2}{n}\right)$$

The standardized random variable Z approaches the standard normal distribution:

$$Z = \frac{X - \mu}{\sigma} \sim N(0, 1)$$

5.4 Applications of the Central Limit Theorem

5.4.1 Example 1: Sum of Exponential Lifetimes

Electronic components follow an Exponential distribution with a mean life of 100 hours. Randomly select 16 independent components. Find the probability that the total lifetime ($\sum X_i$) is greater than 1920 hours.

Solution: For an Exponential distribution $X \sim E(\lambda)$, we have $E(X) = 1/\lambda$ and $D(X) = 1/\lambda^2$. Given $E(X) = 100$, so $D(X) = 100^2 = 10000$. By the CLT, the sum of $n = 16$ independent random variables is approximately:

$$\sum_{i=1}^{16} X_i \sim N(n\mu, n\sigma^2) = N(16 \cdot 100, 16 \cdot 100^2) = N(1600, 160000)$$

We want to find $P(\sum X_i > 1920)$.

$$P(\sum X_i > 1920) = 1 - P(\sum X_i \leq 1920)$$

Standardizing the variable:

$$\begin{aligned} P(\sum X_i \leq 1920) &\approx P\left(\frac{\sum X_i - 1600}{\sqrt{160000}} \leq \frac{1920 - 1600}{\sqrt{160000}}\right) \\ &= P\left(Z \leq \frac{320}{400}\right) = P(Z \leq 0.8) \end{aligned}$$

Using the standard normal CDF $\Phi(z)$:

$$P(\sum X_i > 1920) \approx 1 - \Phi(0.8)$$

5.4.2 Example 2: Probability of Defects (Normal Approximation to Binomial)

100 components are produced, and each one has a 0.1 probability of being defective. Find the probability that the number of defective components is less than 15.

Solution: Let X_i be 1 if the i -th component is defective, and 0 otherwise. $X_i \sim B(1, 0.1)$. The total number of defective components is $S = \sum_{i=1}^{100} X_i$.

$$E(S) = np = 100 \cdot 0.1 = 10$$

$$D(S) = np(1 - p) = 100 \cdot 0.1 \cdot 0.9 = 9$$

Since $n = 100$ is large, we can use the CLT approximation: $S \sim N(10, 9)$. We want to find $P(S < 15)$. Standardizing the variable (without continuity correction):

$$\begin{aligned} P(S < 15) &= P\left(\frac{S - 10}{\sqrt{9}} < \frac{15 - 10}{\sqrt{9}}\right) \\ &= P\left(Z < \frac{5}{3}\right) = \Phi(1.67) \end{aligned}$$

The probability is approximately $\Phi(1.67) \approx 0.9525$.

6 Chapter 6: Samples and Sampling Distributions

6.1 Sample Statistics

Let X_1, X_2, \dots, X_n be a sample of size n from a population.

6.1.1 Definitions of Common Sample Statistics

- **Sample Mean:**

$$\bar{X} = \frac{1}{n} \sum X_i$$

- **Sample Variance:**

$$S^2 = \frac{1}{n-1} \sum (X_i - \bar{X})^2$$

- **Sample k -th Moment (Moment about the Origin):**

$$A_k = \frac{1}{n} \sum X_i^k$$

- **Sample Central k -th Moment:**

$$B_k = \frac{1}{n} \sum (X_i - \bar{X})^k$$

6.1.2 Estimation of Population Characteristics

When the population numerical characteristics are unknown, the following are used as their estimators:

- Sample mean \bar{X} estimates population mean $\mu = E(X)$.
- Sample variance S^2 estimates population variance $\sigma^2 = E(X - \mu)^2$.
- Sample k -th moment A_k estimates population k -th moment $\mu_k = E(X^k)$.
- Sample central k -th moment B_k estimates population central k -th moment $\nu_k = E(X - \mu)^k$.

6.2 Important Sampling Distributions

6.2.1 Distribution of the Sample Mean

If X_1, X_2, \dots, X_n are samples from $N(\mu, \sigma^2)$, then the sample mean follows a Normal distribution:

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$$

6.2.2 Chi-Square (χ^2) Distribution

- If $Y_1 \sim \chi^2(m)$ and $Y_2 \sim \chi^2(n)$ are independent, then:

$$Y_1 + Y_2 \sim \chi^2(m + n)$$

6.2.3 t-Distribution (Student's t-Distribution)

- If $X \sim N(0, 1)$ and $Y \sim \chi^2(n)$ are independent, then the T -statistic follows a t -distribution with n degrees of freedom, $T \sim t(n)$:

$$T = \frac{X}{\sqrt{Y/n}}$$

- The square of a t -distributed variable is an F -distributed variable:

$$t^2 \sim F(1, n)$$

6.2.4 F-Distribution

- If $X \sim \chi^2(n_1)$ and $Y \sim \chi^2(n_2)$ are independent, then the F -statistic follows an F -distribution with (n_1, n_2) degrees of freedom, $F \sim F(n_1, n_2)$:

$$F = \frac{X/n_1}{Y/n_2}$$

- The inverse of an F -distributed variable is also F -distributed:

$$\frac{1}{F} \sim F(n_2, n_1)$$

7 Chapter 7: Parameter Estimation

7.1 Point Estimation

7.1.1 Method of Moments

The method of moments estimates the unknown parameters $\theta_1, \dots, \theta_k$ by equating the first k population moments to the corresponding sample moments.

1. **Population l -th Moment (Moment about the Origin):**

$$\mu_l = E(X^l) = \int_{-\infty}^{+\infty} X^l f(x; \theta_1, \dots, \theta_k) dx$$

2. Sample l -th Moment (Moment about the Origin):

$$A_l = \frac{1}{n} \sum_{i=1}^n X_i^l$$

3. **Solving for Estimators:** Set $A_l = \mu_l(\theta_1, \dots, \theta_k)$ and solve the system of equations for the parameters $\hat{\theta}_1, \dots, \hat{\theta}_k$.

Example: Moment Estimator for a Distribution Parameter Given the PDF:

$$f(x) = \begin{cases} \sqrt{\theta} x^{\sqrt{\theta}-1} & 0 < x < 1 \\ 0 & \text{others} \end{cases}$$

Find the moment estimator $\hat{\theta}$.

Solution: Equate the first population moment $E(X)$ to the first sample moment \bar{X} :

$$E(X) = \int_0^1 x \left(\sqrt{\theta} x^{\sqrt{\theta}-1} \right) dx = \sqrt{\theta} \int_0^1 x^{\sqrt{\theta}} dx = \frac{\sqrt{\theta} X^{\sqrt{\theta}+1}}{\sqrt{\theta}+1} \Big|_0^1 = \frac{\sqrt{\theta}}{\sqrt{\theta}+1}$$

Setting $E(X) = \bar{X}$:

$$\bar{X} = \frac{\sqrt{\theta}}{\sqrt{\theta}+1}$$

Solving for $\sqrt{\theta}$:

$$\begin{aligned} \bar{X}(\sqrt{\theta}+1) &= \sqrt{\theta} \implies \bar{X}\sqrt{\theta} + \bar{X} = \sqrt{\theta} \\ \bar{X} &= \sqrt{\theta}(1 - \bar{X}) \implies \sqrt{\theta} = \frac{\bar{X}}{1 - \bar{X}} \end{aligned}$$

The moment estimator is:

$$\hat{\theta} = \left(\frac{\bar{X}}{1 - \bar{X}} \right)^2$$

7.1.2 Maximum Likelihood Estimation (MLE)

1. Construct the Likelihood Function $L(\theta)$:

$$L(\theta) = \prod_{i=1}^n f(x_i; \theta) \quad (\text{Continuous Case})$$

$$L(\theta) = \prod_{i=1}^n P(X = x_i; \theta) \quad (\text{Discrete Case})$$

2. Take the Log-Likelihood Function $\ln L(\theta)$:

3. Calculate the First Derivative:

$$\frac{d \ln L}{d \theta}$$

4. Solve the Likelihood Equation: Set $\frac{d \ln L}{d \theta} = 0$ and solve for $\hat{\theta}$.

7.2 Properties of Estimators

7.2.1 Unbiasedness and Efficiency

Example: Comparing Unbiased Estimators Let X_1 and X_2 be independent random samples from $N(\mu, 1)$. Consider two estimators for μ :

$$\hat{\mu}_1 = \frac{1}{3}X_1 + \frac{2}{3}X_2 \quad \text{and} \quad \hat{\mu}_2 = \frac{1}{4}X_1 + \frac{3}{4}X_2$$

1. **Check for Unbiasedness:**

$$E(\hat{\mu}_1) = E\left(\frac{1}{3}X_1 + \frac{2}{3}X_2\right) = \frac{1}{3}E(X_1) + \frac{2}{3}E(X_2) = \frac{1}{3}\mu + \frac{2}{3}\mu = \mu$$

$$E(\hat{\mu}_2) = E\left(\frac{1}{4}X_1 + \frac{3}{4}X_2\right) = \frac{1}{4}E(X_1) + \frac{3}{4}E(X_2) = \frac{1}{4}\mu + \frac{3}{4}\mu = \mu$$

Both estimators are unbiased.

2. **Compare Efficiency (Variance):**

$$D(\hat{\mu}_1) = D\left(\frac{1}{3}X_1 + \frac{2}{3}X_2\right) = \left(\frac{1}{3}\right)^2 D(X_1) + \left(\frac{2}{3}\right)^2 D(X_2) = \frac{1}{9}(1) + \frac{4}{9}(1) = \frac{5}{9}$$

$$D(\hat{\mu}_2) = D\left(\frac{1}{4}X_1 + \frac{3}{4}X_2\right) = \left(\frac{1}{4}\right)^2 D(X_1) + \left(\frac{3}{4}\right)^2 D(X_2) = \frac{1}{16}(1) + \frac{9}{16}(1) = \frac{10}{16} = \frac{5}{8}$$

Since $D(\hat{\mu}_1) = 5/9 \approx 0.556$ and $D(\hat{\mu}_2) = 5/8 = 0.625$, we have $D(\hat{\mu}_1) < D(\hat{\mu}_2)$. Therefore, $\hat{\mu}_1$ is a **better** (more efficient) estimator.

7.3 Interval Estimation (Confidence Intervals)

For a confidence level of $1 - \alpha$, the confidence interval is calculated using the following formulas, typically for a Normal population $X \sim N(\mu, \sigma^2)$.

7.3.1 Confidence Interval for the Population Mean μ

- **σ^2 is Known:** Uses the Z (Standard Normal) distribution.

$$\left(\bar{X} \pm \frac{\sigma}{\sqrt{n}} Z_{\alpha/2}\right)$$

- **σ^2 is Unknown:** Uses the t (Student's t) distribution with $n - 1$ degrees of freedom.

$$\left(\bar{X} \pm \frac{S}{\sqrt{n}} t_{\alpha/2}(n-1)\right)$$

7.3.2 Confidence Interval for the Population Variance σ^2

- **μ is Known:** Uses the χ^2 (Chi-square) distribution with n degrees of freedom.

$$\left(\frac{\sum_{i=1}^n (X_i - \mu)^2}{\chi_{\alpha/2}^2(n)}, \frac{\sum_{i=1}^n (X_i - \mu)^2}{\chi_{1-\alpha/2}^2(n)}\right)$$

- **μ is Unknown:** Uses the χ^2 distribution with $n - 1$ degrees of freedom, where $(n - 1)S^2 = \sum (X_i - \bar{X})^2$.

$$\left(\frac{(n-1)S^2}{\chi_{\alpha/2}^2(n-1)}, \frac{(n-1)S^2}{\chi_{1-\alpha/2}^2(n-1)}\right)$$

Example: Calculating Confidence Intervals

$X \sim N(\mu, \sigma^2)$, sample size $n = 6$. Given observed sample variance S^2 corresponding to $S = 0.226$. The confidence level is $1 - \alpha = 95\%$.

1. **Confidence Interval for μ (σ^2 unknown):** The calculation uses the t -distribution with $n - 1 = 5$ degrees of freedom and $t_{0.025}(5)$.

$$\left(\bar{X} \pm \frac{S}{\sqrt{n}} t_{\alpha/2}(n-1)\right) = \left(14.95 \pm \frac{0.26}{\sqrt{6}} t_{0.025}(5)\right)$$

2. **Confidence Interval for σ^2 (μ unknown):** The calculation uses the χ^2 distribution with $n - 1 = 5$ degrees of freedom.

$$\left(\frac{5S^2}{\chi_{\alpha/2}^2(5)}, \frac{5S^2}{\chi_{1-\alpha/2}^2(5)}\right)$$

8 Chapter 8: Hypothesis Testing

8.1 Concepts and Structure of Hypothesis Testing

- Assumption \rightarrow Sampling \rightarrow Decision (Small probability of refusal)
- Left-sided test $H_0 : \theta \geq \theta_0, H_1 : \theta < \theta_0$
- Right-sided test $H_0 : \theta \leq \theta_0, H_1 : \theta > \theta_0$
- Two-sided test $H_0 : \theta = \theta_0, H_1 : \theta \neq \theta_0$

Table 1: Decision Table

Decision	H_0 True	H_0 False
Reject H_0	Type I Error (α)	Correct Decision
Accept H_0	Correct Decision	Type II Error (β)

8.2 Inference for the Population Mean (μ)

1. **Case 1: Population Variance σ^2 is Known (σ^2 known)**

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1)$$

Hypothesis Test (Critical Region):

$$|Z| \geq Z_{\alpha/2}$$

2. **Case 2: Population Variance σ^2 is Unknown (σ^2 unknown)**

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \sim t(n-1)$$

Hypothesis Test (Critical Region):

$$|T| \geq t_{\alpha/2}(n-1)$$

8.3 Inference for the Population Variance (σ^2)

1. **Case 3: Population Mean μ is Known (μ known)**

$$W = \frac{\sum_{i=1}^n (X_i - \mu)^2}{\sigma_0^2} \sim \chi^2(n)$$

Hypothesis Test (Confidence Interval Lower/Upper Bounds for σ^2):

$$\chi_{1-\alpha/2}^2(n) \leq W \leq \chi_{\alpha/2}^2(n)$$

2. **Case 4: Population Mean μ is Unknown (μ unknown)**

$$V = \frac{(n-1)S^2}{\sigma_0^2} \sim \chi^2(n-1)$$

Hypothesis Test (Confidence Interval Lower/Upper Bounds for σ^2):

$$\chi_{1-\alpha/2}^2(n-1) \leq V \leq \chi_{\alpha/2}^2(n-1)$$

8.4 Examples

8.4.1 Example: Mean μ Test (Variance Known)

- **Problem:** The weight of salt in each bag $X \sim N(500, 15^2)$ grams. A sample of $n = 25$ bags is taken, with a sample mean $\bar{x} = 511.22$ grams. Is this normal? Use $\alpha = 0.05$.
- **Hypothesis:** $H_0 : \mu = \mu_0 = 500, H_1 : \mu \neq \mu_0$.
- **Test Statistic:** $Z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1)$.
- **Critical Value:** $Z_{\frac{\alpha}{2}} = 1.96$.
- **Rejection Region (W):** $W = (-\infty, -1.96) \cup (1.96, +\infty)$.
- **Calculation:**

$$Z = \frac{511.22 - 500}{15/\sqrt{25}} = 3.74$$

- **Decision:** Since $3.74 \in W$, reject H_0 (conclude H_1). The process is not normal.

8.4.2 Example: Variance σ^2 Test

- **Problem:** Required $\sigma_0^2 = 0.0004$. A sample of $n = 20$ bags has a sample variance $S = 0.003$. Is the standard met? Let $\alpha = 0.05$.
- **Hypothesis:** $H_0 : \sigma^2 = \sigma_0^2 = 0.0004; H_1 : \sigma^2 \neq \sigma_0^2$.
- **Test Statistic:** $\chi^2 = \frac{(n-1)S^2}{\sigma_0^2} \sim \chi^2(n-1)$.
- **Rejection Region (W):** $W = (0, \chi_{1-\frac{\alpha}{2}}^2(n-1)) \cup (\chi_{\frac{\alpha}{2}}^2(n-1), +\infty)$.
- **Critical Values ($n-1 = 19, \alpha = 0.05$):**

$$- \chi_{0.975}^2(19) = 8.907$$

$$- \chi_{0.025}^2(19) = 32.852$$

- **Rejection Region (W):** $W = (0, 8.907) \cup (32.852, +\infty)$.
- **Calculation:**

$$\chi^2 = \frac{19 \times 0.003}{0.0004} = 142.5$$

- **Decision:** Since $142.5 \in W$, reject H_0 (conclude H_1). The standard is not met.