

Information Electronics Physics

Fundamentals

Lecture Notes

Abstract

This document serves as a summarized translation of the lecture notes for "Information Electronics Physics Fundamentals". It covers Solid State Types, Quantum Mechanics Basics, Band Theory, Carrier Transport, and PN Junction/Semiconductor Device Physics.

Contents

1	Solid State Types and Crystal Structure	3
1.1	Types of Solids	3
1.2	Space Lattice	3
1.2.1	Unit Cells	3
1.3	Basic Crystal Structures	3
1.3.1	Diamond Structure	4
1.4	Crystal Planes and Miller Indices	4
1.5	Defects and Impurities	4
2	Quantum Mechanics Basics	6
2.1	Wave-Particle Duality	6
2.2	The Schrödinger Equation	6
2.2.1	Physical Meaning of ψ	6
2.3	Applications	6
2.3.1	Infinite Potential Well	6
2.3.2	Potential Barrier and Tunneling	6
2.4	Atomic Theory	7
3	Band Theory and Carrier Transport	8
3.1	Energy Band Formation	8
3.2	Classifications of Solids	8

3.3	Effective Mass and Holes	8
3.4	Carrier Concentration	9
3.4.1	Fermi-Dirac Distribution	9
3.4.2	Equilibrium Concentrations	9
3.5	Doping	9
3.6	Transport Properties	9
4	PN Junctions and Semiconductor Devices	10
4.1	The PN Junction at Equilibrium	10
4.2	Biasing the PN Junction	10
4.3	Junction Capacitance	10
4.4	Breakdown Mechanisms	11
4.5	Metal-Semiconductor Contacts	11
4.6	Heterojunctions	11
5	Optical Properties of Solids and Optoelectronics	11
5.1	Optical Absorption	11
5.1.1	Absorption Coefficient	12
5.1.2	Direct and Indirect Transitions	12
5.2	Photovoltaic Effect and Solar Cells	12
5.2.1	Characteristics	13
5.2.2	Advanced Structures	13
5.3	Photodetectors	13
5.3.1	Non-Gain Detectors (Photodiodes)	13
5.3.2	Gain Detectors (Avalanche Photodiodes - APD)	14
5.4	Light Emission (LEDs and Lasers)	14
5.4.1	Light Emitting Diodes (LED)	14
5.4.2	Laser Diodes (LD)	14
6	Magnetoelectronics	15
6.1	Magnetic Properties of Materials	15
6.2	Ferromagnetism and Hysteresis	15
7	Superconductivity	16
7.1	Basic Properties	16
7.2	Critical Parameters	16
7.3	Types of Superconductors	16
7.4	Theoretical Models	16
7.4.1	London Equations	16
7.4.2	BCS Theory (Bardeen-Cooper-Schrieffer)	16
7.5	Tunneling and Josephson Effect	17

1 Solid State Types and Crystal Structure

1.1 Types of Solids

Solids are classified based on the arrangement of their atoms:

- **Amorphous:** Short-range order only (no long-range periodicity).
- **Polycrystalline:** Composed of many small crystallites or grains.
- **Single Crystal:** Long-range order extends throughout the entire material. This is crucial for semiconductor devices.

1.2 Space Lattice

A crystal structure is described by:

$$\text{Crystal Structure} = \text{Lattice} + \text{Basis} \quad (1)$$

- **Lattice:** A set of points (grid) where every point has an identical environment.
- **Basis:** An atom or group of atoms attached to every lattice point.
- **Bravais Lattices:** 14 unique lattice structures in 3D space where all lattice points are equivalent.

1.2.1 Unit Cells

- **Primitive Cell:** The smallest volume cell that can reproduce the lattice (contains 1 lattice point). Vectors: $\vec{a}, \vec{b}, \vec{c}$.
- **Unit Cell:** A convenient repeating volume (may contain > 1 lattice point) chosen to display symmetry.
- **Lattice Constant (a):** The physical dimension of the unit cell side length. Typical magnitude: 10^{-8} cm or 1 Å.

1.3 Basic Crystal Structures


Common structures in cubic systems:

1. **Simple Cubic (SC):** Atoms at corners only.
2. **Body-Centered Cubic (BCC):** Atoms at corners + 1 atom in the center.
3. **Face-Centered Cubic (FCC):** Atoms at corners + atoms at the center of each face.

1.3.1 Diamond Structure

Crucial for Silicon (Si) and Germanium (Ge).

- Can be visualized as two interpenetrating FCC sub-lattices shifted by a diagonal vector $(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$.
- Each atom has 4 nearest neighbors (tetrahedral bonding).
- Nearest neighbor distance: $\frac{\sqrt{3}}{4}a$.



1.4 Crystal Planes and Miller Indices

To determine the Miller indices (hkl) of a plane:

1. Find intercepts on axes in terms of lattice constants (e.g., p, q, r).
2. Take reciprocals $(\frac{1}{p}, \frac{1}{q}, \frac{1}{r})$.
3. Reduce to the smallest integers.

Note: For cubic systems, direction $[hkl]$ is perpendicular to plane (hkl) .

1.5 Defects and Impurities

Real crystals deviate from perfect periodicity.

- **Thermal Vibration:** Atoms vibrate around equilibrium positions.
- **Point Defects:**
 - *Vacancy:* An atom is missing from a lattice site.
 - *Interstitial:* An atom is squeezed into a void between sites.
 - *Frenkel Defect:* Pair of Vacancy + Interstitial (atom moves to interstitial site).
 - *Schottky Defect:* Atom moves from interior to the surface, leaving a vacancy. Requires less energy than Frenkel.
- **Impurities (Doping):**

- *Substitutional*: Impurity replaces a host atom.
- *Interstitial*: Impurity sits in voids.
- *Ion Implantation*: High-energy method to introduce impurities (precise but damages lattice).
- *Diffusion*: Thermal movement from high to low concentration.

2 Quantum Mechanics Basics

2.1 Wave-Particle Duality

- **Energy Quanta:** $E = h\nu = \hbar\omega$.
- **Momentum:** $p = \frac{h}{\lambda} = \hbar k$.
- **Heisenberg Uncertainty Principle:** $\Delta x \Delta p \geq \frac{\hbar}{2}$.

2.2 The Schrödinger Equation

The fundamental equation governing the behavior of electrons.

$$-\frac{\hbar^2}{2m}\nabla^2\Psi(x,t) + U(x)\Psi(x,t) = j\hbar\frac{\partial\Psi(x,t)}{\partial t} \quad (2)$$

For a time-independent potential $U(x)$, the wavefunction separates into spatial and temporal parts. The time-independent form is:

$$\frac{d^2\psi(x)}{dx^2} + \frac{2m}{\hbar^2}(E - U(x))\psi(x) = 0 \quad (3)$$

2.2.1 Physical Meaning of ψ

- $\psi(x)$ itself is a complex quantity with no direct physical meaning.
- $|\psi(x)|^2 = \psi(x)\psi^*(x)$ represents the **probability density** of finding the electron at position x .
- Normalization condition: $\int_{-\infty}^{\infty} |\psi(x)|^2 dx = 1$.

2.3 Applications

2.3.1 Infinite Potential Well

Electron confined in region $0 < x < a$ with $U(x) = 0$, and $U(x) = \infty$ elsewhere.

- Solution: $\psi_n(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi x}{a}\right)$.
- Energy is quantized: $E_n = \frac{\hbar^2\pi^2 n^2}{2ma^2}$ where $n = 1, 2, 3, \dots$

2.3.2 Potential Barrier and Tunneling

When an electron with energy $E < U_0$ encounters a potential barrier of finite width:

- Classically: Reflection $R = 1$.

- **Quantum Mechanically:** The wavefunction decays exponentially inside the barrier but is non-zero on the other side.
- **Tunneling Effect:** There is a finite probability of transmission.

2.4 Atomic Theory

For a single electron atom (Hydrogen model), states are defined by quantum numbers:

- **Principal (n):** Energy shell ($n = 1, 2, \dots$).
- **Orbital (l):** Angular momentum ($l = 0, 1, \dots, n - 1$).
- **Magnetic (m):** Orientation ($m = -l, \dots, +l$).
- **Spin (s):** Intrinsic property ($s = \pm 1/2$).

3 Band Theory and Carrier Transport

3.1 Energy Band Formation


As atoms are brought together to form a crystal, discrete atomic energy levels split into continuous bands due to the interaction of wavefunctions (Pauli Exclusion Principle).

- **Bloch Theorem:** In a periodic potential, the wavefunction is a plane wave modulated by a periodic function: $\psi(x) = u(x)e^{jkx}$.
- **Kronig-Penney Model:** Uses a periodic square well potential to mathematically demonstrate the existence of allowed energy bands and forbidden gaps.

3.2 Classifications of Solids

Based on band structure at 0K:

- **Insulator:** Filled valence band, empty conduction band, large band gap (E_g).
- **Semiconductor:** Similar to insulators but small E_g . Electrons can be thermally excited.
- **Metal:** Partially filled band or overlapping conduction/valence bands.



3.3 Effective Mass and Holes

- Electrons in a crystal respond to external forces as if they had a mass m^* , different from vacuum mass m_0 .
- $m^* = \hbar^2 \left(\frac{d^2 E}{dk^2} \right)^{-1}$. Curvature of the band determines mass.
- **Hole:** A missing electron in the valence band behaves like a positively charged particle with positive effective mass.

3.4 Carrier Concentration

3.4.1 Fermi-Dirac Distribution

The probability that a state at energy E is occupied:

$$f(E) = \frac{1}{1 + \exp\left(\frac{E - E_F}{k_B T}\right)} \quad (4)$$

Where E_F is the Fermi Level. At $T = 0\text{K}$, $f(E) = 1$ for $E < E_F$ and 0 for $E > E_F$.

3.4.2 Equilibrium Concentrations

- Electron conc. (n_0): $n_0 = N_c \exp\left(-\frac{E_c - E_F}{k_B T}\right)$
- Hole conc. (p_0): $p_0 = N_v \exp\left(-\frac{E_F - E_v}{k_B T}\right)$
- **Law of Mass Action:** $n_0 p_0 = n_i^2 = N_c N_v \exp\left(-\frac{E_g}{k_B T}\right)$. This holds independent of doping in equilibrium.
- **Intrinsic Fermi Level (E_i):** Located near the middle of the band gap.

3.5 Doping

- **Donors (N-type):** Group V elements (e.g., P, As in Si). Add electrons. Level E_D near E_c .
- **Acceptors (P-type):** Group III elements (e.g., B in Si). Create holes. Level E_A near E_v .
- **Temperature Regimes:**
 1. *Freeze-out (Low T):* Dopants not fully ionized.
 2. *Extrinsic (Medium T):* Constant carrier density determined by doping ($n \approx N_D$).
 3. *Intrinsic (High T):* Thermally generated carriers dominate ($n_i > N_D$).

3.6 Transport Properties

- **Drift:** Motion caused by electric field. $v_d = \mu E$. (μ = mobility).
- **Resistivity:** $\rho = \frac{1}{\sigma} = \frac{1}{q(n\mu_n + p\mu_p)}$.
- **Hall Effect:** Used to measure carrier type and concentration.
 - N-type: Hall coefficient $R_H < 0$.
 - P-type: Hall coefficient $R_H > 0$.


4 PN Junctions and Semiconductor Devices

4.1 The PN Junction at Equilibrium

When P-type and N-type materials join:

- **Diffusion:** Holes diffuse $P \rightarrow N$, Electrons diffuse $N \rightarrow P$.
- **Space Charge Region (SCR):** Leaving behind ionized dopants (N_A^- in P-side, N_D^+ in N-side) creates a built-in electric field preventing further diffusion.
- **Energy Bands:** The Fermi level must be constant ($E_{FP} = E_{FN}$). This causes bands to bend.
- **Built-in Potential (V_{bi} or V_D):**

$$V_{bi} = \frac{k_B T}{q} \ln \left(\frac{N_A N_D}{n_i^2} \right) \quad (5)$$



4.2 Biasing the PN Junction

- **Forward Bias ($V_F > 0$):** P connected to (+). Barrier lowers. Diffusion current dominates.
- **Reverse Bias ($V_R > 0$):** P connected to (-). Barrier increases. Current is negligible (leakage).
- **Shockley Diode Equation:**

$$I = I_s \left[\exp \left(\frac{qV}{k_B T} \right) - 1 \right] \quad (6)$$

4.3 Junction Capacitance

- **Barrier/Junction Capacitance (C_j):** Dominant in reverse bias. Acts like a parallel plate capacitor with changing width W . Used in **Varactor Diodes**.
- **Diffusion Capacitance (C_d):** Dominant in forward bias due to stored minority carriers.

4.4 Breakdown Mechanisms

Occurs at high reverse voltage.

1. **Zener Breakdown:** Tunneling of electrons from valence band to conduction band in heavily doped junctions.
2. **Avalanche Breakdown:** Impact ionization. Carriers gain enough kinetic energy to knock bound electrons free, creating pairs.

4.5 Metal-Semiconductor Contacts

- **Schottky Contact:** Rectifying (diode-like). Occurs when work functions differ such that a barrier forms ($\Phi_m > \Phi_s$ for n-type).
- **Ohmic Contact:** Linear I-V curve. Achieved by heavy doping (tunneling) or matching work functions. Essential for connecting devices to circuits.

4.6 Heterojunctions

Junctions between two different semiconductor materials (e.g., GaAs / AlGaAs).

- **Band Offsets:** Discontinuities in conduction (ΔE_c) and valence (ΔE_v) bands.
- **Quantum Wells:** Can trap carriers in 2D layers (2D Electron Gas).
- Applications: High Electron Mobility Transistors (HEMTs), Lasers.

5 Optical Properties of Solids and Optoelectronics

5.1 Optical Absorption

When light interacts with a solid, photons may be absorbed if their energy matches available energy states in the material.

- **Condition for Fundamental Absorption:** $h\nu \geq E_g$.
- **Process:** An electron in the valence band absorbs a photon and transitions to the conduction band, creating an electron-hole pair.
- **Transparency:** If $h\nu < E_g$, the material is typically transparent (ignoring impurities or phonon absorption).

5.1.1 Absorption Coefficient

The intensity of light $I(x)$ decays as it propagates through the material according to the Beer-Lambert Law:

$$I(x) = I_0 e^{-\alpha x} \quad (7)$$


Where α is the absorption coefficient.

- Relation to complex refractive index $\tilde{n} = n + jk$: $\alpha = \frac{4\pi k}{\lambda_0}$.
- **Penetration Depth:** The distance $1/\alpha$ where intensity drops to $1/e$.

5.1.2 Direct and Indirect Transitions

Conservation of energy and crystal momentum ($\hbar k$) must be satisfied.

1. **Direct Transition:** The conduction band minimum and valence band maximum occur at the same wavevector k (usually $k = 0$).
 - Momentum conservation: $k_f \approx k_i$ (since photon momentum is negligible).
 - Example materials: GaAs, InP (efficient for LEDs/Lasers).
2. **Indirect Transition:** The band edges are at different k values.
 - Requires a **phonon** (lattice vibration) to conserve momentum: $E_f = E_i + \hbar\nu \pm E_{\text{phonon}}$ and $\vec{k}_f = \vec{k}_i + \vec{q}_{\text{phonon}}$.
 - Less efficient process. Example material: Si, Ge.



5.2 Photovoltaic Effect and Solar Cells

Solar cells operate on the photovoltaic effect within a PN junction.

- **Mechanism:** Photogenerated electron-hole pairs in the depletion region are separated by the built-in electric field (Electrons \rightarrow N-side, Holes \rightarrow P-side).

- This accumulation creates a forward voltage (photovoltage) opposing the built-in potential.

5.2.1 Characteristics

- **Short Circuit Current (I_{SC}):** Current when $V = 0$. Proportional to light intensity.
- **Open Circuit Voltage (V_{OC}):** Voltage when $I = 0$.
- **Fill Factor (FF):** A measure of "squareness" of the I-V curve.

$$FF = \frac{I_m V_m}{I_{SC} V_{OC}} \quad (8)$$

- **Efficiency (η):** Ratio of max electrical power output to incident optical power (P_{in}).

$$\eta = \frac{P_{max}}{P_{in}} = \frac{FF \cdot I_{SC} \cdot V_{OC}}{P_{in}} \quad (9)$$

5.2.2 Advanced Structures

- **Heterojunctions:** Using a wide bandgap material (e.g., AlGaAs) on top of a narrower one (GaAs). The top layer acts as a "window," allowing high-energy photons to reach the active region without surface recombination.
- **PIN Structure (Amorphous Si):** An intrinsic (i) layer is sandwiched between p and n layers to extend the electric field region, improving carrier collection efficiency in materials with low diffusion lengths.

5.3 Photodetectors

5.3.1 Non-Gain Detectors (Photodiodes)

Operate in reverse bias to widen the depletion region and increase speed.

- **Quantum Efficiency (η):** Number of electron-hole pairs generated per incident photon.
- **Responsivity (R):** Photocurrent per unit optical power (A/W).
- **PIN Photodiode:** The intrinsic layer width W can be tailored to optimize quantum efficiency (absorption volume) and response speed (transit time vs. capacitance).

5.3.2 Gain Detectors (Avalanche Photodiodes - APD)

Operate at high reverse bias near breakdown.

- **Mechanism:** Primary photogenerated carriers gain sufficient kinetic energy to cause *impact ionization*, creating secondary pairs.
- **Result:** Internal gain M . High sensitivity but higher noise.

5.4 Light Emission (LEDs and Lasers)

5.4.1 Light Emitting Diodes (LED)

- Based on **Spontaneous Emission:** An electron falls to the valence band randomly, emitting a photon.
- Operates in Forward Bias (carrier injection).
- Efficiency depends on the ratio of radiative to non-radiative recombination rates.

5.4.2 Laser Diodes (LD)

Based on **Stimulated Emission:** An incoming photon triggers an electron to drop, emitting an identical photon (same phase, frequency, direction). **Three Conditions for Lasing:**

1. **Population Inversion:** More electrons in the excited state than the ground state ($N_2 > N_1$). Achieved by heavy doping (degenerate) and high injection ($E_{Fc} - E_{Fv} > E_g$).
2. **Optical Cavity:** Fabry-Perot resonator (cleaved facets) provides feedback/amplification.
3. **Threshold Condition:** Gain \geq Loss (absorption + mirror loss).

6 Magnetoelectronics

6.1 Magnetic Properties of Materials


Magnetization M relates to magnetic field H via susceptibility χ : $M = \chi H$.

1. **Diamagnetism** ($\chi < 0$): Weak repulsion. Induced magnetic moment opposes external field (Lenz's law at atomic scale). Present in all matter but often masked.
2. **Paramagnetism** ($\chi > 0$): Weak attraction. Randomly oriented permanent dipoles align partially with the field.
3. **Ferromagnetism** ($\chi \gg 0$): Strong attraction. Spontaneous magnetization exists below the Curie Temperature (T_c).

6.2 Ferromagnetism and Hysteresis

Ferromagnetic materials exhibit a memory effect.

- **Domain Theory:** Below T_c , microscopic regions (domains) are fully magnetized. An external field aligns these domains.
- **Hysteresis Loop:** The relationship between B and H is non-linear and irreversible.
 - **Saturation** (B_s): All domains aligned.
 - **Remanence** (B_r): Magnetization remaining at $H = 0$.
 - **Coercivity** (H_c): Reverse field needed to demagnetize ($B = 0$).
- **Origin:** Quantum mechanical **Exchange Interaction** (Weiss Mean Field). The Pauli Exclusion Principle forces parallel spins in certain orbital overlaps to lower electrostatic energy.



7 Superconductivity

7.1 Basic Properties

1. **Zero Resistance:** Below a critical temperature T_c , DC electrical resistance vanishes ($R = 0$).
2. **Meissner Effect (Perfect Diamagnetism):** A superconductor expels all magnetic flux from its interior ($\chi = -1$). $B = 0$ inside.

7.2 Critical Parameters

Superconductivity is destroyed if any of these exceed their critical value:

- Temperature ($T > T_c$).
- Magnetic Field ($H > H_c$). Relationship: $H_c(T) \approx H_c(0)[1 - (T/T_c)^2]$.
- Current Density ($J > J_c$).

7.3 Types of Superconductors

- **Type I:** Abrupt transition from superconducting state to normal state at H_c . (Mostly pure metals).
- **Type II:** Exists in a "Mixed State" (or Vortex State) between H_{c1} and H_{c2} . Magnetic flux penetrates in quantized vortices. (Allows for high-field magnets).

7.4 Theoretical Models

7.4.1 London Equations

Phenomenological description of electromagnetic fields in superconductors. Predicts the **Penetration Depth** (λ_L), the thin surface layer where supercurrents flow to screen the magnetic field.

7.4.2 BCS Theory (Bardeen-Cooper-Schrieffer)

Microscopic quantum theory.

- **Cooper Pairs:** Electrons (fermions) pair up to form bosons with opposite spin and momentum.
- **Mechanism:** Electron-Phonon Interaction. An electron distorts the lattice (attracts positive ions), creating a region of excess positive charge that attracts a second electron.

- **Energy Gap (2Δ):** A gap forms at the Fermi level, preventing small excitations (scattering) which causes resistance. $2\Delta \approx 3.5k_B T_c$.

7.5 Tunneling and Josephson Effect

- **Giaever Tunneling (SIS):** Tunneling of single electrons (quasiparticles) only occurs when voltage bias $eV \geq \Delta$ (or 2Δ for S-I-S), revealing the density of states gap.
- **Josephson Effect:** Tunneling of **Superconducting Cooper Pairs** through a thin insulating barrier.
 - **DC Josephson Effect:** Current flows with zero voltage drop ($V = 0$) up to a critical current I_c .
 - **AC Josephson Effect:** If a DC voltage V is applied, the current oscillates at frequency $f = 2eV/h$.