

CSCI 467 Problem Set 4

Teresa Liu

March 13, 2020

True/False Questions

1. For linear regression, when including interactions in our model, we can omit the main effects if the p-values associated with those effects are not statistically significant
 - a. **False:** *according to the Hierarchical Principle, you must include the main effect associated with interactions even if p-values are not significant*
2. KNN Regression is an example of a parametric model.
 - a. **False:** *KNN Regression is a non-parametric model.*
3. Discriminant Analysis is preferred for data sets that are well separated, since the parameter estimates for logistic regression are unstable.
 - a. **True:** *logistic regression does not perform well with well separated data sets.*
4. Naïve Bayes is robust to isolated noise samples, irrelevant attributes, and redundant attributes.
 - a. **False:** *Naïve Bayes is not robust to redundant attributes since this model assumes that features are independent in each class, and redundancy violates this assumption.*
5. Leave One Out Cross Validation (LOOCV) produces less bias and doesn't overestimate errors. However, this method is very computationally intensive, which is why K-Fold Cross Validation is more popular.
 - a. **True:** *Since LOOCV uses every data point individually as validation sets, it produces the lowest bias of the resampling methods, but this is too computationally difficult to perform, so K-Fold provides a compromise between bias and computational challenge.*
6. All algorithms perform equally when averaged over all problems
 - a. **True:** *This is the fundamental No Free Lunch Theorem*
7. KNN Regression is known as a Lazy Learning algorithm as opposed to an Eager Learning algorithm.
 - a. **True:** *KNN is known as lazy learning, since generalization beyond the training data is delayed until a query is made to the system, as opposed to in Eager Learning, where the system tries to generalize the training data before receiving queries.*
8. Dummy encoding and one-hot encoding are a commonly used method for converting a categorical input variable into continuous variables. With k-groups for a predictor, dummy encoding requires k coded variables and one-hot encoding requires k-1 coded variables.

- a. *False: The second statement should be flipped. With k -groups, dummy encoding requires $k-1$ coded variables and one-hot encoding requires k coded variables.*

Conceptual/Fundamental Midterm Exam Questions

1. Which of the following examples are considered unsupervised learning?
 - a. Taking a dataset of people's movie-watching history and transforming that data into a recommendations feature based on overlapping movie histories.
 - b. Anomaly detection when researching a medical data set for cancer cells, but we don't know what the anomaly looks like.
 - c. Using stock market data from the past 20 years to predict how the market will perform in the next 5 years.
 - d. Taking a large set of images of people and grouping faces together so that users can search photos by people's faces.
 - e. Training a factory robot to pick up and drop off boxes based on a set of rewards.
 - f. *Answer: A B D*
2. The following are indications that you should use discriminant analysis over logistic regression for a dataset.
 - a. When classes are well-separated.
 - b. If n is small and the distribution of the predictors X is approximately normal in each of the classes.
 - c. When we have more than 2 response classes and we wish to categorize data.
 - d. When you have many outliers in your data.
 - e. *Answer: all of the above*

Observe the following multiple linear regression model

$$\hat{y} = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3$$

- \hat{y} : the predicted number of sales in 1000's of dollars
 - x_1 : the number of ads put up on billboards
 - x_2 : the number of TV commercials that run on a channel
 - x_3 : a dummy variable with the ad containing a special deal
 - the value of coefficients: $\beta_0 = 1.1$, $\beta_1 = 5.2$, $\beta_2 = 1.3$, $\beta_3 = -0.2$
3. Which of the following statements are correct interpretations of β_1 ?
 - a. If the value of the number of ads put up on billboards increases by one unit, then the number of sales is predicted to increase by \$5,200 assuming that all other predictors remain fixed.
 - b. If the value of the number of ads put up on billboards decreases by 5.2, then the number of sales is predicted to decrease by \$1,000 assuming that all other predictors remain fixed.
 - c. If the value of the number of ads put up on billboards increases by one unit, then the number of sales is predicted to increase by \$5.20.

- d. If the value of the number of ads put up on billboards decreases by one unit, then the number of sales is predicted to decrease by \$5,200 assuming that all other predictors remain fixed.
 - e. **Answer:** A D
4. Which of the following are valid methods of selecting important variables in linear regression?
- a. Sideways Selection, in which you start with a model with $p/2$ variables, and add variable or remove a variable when the p-value associated with that variable is outside of an accepted range, refit the model, and continue until a stopping rule is reached.
 - b. Forward Selection, in which you begin with a null model, fit p simple linear regressions and add to the model the variable that results in the lowest RSS, and then repeat until a stopping rule is reached.
 - c. Backward Selection, in which you begin with all variables, remove the variable with the largest p-value, fit the new $p-1$ model, and continue until a stopping rule is reached.
 - d. LOOCV, in which you leave a variable out of the regression model and repeat N times to fit the model until you figure out which variable caused the most error, and then remove that variable, and repeat until stopping rule is reached.
 - e. **Answer:** B C