
Grade 12 GS

Probability ex 20

By Mr.Kassem Hodeib K.H



Exercise 20: "If then question or two actions:

Let C be a given fake coin such that the probability of having a head is three times having a tail.

Let U_1 and U_2 be two given urns such that:

U₁ contains 10 balls: 6 red and 4 yellow.

U₂ contains 10 balls: 5 red, 4 black and 1 green.

We throw the coin C. If we get a tail, then we pick up, randomly and **simultaneously**, two balls from urn U_1 ; otherwise, if we get a head, we select at random two balls from U_2 one after the other with replacement.

Consider the following events:


T: "The coin turns on a tail"

H: "The coin turns on a Head"

R: "The chosen balls are all red."

1) Show that
$$P(H) = \frac{3}{4}$$
 and $P(T) = \frac{1}{4}$.

$$P(H) + P(T) = 1$$
 Then: $3P(T) + P(T) = 1 \rightarrow 4P(T) = 1$
but $P(H) = 3P(T)$ $P(T) = \frac{1}{4}$ and $P(H) = 3P(T) = \frac{3}{4}$



 U_2

• Toss the coin C:

$$\frac{3}{4}$$
 H $\frac{/H}{}$

1) Calculate P(R/T), $P(R \cap T)$, and $P(R \cap H)$. Deduce P(R).

•
$$P(R/T) = \frac{C_6^2}{C_{10}^2} = \frac{1}{3}$$
 • $P(R \cap T) = P(R/T) \times P(T) = \frac{1}{3} \times \frac{1}{4} = \frac{1}{12}$

•
$$P(R \cap H) = P(R/H) \times P(H) = \frac{5}{10} \times \frac{5}{10} \times \frac{3}{4} = \frac{3}{16}$$

•
$$P(R) = P(R \cap T) + P(R \cap H) = \frac{1}{12} + \frac{3}{16} = \frac{13}{48}$$

2) The two selected balls are red. Calculate the probability that they come from U_1 .

•
$$P(T/R) = \frac{P(T \cap R)}{P(R)} = \frac{\frac{1}{12}}{\frac{13}{48}} = \frac{4}{13}$$

- 3) Calculate the probability of getting only one red ball.
- $P(N) = P(N \cap T) + P(N \cap H) = P(N/T) \times P(T) + P(N/H) \times P(H)$

$$= \frac{C_6^1 \times C_4^1}{C_{10}^2} \times \frac{1}{4} + \left(\frac{5}{10} \times \frac{5}{10} \times 2!\right) \times \frac{3}{4} = \frac{61}{120}$$

