

GENOMEPATH CLINICAL DIAGNOSTICS (SYNTHETIC FACILITY)

3600 Synthetic Boulevard, Suite 500, Philadelphia, PA 19100 | CLIA ID: 00D7654321 | Director: Dr. Sarah Jenkins, MD, FCAP (Synthetic)

Hereditary Cancer Risk 15-Gene NGS Panel

Germline Sequence and Deletion/Duplication Analysis (Synthetic Demonstration)

Patient Demographics	Ordering Geneticist	Specimen Details
Name: Brenda S. Sample DOB: 09/30/1985 Sex: Female Patient ID: SYN-MRN-9912048	Provider: Dr. Ethan Ross Facility: Pennsylvania Genetics Institute NPI: 9847392015 Location: Philadelphia, PA	Specimen ID: SYN-SPEC-NGS-2098 Collection Date: Jul 15, 2026 Received Date: Jul 16, 2026 Report Date: Jul 24, 2026 Specimen Type: Whole Blood

Result: Pathogenic Variant Identified

Status: Final
Gene: BRCA1 (Heterozygous)

Test Result Summary: A heterozygous pathogenic frameshift variant was identified in the BRCA1 gene in this synthetic sample. No other pathogenic alterations or copy number variants were detected in the remaining 14 genes tested.

Detailed Genetic Variant Findings

Gene Tested	LOINC	Classification & HGVS Variant	Zygosity	Interpretation
BRCA1 Genetic Analysis	69548-6	Pathogenic - c.5266dupC (p.Gln1756Profs*74)	Heterozygous	Abnormal (Pathogenic)
BRCA2 Genetic Analysis	69548-6	Negative / No Pathogenic Variant	N/A	Normal
PALB2 Genetic Analysis	69548-6	Negative / No Pathogenic Variant	N/A	Normal
TP53 Genetic Analysis	69548-6	Negative / No Pathogenic Variant	N/A	Normal
CHEK2 Genetic Analysis	69548-6	Negative / No Pathogenic Variant	N/A	Normal
CDH1 Genetic Analysis	69548-6	Negative / No Pathogenic Variant	N/A	Normal

Clinical Interpretation

The BRCA1 c.5266dupC (p.Gln1756Profs*74) variant is a well-characterized founder mutation in the BRCA1 tumor suppressor gene that results in premature protein truncation. This finding confirms a diagnosis of Hereditary Breast and Ovarian Cancer (HBOC) syndrome. High-risk surveillance protocols (e.g. annual breast MRI, risk-reducing salpingo-oophorectomy discussions) and cascade testing for at-risk first-degree relatives are strongly indicated.

Methodology & Limitations

Methodology: Genomic DNA is extracted from whole blood. Target enrichment of all coding exons and +/- 20bp flanking intronic boundaries of 15 hereditary cancer predisposition genes is performed using custom hybridization probes, followed by paired-end NextGen sequencing (Illumina NovaSeq). Copy number variations (CNVs) are called using normalized read-depth algorithms.

Limitations: This assay does not detect deep intronic mutations outside the targeted regions, complex chromosomal rearrangements, or low-level somatic mosaicism.