

# RESEARCH STATEMENT

TERRENCE GEORGE

My research lies at the intersection of combinatorics, probability, and algebraic geometry. I have explored combinatorial models in statistical mechanics—specifically the dimer model, electrical networks, and the Ising model—and their connections to discrete geometry, integrable systems, algebraic geometry, etc.

The **dimer model** was originally studied in statistical physics as a model for the adsorption of diatomic molecules on crystal surfaces. It can also be interpreted as a model for a melting crystal. Mathematically, a **dimer cover** (or **perfect matching**) of a graph is a subset of the edges that use every vertex exactly once (Figure 1). Enumeration of perfect matchings is a classical problem in graph theory and known to be computationally hard. A celebrated result of Kasteleyn [Kas63] and of Temperley and Fischer [TF61] says that, for planar bipartite graphs, the number of dimer covers of a can be computed in polynomial time as the determinant of a version of the adjacency matrix of the graph, called the **Kasteleyn matrix**. Underpinning the success of the dimer model is this determinantal structure which effectively reduces probabilistic computations for the dimer model to linear algebra.

Beyond statistical mechanics, recent research has uncovered surprising and profound connections between the determinantal structure of the dimer model and various fields, including total positivity in combinatorics, integrable aspects of discrete differential geometry, discrete complex analysis and graph embeddings, and Harnack curves and Beauville integrable systems in algebraic geometry. *Through my work, I aim to contribute to a deeper understanding of these relationships and the broader implications of the dimer model across these various fields, with a particular emphasis on integrating geometric and algebraic methods.*

## 1. LIMIT SHAPES

A fascinating aspect of dimer models is the emergence of **limit shapes**, a manifestation of the law of large numbers, where a random dimer cover of a large graph concentrates around a deterministic shape. The first result of this kind was the celebrated Arctic Circle Theorem for domino tilings [JPS98]. These limit shapes can be interpreted as modeling melting




FIGURE 1. A dimer cover of a portion of the square lattice graph which models the surface of a crystal with two types of molecules corresponding to black and white vertices.




FIGURE 2. Simulations of random groves and percolation configurations on the triangular lattice showing the emergence of limit shapes. In each picture, there is a “liquid” region in the middle outside which the random object appears frozen or “solid”.

crystals in equilibrium and have distinct macroscopic regions that correspond to solid and liquid phases. A common theme is that results about the dimer model have parallels in other statistical mechanical models, such as electrical networks and Ising models. In the context of electrical networks, the equivalent of dimer covers are spanning-tree-like objects called **groves**. In [Geo21], I used the integrable structure of the model to extend the Arctic Circle Theorem for groves due to Petersen and Speyer [PS05] (Figure 2(a)). While this technique only identifies the boundary between different phases and not the full limit shape, it shows that groves display similarly rich limit shape behavior as dimers and deserve further study using more powerful techniques such as variational methods. With Russkikh, Kenyon and Vu, I am investigating limit shapes in Fortuin–Kasteleyn (FK) percolation models, a one-parameter family of statistical-mechanical models generalizing groves (Figure 2(b)).

## 2. TOTAL POSITIVITY

Many of the connections of the dimer model to other areas of mathematics involve moduli spaces of weighted bipartite graphs on surfaces, which possess a rich underlying structure. These spaces are instances of Fock and Goncharov’s **cluster varieties** [FG09], the geometric analogs of Fomin and Zelevinsky’s cluster algebras [FZ02].

The **Grassmannian**, the space parameterizing all subspaces of a fixed dimension of a vector space, is a classical object in algebraic geometry. An explicit way to present an element of the Grassmannian is to specify a rectangular matrix whose rows span the subspace. The maximal minors of the matrix provides a set of homogeneous coordinates called **Plücker coordinates**. A totally nonnegative matrix is a matrix whose minors are all nonnegative. Generalizing this notion, Postnikov [Pos06] defined the **totally nonnegative Grassmannian** to be the set of elements of the Grassmannian with all Plücker coordinates nonnegative. Roughly, the Kasteleyn matrix of a dimer model on a disk is a matrix representative of an element of the totally nonnegative Grassmannian. Postnikov showed, in an equivalent formulation in terms of nonintersecting paths, that this construction, called **boundary**




FIGURE 3. Degeneration of a weighted bipartite graph with weights as  $t \rightarrow 0$ , and the corresponding degeneration of the (amoeba of the) Harnack curve and line bundle (drawn as a divisor).




FIGURE 4. An example of a stratified “totally nonnegative” space parameterizing degenerations of weighted bipartite graphs on a torus/Harnack curves and line bundles. Each cell is labeled by an equivalence class of bipartite graphs, a representative of which is shown.

**measurement**, is a bijection between the space of weighted bipartite graphs and the totally nonnegative Grassmannian. Moreover, degenerations of weighted bipartite graphs induce a beautiful stratification of the totally nonnegative Grassmannian into **positroid** cells. Lam [Lam18] showed that the space of electrical networks gives rise to a stratified subspace of the totally nonnegative Grassmannian which, with Chepuri and Speyer [CGS21], we showed is a totally nonnegative Lagrangian Grassmannian. Together with work of Galashin and Pylyavskyy on the Ising model [GP20], this completes a compelling classification of totally nonnegative Grassmannians associated to statistical-mechanical models in the disk.

In [Geo24c], I studied the problem of reconstructing electrical networks from boundary measurements, a classical problem for which I provided a new solution using ideas from cluster algebras.

### 3. TOTAL POSITIVITY ON A TORUS

Aside from the disk, the best understood dimer models are those on a torus. Classically, they were studied because in statistical mechanics, one is interested in taking limits of planar graphs as the graphs get larger. A way to take the limit that avoids boundary issues is to take larger and larger covers of a graph on a torus. In this setting, it is natural to consider action of the Kasteleyn matrix on translation eigenspaces. This modified Kasteleyn matrix contains two variables  $z, w$  (translation eigenvalues) so its determinant is a polynomial in  $z, w$  and therefore defines an algebraic curve called the **spectral curve**. Kenyon, Okounkov and Sheffield [KOS06] proved a surprising and deep connection between the dimer model and real algebraic geometry. They showed that the spectral curve is a real algebraic curve with a special topology called a **Harnack curve**, and used this fact to obtain a universal classification of the phases of the dimer model. Later, Kenyon and Okounkov [KO06] showed that there is a bijection, called **spectral transform**, between the space of weighted bipartite graphs on the torus and the moduli space of Harnack curves along with certain line bundles on them, or equivalently, points in the Jacobian variety. **The spectral transform is the torus-analog of Postnikov's boundary measurement on the disk and, jointly with Pavel Galashin, one of my long term research goals is to understand degenerations of weighted bipartite graphs on the torus (Figure 3) and the induced positroid-like stratification on the moduli space of Harnack curves with line bundles. As a first step, in [GG24], we study the combinatorics of the stratification and determine the analog of positroids, which is already considerably more complicated and subtle than in Postnikov's setting (Figure 4).**

### 4. THE INVERSE SPECTRAL TRANSFORM

Cluster varieties are equipped with a canonical Poisson structure [GSV10] compatible with local transformations called cluster mutations. The dimer model has a local transformation called the **spider move** which was known from the beginnings of cluster algebras to be an example of a cluster mutation. Upgrading these observations, Goncharov and Kenyon [GK13] constructed, for each convex polygon in the plane with integer vertices, a Hamiltonian integrable system called a **cluster integrable system** from the dimer model on a torus. It was later shown in [FM16] that many well-known integrable systems such as the Toda lattice and the pentagram map are special cases of cluster integrable systems. The **spectral transform** of Kenyon and Okounkov provides action-angle coordinates for the integrable system, a set of coordinates in which the time evolution of the system is linear. Fock [Foc15], using the theory of finite-gap solutions of integrable systems, explicitly constructed the inverse of the spectral transform and showed that the spider move is equivalent to Fay's celebrated trisecant identity for theta functions on Riemann surfaces. **With Goncharov and Kenyon [GGK23], I gave an alternative construction of the inverse spectral transform that only involves rational functions (as opposed to the transcendental theta functions) and is hence amenable to symbolic computation.**

In [Geo24a], I studied the generalization of cluster integrable systems to electrical networks. I showed that the spectral curve is a Harnack curve with an involution. Whenever a curve carries an involution, it determines a linear subvariety of the Jacobian variety called the Prym variety. I explicitly constructed the inverse of the spectral transform using Prym theta functions and showed that the local move for electrical networks, the star-triangle transformation, is equivalent to Fay's quadriseccant identity for Prym theta functions. In [Geo24b], I investigated the Ising model on the torus and showed, through non-explicit

methods, that the spectral transform forms a bijection with a space of symmetric Harnack curves and Prym varieties. These papers reveal that these models possess many, if not all, of the integrable structures present in the dimer model, and I believe there is still much to be explored in this area. An interesting open problem is to construct an explicit inverse map using theta functions and to find the theta function identity describing the Ising star-triangle transformation. Another significant problem is to find Poisson structures compatible with electrical and Ising star-triangle transformations and to prove Arnold–Liouville integrability.

Associated with a convex polygon having integer coordinates in the plane is another integrable system from algebraic geometry: the **Beauville integrable system** [Bea91, Bea90] on the corresponding projective toric surface. Goncharov and Kenyon [GK13] conjectured that the spectral transform is an isomorphism between integrable systems. After initial strategy of Goncharov and Kenyon to prove this conjecture through combinatorial methods were unsuccessful, Inchiostro and I proved it in [GI22] using homological algebra machinery for the dimer model on the hexagonal lattice, i.e., in the case where the toric surface is the projective plane. Establishing the equivalence of cluster integrable systems with Beauville integrable systems in general would be very interesting. I also believe that these powerful, though abstract, algebraic techniques could have broader applications than what we have explored so far.

## 5. DISCRETE INTEGRABLE SYSTEMS

We obtain discrete dynamical systems by applying sequences of spider moves on weighted bipartite graphs, known as **cluster modular transformations** as defined in [FG09]. An example is the pentagram map [OST10], which is a cluster modular transformation on a specific square lattice. In addition to integrable systems in geometry, cluster modular transformations also emerge from the combinatorial and probabilistic study of the dimer model. These transformations have been used to compute partition functions, perform exact sampling via shuffling algorithms, and prove limit shape results (see, e.g., [EKL92, JPS98, PS05, KP16]). Fock and Marshakov [FM16] investigated the group of cluster modular transformations and conjectured its isomorphism type, a conjecture that Inchiostro and I confirmed in [GI24].

Recently, several integrable systems have been discovered in discrete differential geometry (see, e.g., [ILP16, AFIT22, Izo23]) that belong to the class of cluster integrable systems. However, their discrete dynamics are governed not only by sequences of spider moves but also by an additional non-local transformation known as the geometric R-matrix, first introduced by Inoue, Lam, and Pylyavskyy [ILP16]. The group of discrete dynamics that arises in this context is termed the generalized cluster modular group, which Ramassamy and I computed in [GR23], extending the results from [GI24].

Together with Affolter and Ramassamy, I developed a general framework in [AGR21] to prove the equivalence of these integrable systems with cluster integrable systems. Unlike previous approaches that relied on algebraic methods to compare coordinates, our work introduces a novel geometric perspective to make this identification. This approach, along with [AGR] and [AGPR24], provides a unified understanding of the integrable aspects of discrete differential geometry, dimer models, and cluster algebras.

## 6. OTHER SHORT AND MEDIUM TERM PROJECTS

Below, I list some future projects that align with my research interests and knowledge. They could be suitable for either a PhD or strong Masters thesis.

**6.1. The twist map for electrical networks/Ising models.** Cluster varieties are part of an ensemble  $\mathcal{A} \rightarrow \mathcal{X}$ , comprising two varieties,  $\mathcal{A}$  and  $\mathcal{X}$ , with a canonical map between them. In the case of the dimer model on a disk, both  $\mathcal{A}$  and  $\mathcal{X}$  are subvarieties of the Grassmannian, called open positroid varieties [KLS13], and the canonical map is an automorphism between these varieties, called the *twist* [MS16, MS17]. Although electrical networks and Ising models are part of similar cluster-like ensembles [GK13, KP16], where the  $\mathcal{X}$  varieties are relatively well understood [BGKT23, GP20, Lam18, CGS21], the  $\mathcal{A}$  varieties and the twist require further investigation. I worked out the simplest case in [Geo24c].

**6.2. Positive geometry.** An exciting development at the intersection of combinatorics and algebraic geometry, inspired by the study of scattering amplitudes in physics, is the emerging field of Positive Geometry [AHBL17]. Lam [Lam16] showed that positroid varieties are instances of positive geometries. It would be interesting to prove that electrical networks and Ising models are also examples of positive geometries. Doing this would likely require new approaches to investigate the geometric properties of these spaces, as the representation theoretic methods used for positroid varieties appear to be inadequate.

**6.3. Electrical matrix Schubert varieties.** Matrix Schubert varieties, introduced by Fulton [Ful92], are determinantal varieties whose Gröbner degenerations were shown by Knutson and Miller [KM05] to provide a geometric explanation for the positive expansion of Schubert polynomials using pipe dreams. Snyder [Sni10] extended this to open patches in positroid varieties and affine pipe dreams. Together with Speyer, we computed several examples of analogous degenerations of response matrices in electrical networks, characterized by objects resembling pipe dreams, which would be very interesting to investigate further.

**6.4. Limit shapes for cube groves using the spectral transform.** Recently, Boutillier and de Tilière [BdT24] applied Fock’s inverse spectral transform [Foc15] for the dimer model to investigate domino tilings of the Aztec diamond, extending the work of Borodin and Berggren [BB23]. It would be interesting to use the spectral transform for electrical networks, which I studied in [Geo24a], to establish stronger limit shape results for groves.

## REFERENCES

- [AFIT22] Maxim Arnold, Dmitry Fuchs, Ivan Izmistiev, and Serge Tabachnikov. Cross-ratio dynamics on ideal polygons. *Int. Math. Res. Not. IMRN*, (9):6770–6853, 2022.
- [AGPR24] Niklas Affolter, Max Glick, Pavlo Pylyavskyy, and Sanjay Ramassamy. Vector-relation configurations and plabic graphs. *Selecta Math. (N.S.)*, 30(1):Paper No. 9, 55, 2024.
- [AGR] Niklas Affolter, Max Glick, and Sanjay Ramassamy. To appear.
- [AGR21] Niklas Affolter, Terrence George, and Sanjay Ramassamy. Integrable dynamics in projective geometry via dimers and triple crossing diagram maps on the cylinder. 2021.
- [AHBL17] Nima Arkani-Hamed, Yuntao Bai, and Thomas Lam. Positive geometries and canonical forms. *J. High Energy Phys.*, (11):039, front matter+121, 2017.
- [BB23] Tomas Berggren and Alexei Borodin. Geometry of the doubly periodic aztec dimer model, 2023.
- [BdT24] Cédric Boutillier and Béatrice de Tilière. Fock’s dimer model on the aztec diamond, 2024.
- [Bea90] Arnaud Beauville. Jacobiennes des courbes spectrales et systèmes hamiltoniens complètement intégrables. *Acta Math.*, 164(3-4):211–235, 1990.
- [Bea91] A. Beauville. Systèmes hamiltoniens complètement intégrables associés aux surfaces  $K3$ . In *Problems in the theory of surfaces and their classification (Cortona, 1988)*, Sympos. Math., XXXII, pages 25–31. Academic Press, London, 1991.
- [BGKT23] B. Bychkov, V. Gorbounov, A. Kazakov, and D. Talalaev. Electrical networks, Lagrangian Grassmannians, and symplectic groups. *Mosc. Math. J.*, 23(2):133–167, 2023.

- [CGS21] Sunita Chepuri, Terrence George, and David E Speyer. Electrical networks and lagrangian grassmannians. 2021.
- [EKLP92] Noam Elkies, Greg Kuperberg, Michael Larsen, and James Propp. Alternating-sign matrices and domino tilings. I. *J. Algebraic Combin.*, 1(2):111–132, 1992.
- [FG09] Vladimir V. Fock and Alexander B. Goncharov. Cluster ensembles, quantization and the dilogarithm. *Ann. Sci. Éc. Norm. Supér. (4)*, 42(6):865–930, 2009.
- [FM16] Vladimir V. Fock and Andrey Marshakov. Loop groups, clusters, dimers and integrable systems. In *Geometry and quantization of moduli spaces*, Adv. Courses Math. CRM Barcelona, pages 1–66. Birkhäuser/Springer, Cham, 2016.
- [Foc15] V. V. Fock. Inverse spectral problem for gk integrable system. 2015.
- [Ful92] William Fulton. Flags, Schubert polynomials, degeneracy loci, and determinantal formulas. *Duke Math. J.*, 65(3):381–420, 1992.
- [FZ02] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. I. Foundations. *J. Amer. Math. Soc.*, 15(2):497–529, 2002.
- [Geo21] Terrence George. Grove arctic curves from periodic cluster modular transformations. *Int. Math. Res. Not. IMRN*, (20):15301–15336, 2021.
- [Geo24a] Terrence George. Spectra of biperiodic planar networks. *Comm. Math. Phys.*, 405(1):Paper No. 10, 47, 2024.
- [Geo24b] Terrence George. Spectral transform for the ising model. 2024.
- [Geo24c] Terrence George. The twist for electrical networks and the inverse problem. *Int. Math. Res. Not. IMRN*, (8):7001–7031, 2024.
- [GG24] Pavel Galashin and Terrence George. Move-reduced graphs on a torus. *Trans. Amer. Math. Soc.*, 377(6):4055–4099, 2024.
- [GGK23] T. George, A. B. Goncharov, and R. Kenyon. The inverse spectral map for dimers. *Math. Phys. Anal. Geom.*, 26(3):Paper No. 24, 51, 2023.
- [GI22] Terrence George and Giovanni Inchiostro. Dimers and beauville integrable systems. 2022.
- [GI24] Terrence George and Giovanni Inchiostro. The cluster modular group of the dimer model. *Ann. Inst. Henri Poincaré D*, 11(1):147–198, 2024.
- [GK13] Alexander B. Goncharov and Richard Kenyon. Dimers and cluster integrable systems. *Ann. Sci. Éc. Norm. Supér. (4)*, 46(5):747–813, 2013.
- [GP20] Pavel Galashin and Pavlo Pylyavskyy. Ising model and the positive orthogonal Grassmannian. *Duke Math. J.*, 169(10):1877–1942, 2020.
- [GR23] Terrence George and Sanjay Ramassamy. Discrete dynamics in cluster integrable systems from geometric  $R$ -matrix transformations. *Comb. Theory*, 3(2):Paper No. 12, 29, 2023.
- [GSV10] Michael Gekhtman, Michael Shapiro, and Alek Vainshtein. *Cluster algebras and Poisson geometry*, volume 167 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.
- [ILP16] Rei Inoue, Thomas Lam, and Pavlo Pylyavskyy. Toric networks, geometric  $R$ -matrices and generalized discrete Toda lattices. *Comm. Math. Phys.*, 347(3):799–855, 2016.
- [Izo23] Anton Izosimov. Polygon recutting as a cluster integrable system. *Selecta Math. (N.S.)*, 29(2):Paper No. 21, 31, 2023.
- [JPS98] William Jockusch, James Propp, and Peter Shor. Random domino tilings and the arctic circle theorem. 1998.
- [Kas63] P. W. Kasteleyn. Dimer statistics and phase transitions. *J. Mathematical Phys.*, 4:287–293, 1963.
- [KLS13] Allen Knutson, Thomas Lam, and David E. Speyer. Positroid varieties: juggling and geometry. *Compos. Math.*, 149(10):1710–1752, 2013.
- [KM05] Allen Knutson and Ezra Miller. Gröbner geometry of Schubert polynomials. *Ann. of Math. (2)*, 161(3):1245–1318, 2005.
- [KO06] Richard Kenyon and Andrei Okounkov. Planar dimers and Harnack curves. *Duke Math. J.*, 131(3):499–524, 2006.
- [KOS06] Richard Kenyon, Andrei Okounkov, and Scott Sheffield. Dimers and amoebae. *Ann. of Math. (2)*, 163(3):1019–1056, 2006.
- [KP16] Richard Kenyon and Robin Pemantle. Double-dimers, the Ising model and the hexahedron recurrence. *J. Combin. Theory Ser. A*, 137:27–63, 2016.

- [Lam16] Thomas Lam. Totally nonnegative Grassmannian and Grassmann polytopes. In *Current developments in mathematics 2014*, pages 51–152. Int. Press, Somerville, MA, 2016.
- [Lam18] Thomas Lam. Electroid varieties and a compactification of the space of electrical networks. *Adv. Math.*, 338:549–600, 2018.
- [MS16] R. J. Marsh and J. S. Scott. Twists of Plücker coordinates as dimer partition functions. *Comm. Math. Phys.*, 341(3):821–884, 2016.
- [MS17] Greg Muller and David E. Speyer. The twist for positroid varieties. *Proc. Lond. Math. Soc. (3)*, 115(5):1014–1071, 2017.
- [OST10] Valentin Ovsienko, Richard Schwartz, and Serge Tabachnikov. The pentagram map: a discrete integrable system. *Comm. Math. Phys.*, 299(2):409–446, 2010.
- [Pos06] Alexander Postnikov. Total positivity, grassmannians, and networks. 2006.
- [PS05] T. Kyle Petersen and David Speyer. An arctic circle theorem for Groves. *J. Combin. Theory Ser. A*, 111(1):137–164, 2005.
- [Sni10] Michelle Snider. Affine patches on positroid varieties and affine pipe dreams (thesis), 2010.
- [TF61] H. N. V. Temperley and Michael E. Fisher. Dimer problem in statistical mechanics—an exact result. *Philos. Mag. (8)*, 6:1061–1063, 1961.