

Karim Andre Drouin
825 Brickell Bay Drive
Miami, FL 33130

Test Report

Patient	Date of Birth	Report Date/Time	Collection Date/Time	Medical Record Number
Karim Andre Drouin	3/2/1992	4/16/2022 10:36AM	4/16/2022 10:02AM	20359-00423
Test	Employee ID	Result		Type of Test
SARS-CoV-2	N/A	SARS-CoV-2 Not Detected		COVID-Ag

Name and Address of Lab Where Test Was Performed

Salvation Army: 911 West Flagler St, Miami, FL 33130, USA
If your result is **Not Detected** for SARS-CoV-2 (Coronavirus 2), the virus that causes COVID-19 this may mean you were not infected at the time your test was performed or there was not enough virus present to be detected. It does not mean you will not get infected or sick. It is possible that you were very early in your infection at the time of your test and that you could test positive later, or you could be exposed later and then develop the illness.

If you had close contact with someone with COVID-19 you must continue to quarantine. You should quarantine for 14 days from your last contact with an infected individual. Please see the CDC website for details on what constitutes close contact and quarantine information. <https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/quarantine.html>

Monitor your health daily for symptoms of COVID. Possible symptoms include fever, chills, cough, shortness of breath, fatigue, body aches, headache, new loss of taste or smell, sore throat, nasal congestion, nausea, vomiting or diarrhea.

If your result is **Detected** for SARS-CoV-2 (Coronavirus 2), the virus that causes COVID-19, please follow these recommendations:

Stay at home (self-isolation): Most people with COVID-19 have mild illness and recover at home without medical care.

- Do not leave your home, except to get medical care.
- Do not go to work or school, use public transportation, or visit public places.
- Avoid contact with other household members, if possible.
- Use a separate bedroom and bathroom than other household contacts, if possible.
- Wear a mask when around other people.

When to end isolation:

- 10 days since symptoms first appeared and
- 24 hours with no fever without the use of fever-reducing medications and
- Other symptoms of COVID-19 are improving. *Loss of taste and smell may persist for weeks or months after recovery and need not delay the end of isolation.*
- If you had severe illness or are immunocompromised consult your doctor on when it is safe to end isolation.

Monitor your symptoms:

Look for **emergency warning signs** for COVID-19. If someone is showing any of these signs, **seek emergency medical care immediately**:

- Trouble breathing
- Persistent pain or pressure in the chest
- New confusion
- Inability to wake or stay awake
- Bluish lips or face

The CDC self-checker is a tool to help you make decisions and seek appropriate care. It can be accessed at <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/coronavirus-self-checker.html>. Contact your doctor if you have questions about your illness or on whether you need to seek medical care. Call ahead and tell them you have COVID-19.

Notify your close contacts: Alert people that you have been in close contact with while ill that you have tested positive. Tell them to quarantine for 14 days and monitor their health for signs of COVID-19.

If your result is **INVALID** for SARS-CoV-2 (Coronavirus 2), the virus that causes COVID-19, please refer to the reason listed above. We recommend that you reschedule for additional testing.

If your result is **REJECTED** for SARS-CoV-2 (Coronavirus 2), the virus that causes COVID-19, please refer to the reason listed above. We recommend that you reschedule for additional testing.

Our knowledge of the Coronavirus 2 and the disease it causes is rapidly evolving. For the most up-to-date information, please visit the Centers for Disease Control and Prevention website <https://www.cdc.gov/coronavirus/2019-nCoV/index.html>

Method: The CareStart™ COVID-19 Test is a type of test called an antigen test. Antigen tests are designed to detect proteins from the virus that causes COVID-19 in respiratory specimens, for example nasopharyngeal swabs.

Limitations: There are different kinds of tests for COVID-19. Molecular tests (also known as PCR tests) detect genetic material from the virus. Antigen tests detect proteins from the virus. Antigen tests are specific for the virus but are not as sensitive as molecular tests. This means that a positive result is highly accurate, but a negative result does not rule out infection.

Disclaimer: The CareStart™ COVID-19 Ag Test EUA has not been FDA cleared or approved. It has been authorized by the FDA under an emergency use authorization for use by authorized entities. The test has been authorized only for the detection of proteins from SARS-CoV-2, not for any other viruses or pathogens, and is only authorized for the duration of the declaration that circumstances exist justifying the authorization of emergency use of in vitro diagnostic tests for detection and/or diagnosis of COVID-19 under Section 564(b)(1) of the Act, 21 U.S.C. § 360bbb-3(b)(1), unless the authorization is terminated or revoked sooner.

Fact Sheets

<https://www.fda.gov/media/142918/download>
<https://www.fda.gov/media/142917/download>