

QUANTITATIVE RESEARCH METHODS

DR. MEIKE MORREN

Lecture 5

contents



- Multiple regression
- Mean centering
- Standardization

MULTIPLE REGRESSION



Multiple vs simple regression

- Estimation becomes more complicated when multiple explanatory variables are included
- A general method would be the least squares (ordinary least squares – OLS) where one obtains the regression coefficients by minimizing the errors
- In order to compute the coefficients, we need to use derivations

OLS (1)

- Minimizing the sum of the squared deviations of the Y_i 's
- This minimized solution provides reliable and stable estimates of β_n
- The estimated regression function is written

$$\hat{Y}_i = b_0 + b_1x_1 + \cdots + b_nx_n + \varepsilon_i$$

- Another way to model this the relationship is

$$f_{\theta}(x) = \theta_1x_1 + \cdots + \theta_nx_n$$

OLS(2)

- We want to minimize the least-squares cost function

$$J(\theta) = \frac{1}{2m} \sum_{i=1}^m (f_{\theta}(x^{(i)}) - y^{(i)})^2$$

Where $x^{(i)}$ is the i th observation and
 $y^{(i)}$ is the i th expected result

OLS (3)

- We can rewrite this loss function J as

$$f_{\theta}(x) = \theta^T x$$

- With this we can rewrite the least-squares cost function using matrix multiplication

$$J(\theta_{0..n}) = \frac{1}{2m} (X\theta - y)^T (X\theta - y)$$

Derivatives

$$\partial J / \partial \theta = 2X^T X \theta - 2X^T y = 0$$

$$X^T X \theta = X^T y$$

If the matrix $X^T X$ is invertible. we can multiply both sides by $(X^T X)^{-1}$ and get

$$\theta = (X^T X)^{-1} X^T y$$

OLS estimation: step by step

- This formula can be used to estimate multiple regression coefficients
- 1. Combine all k independent variables in columns in a matrix ($n \times k$)
- 2. Add a vector of 1s to estimate the intercept
- 3. Make a vector of the dependent variable
- 4. Solve the formula

Example mtcars

- Include both horsepower and weight
- Estimate linear regression using the formula
- Plot two variables
- Add regression line using the coefficients that you found

Y and X1 and X2 (hp and wt)

```
> x
```

```
      [,1] [,2] [,3]  
[1,]      1  110 2.620  
[2,]      1  110 2.875  
[3,]      1   93 2.320  
[4,]      1  110 3.215  
[5,]      1  175 3.440  
[6,]      1  105 3.460  
[7,]      1  245 3.570  
[8,]      1   62 3.190  
[9,]      1   95 3.150  
[10,]     1  123 3.440  
[11,]     1  123 3.440  
[12,]     1  180 4.070  
[13,]     1  180 3.730  
[14,]     1  180 3.780  
[15,]     1  205 5.250  
[16,]     1  215 5.424  
[17,]     1  230 5.345  
[18,]     1   66 2.200  
[19,]     1   52 1.615  
[20,]     1   65 1.835  
[21,]     1   97 2.465  
[22,]     1  150 3.520  
[23,]     1  150 3.435  
[24,]     1  245 3.840  
[25,]     1  175 3.845  
[26,]     1   66 1.935  
[27,]     1   91 2.140  
[28,]     1  113 1.513  
[29,]     1  264 3.170  
[30,]     1  175 2.770  
[31,]     1  335 3.570  
[32,]     1  109 2.780
```

$$\hat{Y}_i = b_0 + b_1x_1 + b_2x_2 + \varepsilon_i$$

```
> y
```

```
[1] 21.0 21.0 22.8 21.4 18.7 18.1 14.3 24.4 22.8 19.2 17.8 16.4 17.3 15.2 10.4 10.4 14.7 32.4 30.4 33.9 21.5 15.5 15.  
[24] 13.3 19.2 27.3 26.0 30.4 15.8 19.7 15.0 21.4
```

```
>
```

Scatterplot of y and X_1

> x


	[,1]	[,2]	[,3]
[1,]	1	110	2.620
[2,]	1	110	2.875
[3,]	1	93	2.320
[4,]	1	110	3.215
[5,]	1	175	3.440
[6,]	1	105	3.460
[7,]	1	245	3.570
[8,]	1	62	3.190
[9,]	1	95	3.150
[10,]	1	123	3.440
[11,]	1	123	3.440
[12,]	1	180	4.070
[13,]	1	180	3.730
[14,]	1	180	3.780
[15,]	1	205	5.250
[16,]	1	215	5.424
[17,]	1	230	5.345
[18,]	1	66	2.200
[19,]	1	52	1.615
[20,]	1	65	1.835
[21,]	1	97	2.465
[22,]	1	150	3.520
[23,]	1	150	3.435
[24,]	1	245	3.840
[25,]	1	175	3.845
[26,]	1	66	1.935
[27,]	1	91	2.140
[28,]	1	113	1.513
[29,]	1	264	3.170
[30,]	1	175	2.770
[31,]	1	335	3.570
[32,]	1	109	2.780

> y

[1]	21.0	21.0	22.8	21.4	18.7	18.1	14.3	24.4	22.8	19.2	17.8	16.4	17.3	15.2	10.4	10.4	14.7	32.4	30.4	33.9	21.5	15.5	15.
[24]	13.3	19.2	27.3	26.0	30.4	15.8	19.7	15.0	21.4														

>

OLS: miles/gallon and horsepower



Scatterplot of y and X_2

> x


	[,1]	[,2]	[,3]
[1,]	1	110	2.620
[2,]	1	110	2.875
[3,]	1	93	2.320
[4,]	1	110	3.215
[5,]	1	175	3.440
[6,]	1	105	3.460
[7,]	1	245	3.570
[8,]	1	62	3.190
[9,]	1	95	3.150
[10,]	1	123	3.440
[11,]	1	123	3.440
[12,]	1	180	4.070
[13,]	1	180	3.730
[14,]	1	180	3.780
[15,]	1	205	5.250
[16,]	1	215	5.424
[17,]	1	230	5.345
[18,]	1	66	2.200
[19,]	1	52	1.615
[20,]	1	65	1.835
[21,]	1	97	2.465
[22,]	1	150	3.520
[23,]	1	150	3.435
[24,]	1	245	3.840
[25,]	1	175	3.845
[26,]	1	66	1.935
[27,]	1	91	2.140
[28,]	1	113	1.513
[29,]	1	264	3.170
[30,]	1	175	2.770
[31,]	1	335	3.570
[32,]	1	109	2.780

> y

[1]	21.0	21.0	22.8	21.4	18.7	18.1	14.3	24.4	22.8	19.2	17.8	16.4	17.3	15.2	10.4	10.4	14.7	32.4	30.4	33.9	21.5	15.5	15.
[24]	13.3	19.2	27.3	26.0	30.4	15.8	19.7	15.0	21.4														


>

OLS: miles/gallon and weight



Equation (1)

$$Y_i = b_0 + b_1x_1 + b_2x_2 + \varepsilon_i$$




An increase of 1 pt in
 X_1 leads to increase
of b_1 in Y

Interpretation beta coefficients:

- A 1-point increase in weight (measured in 1000lbs) leads to a 3.88 decrease in miles per gallon
- Thus, the heavier the car, the fewer miles you can drive with a gallon of gasoline
- **Controlling for horsepower:**
- This effect **holds** for all values of horsepower. So irrespective of how fast the car can drive, a 1 pt increase in weight will always lead to a 3.88 decrease in miles per gallon

Equation (2)

$$Y_i = b_0 + b_1x_1 + b_2x_2 + \varepsilon_i$$



= Constant
= Intercept

Interpretation constant :

The mean level of the dependent variable where ALL the independent variables are 0

Thus...

The mean level of miles per gallon for 0 weight and 0 horsepower

Exe 5_1.r

- Estimate a multiple regression using two interval predictors (or ordinal treated as interval)
- Compare your results with the model where you do not control for the second variable
- Interpret the intercept and coefficients

MULTIPLE REGRESSION

Mean centering

Mean centering

- Usually a zero has no meaning for the independent variables and does not even occur in your data
- This makes the interpretation of the intercept difficult

Mean centering

- Mean centering allows you to interpret the intercept
- By subtracting the mean from all values on the independent variable, you make the mean zero
- The coefficients remain the same: a one point increase is still a one point increase

Mean centering

X	mean		X mean centered
110	110	$110 - 110$	0
110	110	$110 - 110$	0
931	110	$931 - 110$	821
110	110	$110 - 110$	0
175	110	$175 - 110$	65
105	110	$105 - 110$	-5
245	110	$245 - 110$	135
62	110	$62 - 110$	-48
95	110	$95 - 110$	-15

Example mtcars

```
> summary(lm(mpg ~ mhp + mwt, data = mtcars))
```

Call:

```
lm(formula = mpg ~ mhp + mwt, data = mtcars)
```

Residuals:

Min	1Q	Median	3Q	Max
-3.941	-1.600	-0.182	1.050	5.854

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	20.09062	0.45846	43.822	< 2e-16 ***
mhp	-0.03177	0.00903	-3.519	0.00145 **
mwt	-3.87783	0.63273	-6.129	1.12e-06 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.593 on 29 degrees of freedom
Multiple R-squared: 0.8268, Adjusted R-squared: 0.8148
F-statistic: 69.21 on 2 and 29 DF, p-value: 9.109e-12

```
> summary(lm(mpg ~ hp + wt, data = mtcars))
```

Call:

```
lm(formula = mpg ~ hp + wt, data = mtcars)
```

Residuals:

Min	1Q	Median	3Q	Max
-3.941	-1.600	-0.182	1.050	5.854


Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	37.22727	1.59879	23.285	< 2e-16 ***
hp	-0.03177	0.00903	-3.519	0.00145 **
wt	-3.87783	0.63273	-6.129	1.12e-06 ***


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Example mtcars horsepower

OLS: miles/gallon and horsepower



mean centering



Exe 5_2.r

- Center all your independent variables
- Estimate the model again, and compare with your result obtained in exe 5_1.r

MULTIPLE REGRESSION

Standardized coefficients

Standardized regression

- If you have multiple variables that have a different range of values, the unstandardized coefficients are hard to compare in terms of strength
- A 1-point increase in one variable means something else than a 1-point increase in another variable
- Therefore, ALL variables are standardized
- The 1-point increase becomes a 1-standard deviation increase

Z-scores

(centered and standardized)

- Calculate z-scores:

$$z_i = \frac{x_i - \bar{X}}{\sigma_x}$$


1. Calculate mean
2. Calculate standard deviation (sd)
3. Calculate z scores

Z-scores have a
mean of zero and standard deviation of 1


Example mtcars

- First calculate z-scores
- Create vectors of standardized scores
- Solve equation with standardized values
- Plot the line
- Compare with previous results

OLS: miles/gallon and horsepower




standardized




Example mtcars : weight

OLS: miles/gallon and weight



standardized



Interpretation effect weight:

- A 1 **standard deviation** increase in weight (measured in 1000lbs) leads to a 3.88 **standard deviation** decrease in miles per gallon
- Thus, the heavier the car, the fewer miles you can drive with a gallon of gasoline
- Controlling for horsepower:
- This effect holds for all values of horsepower. So irrespective of how fast the car can drive, an increase in weight will always lead to a decrease in miles per gallon

Exe 5_3.r

- Standardize all your variables (independent and dependent!)
- Estimate the model again, and compare with your result obtained in exe 5_1.r and exe 5_2.r

Next lecture

- moderation
- (if we have the time) mediation
- We will use the other dataset on tablets!