



Pacific Community
Communauté
du Pacifique

A summary of stock status and national implications for the Cook Islands



FAME

Fisheries, Aquaculture and Marine Ecosystems Division

Contents

Executive summary	3
Introduction	4
Summary of tuna stock status	5
Skipjack tuna	5
Yellowfin tuna	8
Bigeye tuna	10
South Pacific albacore tuna	12
Other species	14
References	16
Tables	17
Figures	17

List of Tables

1 Definitions of key terms used in describing the impact of fishing upon and the status of fish stocks	4
2 Key stock and fishery catch statistics for the WCPFC convention area, including the recent period of 2018 – 2022 in the SB EEZ	5

List of Figures

1 test short caption	6
2 test short caption	7
3 test short caption	9
4 test short caption	11
5 test short caption	13
6 test short caption	15

Executive summary

Text

Introduction

Tuna stock assessments provide important information regarding the status of regional tuna stocks and the future predicted impacts of fishing on them. This information can help the Solomon Islands and other WCPFC members decide how best to collectively manage these stocks.

Albacore, bigeye, yellowfin and skipjack tuna are targeted by fisheries operating in the Solomon Islands. The SPC has assessed these stocks in the WCPFC-CA using a modeling tool called MULTIFAN-CL, which is reliant on accurate and comprehensive catch, effort, size and other fisheries data collected by all fishing nations. The assessments provide information on stock status at the WCPFC-CA scale and broad model Regions within that. SPC is working to develop models that can provide similar information at the EEZ scale. However, given that tuna are migratory and may pass through many EEZ's in their lifetime, it will never be feasible to assess and/or manage these species within individual EEZ's in isolation from what is happening in surrounding waters. However, most countries also need to know what the status of the resource is in their immediate vicinity. Therefore we provide here some information for the broad model region that surrounds your EEZ so that you can assess your country's impact on the stock and the state of the stock in your waters.

This report should be used in conjunction with the Tuna Fisheries Assessment Report published annually by the SPC (Hare et al. 2023). Tuna stock assessment results are described using a number of technical terms, and refer to a number of key indicators of stock status (reference points). These terms are described in ??.

Depletion	Depletion describes the level of reduction in the fish stock since fishing first began, typically by comparing current spawning biomass to that which would occur if there was no fishing ($SB_{current}/SB_F = 0$).
Fishing mortality rate	The proportion of the stock removed by fishing in a unit of time. Growth overfished Fish are harvested at an average size that is smaller than the size that would produce the maximum yield per recruit.
Maximum Sustainable Yield (MSY)	The maximum amount of catch that can be taken from the stock per year, on average, in the long-term.
Overfished	Occurs when there are no longer enough adults in the population to produce enough young to replace those fish removed from the population by fishing. In the WCPFC, an overfished fishery is defined as one where the current spawning biomass ($SB_{current}$) is less than 20% of the spawning biomass in the absence of fishing ($SB_F = 0$).
Overfishing	In the WCPFC, overfishing is defined as occurring when the current fishing mortality rate exceeds the fishing mortality rate that would provide the maximum sustainable yield. Sustained overfishing leads to an overfished state.
Recruitment overfished	Occurs when the adult population is depleted to a level where it no longer has the reproductive capacity to replenish itself.

Table 1: Definitions of key terms used in describing the impact of fishing upon and the status of fish stocks

Table 2: Key stock and fishery catch statistics for the WCPFC convention area, including the recent period of 2018 – 2022 in the SB EEZ

Summary description	Albacore	Bigeye	Skipjack	Yellowfin
WCPFC-CA catch	64480	146560	1806281	719736
5-year WCPFC-CA catch trend	Stable	Decreasing	Stable	Increasing
WCPFC-CA Longline catch	60545	60872	3497	87227
WCPFC-CA Purse seine catch	6	65514	1475630	381359
WCPFC-CA Pole and line catch	22	2285	160419	23208
WCPFC-CA Other catch	3906	17889	166735	227942
Catch in SB model regions	15446	80152	1246623	358532
Percent of WCPFC-CA catch in SB model regions	24	54.7	69	49.8
Catch in SB EEZ	4528	3444	73303	29314
Percent of WCPFC-CA catch taken in SB EEZ	7	2.4	4.1	4.1
Percent of Catch in SB model regions taken in SB EEZ	29.3	4.3	5.9	8.2
Catch by SB flagged in WCPFC-CA	2176	1370	30352	19229
Percent of WCPFC-CA catch by SB flagged vessels	3.4	0.9	1.7	2.7

Summary of tuna stock status

The most recent assessments indicate that overfishing is not occurring on albacore, bigeye, skipjack and yellowfin tuna, and these stocks are not in an overfished state (Figure 1). However, catch of each of these species have increased significantly over the past 2 to 3 decades, with corresponding reductions in stock biomass relative to the biomass that would be in the water if no fishing was occurring. The following sections provide more detail regarding the status of these stocks, implications for your national fisheries and briefly reviews assessments of other relevant species. Specific details regarding catch and stock status of each species in the WCPFC and your EEZ are presented in ??.

Skipjack tuna

The most recent skipjack tuna assessment was conducted in 2022 (Castillo-Jordan et al. 2022). The skipjack assessment used an eight region model, which was a change from the five region model previously used in previous assessment; your EEZ is situated in region(s) 6, 7, 8 (Figure 2 - top). Between 2013 and 2022 skipjack catch averaged 1,806,281 mt in the WCPFC-CA (Figure 2 - middle). An average of 1,246,623 mt (69 % of WCPFC-CA catch) comes from region(s) 6, 7, 8. It is estimated that skipjack spawning biomass in the WCPFC-CA and region(s) 6, 7, 8, have been reduced through fishing by **tabl2%** and **tabl2%** respectively (??). The greatest impacts on spawning biomass in the WCPO are equally from the FAD-associated and free school purse seine fisheries, while the FAD-associated purse seine fishery has the greatest impact in region(s) 6, 7, 8 (Figure 2 - bottom).

The SC18 of the WCPFC concluded that overfishing is not occurring on the skipjack stock and the stock is not overfished (Figure 1). The spawning biomass level is below the interim Target Reference Point of 50% ($SB/SB_{F=0} = 0.5$), though well above the Limit Reference Point of 20%, unfished spawning biomass. Previously, the SC12 noted that fishing is having a significant impact on stock size, especially in the western equatorial region and can be expected to negatively affect catch rates. The stock distribution is also influenced by changes in oceanographic conditions associated with El Niño and La Niña events, which impact on catch rates and stock size. Additional purse-seine effort will yield only modest gains in long-term skipjack catch and may result in a corresponding increase in fishing mortality for bigeye and yellowfin tunas (WCPFC 2022).

Annual catch of skipjack in the Solomon Islands has averaged 73,303 mt between 2013 and 2022, representing 4.1 % of the skipjack catch in the WCPFC-CA and 5.9 % of the catch in region(s) 6, 7, 8. Together, skipjack catch by the Solomon Islands fleet (including chartered vessels) both inside and outside the the Solomon Islands EEZ have caught an average of 30,352 mt, which is 1.7 % of the WCPFC skipjack catch. Regional catch of skipjack tuna, including those within or by the Solomon Islands, are




Figure 1: Majuro plot comparing the stock status of the key target tuna species caught in the WCPFC Convention Area. Where current fishing mortality rate exceeds the fishing mortality rate at MSY ($F/F_{MSY} > 1$) then overfishing is occurring. Where the current spawning biomass is less than 0.2 of spawning biomass without fishing, then the stock is overfished ($SB/SB_{F=0} < 0.2$). The large points represent the median estimated stock status in terms of fishing mortality and biomass depletion in the terminal year of the most recent assessment for each species. The error bars indicate stock status uncertainty from all models run in the grid. The blue and green dashed lines indicate the WCPFC interim target reference points for skipjack and albacore tuna, respectively.

considered sustainable at recent average levels. However, the Solomon Islands should note that the FAD component of the regional purse seine fishery (the main fishery for skipjack tuna) catches juvenile bigeye and yellowfin tuna, and that fishery is contributing to the impact on these stocks.




Figure 2: Distribution of total South Pacific albacore catch by fishing method 2013–2022 (Red, pole-and-line; Blue, purse-seine; Green, longline; Orange, troll; Yellow, other) (top). Annual catch of skipjack in the WCPO by fishing method (middle). Percentage impact on spawning biomass due to fishing in the WCPO (bottom - left) and the model Region/s encompassing your EEZ (bottom - right).

Yellowfin tuna

The most recent yellowfin tuna assessment was conducted in 2023 (Magnusson et al. 2023). The yellowfin tuna assessment used a nine region model, with the Solomon Islands situated in region(s) 3, 4, 5 (Figure 3 - top). Between 2013 and 2022 yellowfin catch averaged 719,736 mt in the WCPFC-CA (Figure 3 - middle). An average of 358,532 mt (49.8 % of WCPFC-CA catch) comes from region(s) 3, 4, 5. It is estimated that yellowfin spawning biomass in the WCPFC-CA and region(s) 3, 4, 5, have declined by **tabl2** % and **tabl2** % respectively (??). The greatest impact on the stock is from the fisheries of the Philippines and Indonesia, along with the FAD directed purse-seine fishery in the WCPO and from the FAD directed purse-seine fishery in region(s) 3, 4, 5 (Figure 3 - bottom).

The SC19 (WCPFC 2023) concluded that overfishing is not occurring on the yellowfin stock and the stock is not overfished (Figure 1). However, fishing mortality, exploitation rates and depletion differ between regions, and exploitation rates are highest in the equatorial region (Regions 3, 4, 7 and 8), which account for 94% of the total yellowfin tuna catch. The SC19 recommended that there be no increase in yellowfin catch and that measures be implemented to maintain current spawning biomass levels (WCPFC 2023).

Annual catch of yellowfin in the Solomon Islands have averaged **tabl2t** between **minyear** and **maxyear**, representing **tabl2** % of WCPFC-CA and **tabl2** % of region(s) 3, 4, 5 yellowfin catch. Together, yellowfin catch inside the Solomon Islands and by the Solomon Islands fleet outside your EEZ have accounted for an average **tabl2** % of the WCPFC yellowfin catch. As such, the the Solomon Islands fishery does not contribute significantly to overall regional impacts on the stock. The yellowfin stock in region(s) 3, 4, 5 has low levels of depletion in Region 6 but higher in Region 4 and overall the Solomon Islands catch is a small percentage of the catch in these two regions.





Figure 3: Distribution of total yellowfin catch by fishing method 2013–2022 (Red, pole-and-line; Blue, purse-seine; Green, longline; Orange, troll; Yellow, other) (top). Annual catch of South Pacific albacore in the WCPO by fishing method (middle). Percentage impact on spawning biomass due to fishing in the WCPO (bottom - left) and the model Region/s encompassing your EEZ (bottom - right).

Bigeye tuna

The most recent bigeye tuna assessment was conducted in 2023 (Day et al. 2023). The bigeye assessment used a nine region model, your EEZ is situated in region(s) 3, 4, 5, 8 (Figure 4 - top). Between 2013 and 2022 bigeye catch averaged **tabl2t** in the WCPFC-CA (Figure 4 - middle). An average of **tabl2t** (**tabl2%** of WCPFC-CA catch) comes from 3, 4, 5, 8. It is estimated that bigeye spawning biomass in the WCPFC-CA and 3, 4, 5, 8, have declined by **tabl2%** and **tabl2%** respectively (??). The greatest impact on spawning biomass is from the associated - FAD - purse seine fisheries in the WCPO and from the longline fishery in 3, 4, 5, 8 (Figure 4 - bottom).

The SC14 concluded that overfishing is not occurring on the bigeye stock and the stock is not overfished (Figure 1). However, the increase in juvenile bigeye catch has resulted in a considerable reduction in the potential yield of the WCPO bigeye stock. The loss in yield per recruit due to excess harvest of juvenile fish is substantial.

Annual catch of bigeye in the Solomon Islands have averaged **tabl2t** between **minyear** and **maxyear**, representing **tabl2%** of WCPFC-CA and **tabl2%** of 3, 4, 5, 8 bigeye catch. Together, bigeye catch inside the Solomon Islands and by your fleet outside your EEZ have accounted for an average **tabl2%** of the WCPFC bigeye catch. Regional catch of bigeye tuna, including those within or by the Solomon Islands, are considered to be low if maintained at recent average levels. However, the Solomon Islands should note that the FAD component of the regional purse seine fishery (the main fishery for skipjack tuna) catches juvenile bigeye and yellowfin tuna, and that fishery is contributing to the impact on this stock.

Bigeye tuna is one of two species targeted in the longline fishery operating in the Solomon Islands* and is also taken in the purse seine fishery in the Solomon Islands. The viability of the local and regional purse seine fishery is not dependent on bigeye tuna abundance. However, the prospects for long-term viability (and/or further expansion) of a longline fishery for bigeye in the Solomon Islands will be dependant, in part, on trends in both national and regional fishing mortality on the bigeye stock (and economic and other factors not discussed here).




Figure 4: Distribution of total bigeye catch by fishing method 2013–2022 (Red, pole-and-line; Blue, purse-seine; Green, longline; Orange, troll; Yellow, other) (top). Annual catch of South Pacific albacore in the WCPO by fishing method (middle). Percentage impact on spawning biomass due to fishing in the WCPO (bottom - left) and the model Region/s encompassing your EEZ (bottom - right).

South Pacific albacore tuna

The most recent south Pacific albacore (hereafter, simply “albacore”) assessment was conducted in 2021 (Castillo-Jordan et al. 2021). The albacore assessment used a five region model, your EEZ is situated in 1, 2 (Figure 5 - top). Between 2013 and 2022 albacore catch averaged **tabl2t** in the WCPFC-CA (Figure 5 - middle). An average of **tabl2t** (**tabl2%** of WCPFC-CA catch) comes from 1, 2. It is estimated that albacore spawning biomass in the WCPFC-CA and 1, 2, have declined by **tabl2%** and **tabl2%** respectively (??). The greatest impact on spawning biomass is from the sub-tropical longline fisheries both in the WCPO and in 1, 2 (Figure 5 - bottom).

The WCPFC **SC14** concluded that South Pacific albacore spawning stock is currently above both the level that will support the MSY and the adopted spawning biomass limit reference point, and overfishing is not occurring (Figure 1). But **SC14** also noted that while overfishing is not occurring, further increases in effort will yield little or no increase in long-term catch and result in further reduced catch rates. **SC14** also noted that any increases in catch or effort in sub-tropical longline fisheries are likely to lead to declines in longline catch rates between 10°S-30°S, which will impact vessel profitability **WCPFC2018**. An interim Target Reference Point of $SB/SB_{F=0} = 0.56$ was adopted in 2019 **WCPFC2019**; the current spawning biomass level of 0.52 is estimated to be below the interim TRP

Annual catch of albacore in the Solomon Islands has averaged **tabl2t** between **minyear** and **maxyear**, representing **tabl2%** of WCPFC-CA and **tabl2%** of 1, 2 albacore catch. Together, albacore catch inside the Solomon Islands and by the Solomon Islands fleet outside your EEZ have accounted for an average **tabl2%** of the WCPFC albacore catch. Regional catch of albacore tuna, including those within or by Solomon Islands vessels fishing outside your EEZ, are considered sustainable at recent average levels. But ct.long should note that, deterministic projections¹ **Pilling2018** estimate that there is a 24% chance of the overall stock falling below the LRP by 2045 at recent (status quo) catch and effort levels. While the stock remains in a biologically healthy state, the prospects for any future albacore targeted fishery in the Solomon Islands will depend on local abundance, catch rates and economics.

¹Estimates of risk using deterministic projections are likely to be underestimated as they do not include future uncertainties such as fluctuations in recruitment.





Figure 5: Distribution of total South Pacific albacore catch by fishing method 2013–2022 (Red, pole-and-line; Blue, purse-seine; Green, longline; Orange, troll; Yellow, other) (top). Annual catch of South Pacific albacore in the WCPO by fishing method (middle). Percentage impact on spawning biomass due to fishing in the WCPO (bottom - left) and the model Region/s encompassing your EEZ (bottom - right).

Other species

Stock assessments are available for some other species which may interact with the fisheries in the Solomon Islands (??). These include three shark and three billfish species (Figure 6). Both north and south Pacific swordfish and blue marlin are not overfished and no overfishing was taking place. The SC, has recommended that there should be no increase in fishing mortality for blue marlin and swordfish. The southwest Pacific striped marlin assessment results indicate that the stock is fully exploited, is not experiencing overfishing, but may be overfished (??) **WCPFC2012 WCPFC2013b**. For the sharks that have been assessed to date, blue sharks in the north Pacific are not considered to be overfished and no overfishing is taking place; silky sharks are overfished and overfishing is taking place; and for oceanic whitetip sharks the stock is overfished and overfishing is occurring **WCPFC2012, WCPFC2013, WCPFC2014**. It should also be noted that due to concerns expressed by the Scientific Committee regarding the steep declines in biomass of silky and oceanic whitetip shark, the Commission has adopted CMMs for these species prohibiting their retention, transshipment, storage or landing **WCPFC2011a, WCPFC2013b**. In addition, in an attempt to reduce incidental catch of these species CMM2014-05 prohibits the use of either wire trace branchlines or shark lines on longline sets **this has now changed?**.





Figure 6: Kobe plots comparing the stock status of the key bycatch species caught in the WCPFC Convention Area. Where current fishing mortality rate exceeds the fishing mortality rate at MSY ($F/F_{MSY} > 1$) then overfishing is occurring. Where the current spawning biomass is less than the spawning biomass that would produce MSY, then the stock is overfished ($SB/SB_{MSY} < 1$).

References

- Castillo-Jordan, C., T Teears, J. Hampton, N Davies, J Scutt Phillips, S McKechnie, T Peatman, et al. 2022. "Stock Assessment of Skipjack Tuna in the Western and Central Pacific Ocean: 2022." WCPFC-SC18-2022/SA-WP-01 (REV5). Electronic meeting, 10–18 August 2022.
- Castillo-Jordan, C., H. Xu, J. Hampton, N. Ducharme-Barth, and P. Hamer. 2021. "Stock Assessment of South Pacific Albacore." WCPFC-SC17-2021/SA-WP-02. Electronic meeting, 11–19 August 2021.
- Day, J, A Magnusson, T Teears, J Hampton, N Davies, C Castillo-Jordan, T Peatman, et al. 2023. "Stock assessment of bigeye tuna in the western and central Pacific Ocean: 2023." WCPFC-SC19-2023/SA-WP-05. Koror, Palau, 16-24 August 2023.
- Hare, S., P. G. Williams, C. Castillo-Jordan, P. Hamer, J. Hampton, P. Lehodey, J. Macdonald, et al. 2023. "The western and central Pacific tuna fishery: 2022 overview and status of stocks." Tuna Fisheries Assessment Report no. 23. Online, 20 November 2023.
- Magnusson, A, J Day, T Teears, J Hampton, N Davies, C Castillo-Jordan, T Peatman, et al. 2023. "Stock assessment of yellowfin tuna in the western and central Pacific Ocean: 2023." WCPFC-SC19-2023/SA-WP-04. Koror, Palau, 16-24 August 2023.
- WCPFC. 2022. "Report of the Scientific Committee eighteenth regular session." Summary Report. Online, 10–18 August 2022: Commission for the Conservation; Management of Highly Migratory Fish Stocks in the Western; Central Pacific Ocean.
- . 2023. "Report of the Scientific Committee nineteenth regular session." Summary Report. Koror, Palau, 16-24 August 2023: Commission for the Conservation; Management of Highly Migratory Fish Stocks in the Western; Central Pacific Ocean.

Tables

Figures