
Network ablation effects across sampled semantic clusters
People & portraits (C7) Baseline 33.3%
Tennis action (C33) Baseline 24.5%
Sharks & marine life (C4) Baseline 18.6%
Consumer electronics (C36) Baseline 14.8%
Trains & rail transport (C22) Baseline 12.5%
Water towers (C18) Baseline 7.8%
Rural fields (C35) Baseline 4.8%
Emergency vehicles (C23) Baseline 3.3%
Religious architecture (C25) Baseline 2.9%
Home appliances (C11) Baseline 0.6%
Ventral visual/object −8.0 −8.7 −5.6 −6.3 −3.3 −1.6 −2.0 +0.4 −0.2 −0.3
Early visual −5.0 −6.7 −1.5 −1.6 −8.2 −4.2 −1.6 +0.4 −1.7 −0.6
Parietal attention −2.6 −8.0 −2.6 −1.2 −1.8 −0.9 −0.5 +0.4 −0.6 −0.3
Other +0.5 +1.6 −4.1 −1.5 −2.0 −1.4 −1.1 +0.4 −1.1 −0.4
Motion visual −1.8 −5.6 −1.1 +0.2 +0.2 +0.2 −0.3 −0.4 −0.4 +0.1
Frontal/control 0.0 +1.0 −0.6 −0.8 −1.7 −0.7 0.0 −0.7 −0.2 −0.3
Cerebellum −0.1 +0.3 −1.1 +0.3 +0.6 +0.2 +0.3 −0.4 −0.4 +0.1
Temporal/auditory −0.4 +0.4 +0.2 +0.4 −0.6 −0.4 +0.2 +0.4 −0.5 +0.1
Subcortical +0.3 +0.6 0.0 −0.1 −0.2 −1.2 +0.1 −0.7 −0.3 −0.1
+8.0
+6.0
+4.0
+2.0
0
−2.0
−4.0
−6.0
−8.0
Change in unique-stimulus Top-10 accuracy (ablated − baseline), percentage points
Negative values indicate impaired retrieval; positive values indicate improved retrieval after ablation.