

# From Fields to Trees

Firas Hamze, Nando De Freitas  
UAI 2004

Presented by: samy

February 20, 2014

From Fields to Trees

Firas, Hamze 2004

Covariance Structure of the Gibbs Sampler ...

Liu, Wong, Kong 1994

Markov Chain Strategies in Scientific Computing

Jun Liu, 2001

# The Problem

Given: Graphical Model with known Parameters.


**Goal:** Estimate  $\mathbb{E}[h(X)]$

E.g.:  $\mathbb{1}_A(x)$ ,  $\sum_i X_i$


Generate Samples:  $X^{(1)}, X^{(2)}, \dots, X^{(N)}$

Empirical Estimator

$$\mu_0 = \frac{1}{N} \sum_{i=1}^N h(X^{(i)})$$



# Gibbs Sampling



Current Sample:  $X^{(t)}$

$$X_1^{(t+1)} \mid X_{2,3,4,5,6}^{(t)}$$

$$X_2^{(t+1)} \mid X_1^{(t+1)}, X_{3,4,5,6}^{(t)}$$


...

$$X_6^{(t+1)} \mid X_{1,2,3,4,5}^{(t+1)}$$

Next Sample:  $X^{(t+1)}$

$X^{(1)} \rightarrow X^{(2)} \rightarrow \dots$  is a Markov Chain with eqib<sup>m</sup> distribution  $P$ .

# Blocked Gibbs Sampling



Current Sample:  $X^{(t)}$

$$X_{1,3}^{(t+1)} \mid X_{2,4,5,6}^{(t)}$$


$$X_{2,4,5}^{(t+1)} \mid X_{1,3}^{(t+1)}, X_6^{(t)}$$

$$X_6^{(t+1)} \mid X_{1,2,3,4,5}^{(t+1)}$$

Next Sample:  $X^{(t+1)}$

$X^{(1)} \rightarrow X^{(2)} \rightarrow \dots$  is a Markov Chain with eqlb<sup>m</sup> distribution  $P$ .

# Blocked Gibbs Sampling






Why Blocked Gibbs ?

Sampling from conditional distribution is difficult.

BUT, Chain mixes faster  
 $\Rightarrow$  better samples.

# How to Block (Partition) the Variables?



# Outline

- 1 Digression
  - MCMC Theory
  - Rao-Blackwellisation
- 2 Primary Idea
  - Tree Partitions
- 3 Hamze & Freitas, 2004
- 4 My 701 Course Project



$$X^{(1)} \rightarrow X^{(2)} \rightarrow \dots \rightarrow X^{(t)} \rightarrow X^{(t+1)} \rightarrow \dots$$

$\pi \leftarrow$  Equilibrium Distribution

$$X^{(1)} \rightarrow X^{(2)} \rightarrow \dots \rightarrow X^{(t)} \rightarrow X^{(t+1)} \rightarrow \dots$$

$\pi \leftarrow$  Equilibrium Distribution

$$L_0(\pi) = \{h : \Omega \rightarrow \mathbb{R} : \mathbb{E}_\pi h(X) = 0, \quad \mathbb{V}_\pi h(X) < \infty\}$$

$$\langle h, g \rangle = \text{Covar}_\pi(h(X), g(X))$$

$(L_0(\pi), \langle \cdot, \cdot \rangle)$  is a Hilbert Space.

$$X^{(1)} \rightarrow X^{(2)} \rightarrow \dots \rightarrow X^{(t)} \rightarrow X^{(t+1)} \rightarrow \dots$$

$\pi \leftarrow$  Equilibrium Distribution

$$L_0(\pi) = \{h : \Omega \rightarrow \mathbb{R} : \mathbb{E}_\pi h(X) = 0, \mathbb{V}_\pi h(X) < \infty\}$$

$$\langle h, g \rangle = \text{Covar}_\pi(h(X), g(X))$$

$(L_0(\pi), \langle \cdot, \cdot \rangle)$  is a Hilbert Space.

$$\|h\|^2 = \langle h, h \rangle = \text{Covar}_\pi(h(X), h(X)) = \mathbb{V}_\pi(h(X))$$

## Forward Operator

Define  $F : L(\pi) \rightarrow L(\pi)$ ,  $[Fh](z) = \mathbb{E}[h(X^{(1)}) | X^{(0)} = z]$

## Forward Operator

Define  $F : L(\pi) \rightarrow L(\pi)$ ,  $[Fh](z) = \mathbb{E}[h(X^{(1)}) | X^{(0)} = z]$

Norm of operator:  $\|F\| = \sup_{\|h\| \leq 1} \|Fh\|$ .

## Forward Operator

Define  $F : L(\pi) \rightarrow L(\pi)$ ,  $[Fh](z) = \mathbb{E}[h(X^{(1)})|X^{(0)} = z]$

Norm of operator:  $\|F\| = \sup_{\|h\| \leq 1} \|Fh\|$ .

$$\begin{aligned}\|Fh\|^2 &= \mathbb{V}_{X^{(0)} \sim \pi}(\mathbb{E}[h(X^{(1)})|X^{(0)}]) \\ &\leq \mathbb{V}_{X^{(0)} \sim \pi}(h(X^{(1)})) = \mathbb{V}_{X^{(1)} \sim \pi}(h(X^{(1)})) = \|h\|^2\end{aligned}$$

Hence,  $\|F\| \leq 1$ .

## Forward Operator

Define  $F : L(\pi) \rightarrow L(\pi)$ ,  $[Fh](z) = \mathbb{E}[h(X^{(1)}) | X^{(0)} = z]$

Norm of operator:  $\|F\| = \sup_{\|h\| \leq 1} \|Fh\|$ .

$$\begin{aligned}\|Fh\|^2 &= \mathbb{V}_{X^{(0)} \sim \pi}(\mathbb{E}[h(X^{(1)}) | X^{(0)}]) \\ &\leq \mathbb{V}_{X^{(0)} \sim \pi}(h(X^{(1)})) = \mathbb{V}_{X^{(1)} \sim \pi}(h(X^{(1)})) = \|h\|^2\end{aligned}$$

Hence,  $\|F\| \leq 1$ .

If MC is reversible,  $F$  is self-adjoint  $\implies \|F^n\| = \|F\|^n$ .

**Rate of Convergence:**  $\|F\|$

**Theorem:** Let  $X^{(0)} \sim P^{(0)}$ .  $P^{(n)}, \mathbb{E}^{(n)}$  :  $n^{\text{th}}$  step evolution.

$$|\mathbb{E}^{(n)} h(X) - \mathbb{E}_{\pi} h(X)| \leq C \|F\|^n \|h\|.$$



# Convergence of a MC

**Rate of Convergence:**  $\|F\|$

**Theorem:** Let  $X^{(0)} \sim P^{(0)}$ .  $P^{(n)}, \mathbb{E}^{(n)}$  :  $n^{\text{th}}$  step evolution.

$$|\mathbb{E}^{(n)} h(X) - \mathbb{E}_{\pi} h(X)| \leq C \|F\|^n \|h\|.$$

$$\|F\| = \sup_{f,g} \text{Corr}(f(X^{(t+1)}), g(X^{(t)}))$$

# Convergence of a MC

**Rate of Convergence:**  $\|F\|$

**Theorem:** Let  $X^{(0)} \sim P^{(0)}$ .  $P^{(n)}, \mathbb{E}^{(n)}$  :  $n^{\text{th}}$  step evolution.

$$|\mathbb{E}^{(n)} h(X) - \mathbb{E}_{\pi} h(X)| \leq C \|F\|^n \|h\|.$$

$$\|F\| = \sup_{f,g} \text{Corr}(f(X^{(t+1)}), g(X^{(t)}))$$

Finite State space:  $\|F\| \equiv 2^{\text{nd}}$  eigval of Transition Matrix.

$(X, Y) \sim P$ . Samples:  
 $\{(X^{(1)}, Y^{(1)}), (X^{(2)}, Y^{(2)}), \dots (X^{(N)}, Y^{(N)})\}$

**Goal:** Estimate  $\mathbb{E}_P[h(X)]$

Empirical Estimator  $\mu_0 = \frac{1}{N} \sum_{i=1}^N h(X^{(i)})$

Rao-Blackwellised Estimator  $\mu_{rb} = \frac{1}{N} \sum_{i=1}^N \mathbb{E} \left[ h(X) | Y^{(i)} \right]$

$(X, Y) \sim P$ . Samples:  
 $\{(X^{(1)}, Y^{(1)}), (X^{(2)}, Y^{(2)}), \dots (X^{(N)}, Y^{(N)})\}$

Independent samples: RB Estimator is better.

$(X, Y) \sim P$ . Samples:  
 $\{(X^{(1)}, Y^{(1)}), (X^{(2)}, Y^{(2)}), \dots (X^{(N)}, Y^{(N)})\}$

Independent samples: RB Estimator is better.


2-Stage Gibbs: RB Estimator is better (LKW 1994).

$$\mathbb{V}(\mu_0) - \mathbb{V}(\mu_{rb}) \leq \frac{\mathbb{V}(h(X))}{n^2} \left( n(1 - \|F\|) + 2(n-1)(\|F\| - \|F\|^2) \right. \\ \left. \dots + (\|F\|^{n-1} - \|F\|^n) \right)$$

# Outline


- 1 Digression
  - MCMC Theory
  - Rao-Blackwellisation
- 2 Primary Idea
  - Tree Partitions
- 3 Hamze & Freitas, 2004
- 4 My 701 Course Project

# Primary Idea



Correlated Partitions  $\implies$   
Successive Samples Correlated  $\implies$   
Large  $\|F\|$

# Primary Idea




Correlated Partitions  $\implies$   
Successive Samples Correlated  $\implies$   
Large  $\|F\|$

Blocked Gibbs :  $\|F\|$  is determined by the partitions.

Partition the Graph so that  $\|F\|$  is small !



# Primary Idea




Correlated Partitions  $\implies$   
Successive Samples Correlated  $\implies$   
Large  $\|F\|$

Blocked Gibbs :  $\|F\|$  is determined by the partitions.

Partition the Graph so that  $\|F\|$  is small !

Is this the best strategy ?


# Tree Partitions



Lets focus on Tree Partitions.

Computing conditional distribution on trees is easy.

# Tree Partitions



Lets focus on Tree Partitions.


Computing conditional distribution on trees is easy.

Best Tree Partition ? Not easy.


# Outline

- 1 Digression
  - MCMC Theory
  - Rao-Blackwellisation
- 2 Primary Idea
  - Tree Partitions
- 3 Hamze & Freitas, 2004
- 4 My 701 Course Project

# Grid Graph




Checkerboard



Two-Trees

# Results




H/deF use a a Rao-Blackwellised Estimator.

# Outline


- 1 Digression
  - MCMC Theory
  - Rao-Blackwellisation
- 2 Primary Idea
  - Tree Partitions
- 3 Hamze & Freitas, 2004
- 4 My 701 Course Project

Incorporate Correlations via Weights.





Incorporate Correlations via Weights.



Partitioning into  $K$  trees is NP-Hard ☹ (credit: Elara W)