

Cohesion in Rome

Fosco Loregian



December 1, 2019

Toposes

[...] vi el Aleph, desde todos los puntos,
vi en el Aleph la tierra, y en la tierra otra
vez el Aleph y en el Aleph la tierra, vi mi
cara y mis vísceras, vi tu cara, y sentí
vértigo y lloré...

JLB

Topos theory is a cornerstone of category theory linking together algebra, geometry and logic.

Toposes

[...] vi el Aleph, desde todos los puntos,
vi en el Aleph la tierra, y en la tierra otra
vez el Aleph y en el Aleph la tierra, vi mi
cara y mis vísceras, vi tu cara, y sentí
vértigo y lloré...

JLB

Topos theory is a cornerstone of category theory linking together algebra, geometry and logic.

Simply said, in each topos it is possible to re-enact the totality of known Mathematics; today we focus on

Toposes

[...] vi el Aleph, desde todos los puntos,
vi en el Aleph la tierra, y en la tierra otra
vez el Aleph y en el Aleph la tierra, vi mi
cara y mis vísceras, vi tu cara, y sentí
vértigo y lloré...

JLB

Topos theory is a cornerstone of category theory linking together algebra, geometry and logic.

Simply said, in each topos it is possible to re-enact the totality of known Mathematics; today we focus on

- Logic (better said, a fragment of **dependent type theory**)

Toposes

[...] vi el Aleph, desde todos los puntos,
vi en el Aleph la tierra, y en la tierra otra
vez el Aleph y en el Aleph la tierra, vi mi
cara y mis vísceras, vi tu cara, y sentí
vértigo y lloré...

JLB

Topos theory is a cornerstone of category theory linking together algebra, geometry and logic.

Simply said, in each topos it is possible to re-enact the totality of known Mathematics; today we focus on

- Logic (better said, a fragment of **dependent type theory**)
- Differential geometry

Toposes

[...] vi el Aleph, desde todos los puntos,
vi en el Aleph la tierra, y en la tierra otra
vez el Aleph y en el Aleph la tierra, vi mi
cara y mis vísceras, vi tu cara, y sentí
vértigo y lloré...

JLB

Topos theory is a cornerstone of category theory linking together algebra, geometry and logic.

Simply said, in each topos it is possible to re-enact the totality of known Mathematics; today we focus on

- Logic (better said, a fragment of **dependent type theory**)
- Differential geometry
- (Algebraic topology)

Toposes

[...] vi el Aleph, desde todos los puntos,
vi en el Aleph la tierra, y en la tierra otra
vez el Aleph y en el Aleph la tierra, vi mi
cara y mis vísceras, vi tu cara, y sentí
vértigo y lloré...

JLB

Topos theory is a cornerstone of category theory linking together algebra, geometry and logic.

Simply said, in each topos it is possible to re-enact the totality of known Mathematics; today we focus on

- Logic (better said, a fragment of **dependent type theory**)
- Differential geometry
- (Algebraic topology)
- ...

Definizione di fascio su uno spazio

Let (X, τ) be a topological space; a *sheaf on X* is a functor $F : \tau^{\text{op}} \rightarrow \underline{\text{Set}}$ such that for every $U \in \tau$ and every covering $\{U_i\}$ of U one has

- if $s, t \in FU$ are such that $s|_i = t|_i$ in FU_i for every $i \in I$, then $s = t$ in FU .

¹We denote $s|_i$ the image of $s \in FU$ under the nmeless map $FU \rightarrow FU_i$ induced by the inclusion $U_i \subseteq U$.

Definizione di fascio su uno spazio

Let (X, τ) be a topological space; a *sheaf on X* is a functor $F : \tau^{\text{op}} \rightarrow \underline{\text{Set}}$ such that for every $U \in \tau$ and every covering $\{U_i\}$ of U one has

- if $s, t \in FU$ are such that $s|_i = t|_i$ in FU_i for every $i \in I$, then $s = t$ in FU .
- if $s_i \in FU_i$ is a family of elements such that $s_i|_{ij} = s_j|_{ij}$, then there exists a $s \in FU$ such that $s|_i = s_i$.¹

¹We denote $s|_i$ the image of $s \in FU$ under the nmeless map $FU \rightarrow FU_i$ induced by the inclusion $U_i \subseteq U$.

Examples of sheaves

Every construction in Mathematics that exhibits a local character is a sheaf:

- sending $U \mapsto CU$, continuous functions with domain U
(similarly, differentiable, C^∞ , C^ω , holomorphic...)

Examples of sheaves

Every construction in Mathematics that exhibits a local character is a sheaf:

- sending $U \mapsto CU$, continuous functions with domain U
(similarly, differentiable, C^∞ , C^ω , holomorphic...)
- sending $U \mapsto \Omega^p U$, differential forms supported on U
(similarly: distributions, test functions...)

Examples of sheaves

Every construction in Mathematics that exhibits a local character is a sheaf:

- sending $U \mapsto CU$, continuous functions with domain U
(similarly, differentiable, C^∞ , C^ω , holomorphic...)
- sending $U \mapsto \Omega^P U$, differential forms supported on U
(similarly: distributions, test functions...)
- ... sending $U \mapsto \{f : U \rightarrow \mathbb{R} \mid f \text{ has property } P \text{ locally}\}$ for some P .

Examples of sheaves

Every construction in Mathematics that exhibits a local character is a sheaf:

- sending $U \mapsto CU$, continuous functions with domain U
(similarly, differentiable, C^∞ , C^ω , holomorphic...)
- sending $U \mapsto \Omega^P U$, differential forms supported on U
(similarly: distributions, test functions...)
- ... sending $U \mapsto \{f : U \rightarrow \mathbb{R} \mid f \text{ has property } P \text{ locally}\}$ for some P .

Every construction that does involve global properties, is not a sheaf:

Examples of sheaves

Every construction in Mathematics that exhibits a local character is a sheaf:

- sending $U \mapsto CU$, continuous functions with domain U
(similarly, differentiable, C^∞ , C^ω , holomorphic...)
- sending $U \mapsto \Omega^P U$, differential forms supported on U
(similarly: distributions, test functions...)
- ... sending $U \mapsto \{f : U \rightarrow \mathbb{R} \mid f \text{ has property } P \text{ locally}\}$ for some P .

Every construction that does involve global properties, is not a sheaf:

- sending $U \mapsto \{\text{bounded functions } f : U \rightarrow \mathbb{R}\}$

Examples of sheaves

Every construction in Mathematics that exhibits a local character is a sheaf:

- sending $U \mapsto CU$, continuous functions with domain U
(similarly, differentiable, C^∞ , C^ω , holomorphic...)
- sending $U \mapsto \Omega^P U$, differential forms supported on U
(similarly: distributions, test functions...)
- ... sending $U \mapsto \{f : U \rightarrow \mathbb{R} \mid f \text{ has property } P \text{ locally}\}$ for some P .

Every construction that does involve global properties, is not a sheaf:

- sending $U \mapsto \{\text{bounded functions } f : U \rightarrow \mathbb{R}\}$
- sending $U \mapsto \{L^1 \text{ functions } f : U \rightarrow \mathbb{R}\}$

Examples of sheaves

Every construction in Mathematics that exhibits a local character is a sheaf:

- sending $U \mapsto CU$, continuous functions with domain U
(similarly, differentiable, C^∞ , C^ω , holomorphic...)
- sending $U \mapsto \Omega^P U$, differential forms supported on U
(similarly: distributions, test functions...)
- ... sending $U \mapsto \{f : U \rightarrow \mathbb{R} \mid f \text{ has property } P \text{ locally}\}$ for some P .

Every construction that does involve global properties, is not a sheaf:

- sending $U \mapsto \{\text{bounded functions } f : U \rightarrow \mathbb{R}\}$
- sending $U \mapsto \{L^1 \text{ functions } f : U \rightarrow \mathbb{R}\}$
- ...

Grothendieck topologies

A *sieve* on an object X of a category \mathcal{C} is a subobject S of the hom functor $yX = \mathcal{C}(-, X)$;

Grothendieck topologies

A *sieve* on an object X of a category \mathcal{C} is a subobject S of the hom functor $yX = \mathcal{C}(-, X)$;

A *Grothendieck topology* on a category amounts to the choice of a family of *covering sieves* for every object $X \in \mathcal{C}$; this family of sieves is chosen in such a way that

Grothendieck topologies

A *sieve* on an object X of a category \mathcal{C} is a subobject S of the hom functor $yX = \mathcal{C}(-, X)$;

A *Grothendieck topology* on a category amounts to the choice of a family of *covering sieves* for every object $X \in \mathcal{C}$; this family of sieves is chosen in such a way that

- if $S \Rightarrow yX$ is a covering sieve and $f : Y \rightarrow X$ is a morphism of \mathcal{C} , then the morphism $f^*S \Rightarrow Y$ obtained in the pullback

$$\begin{array}{ccc} f^*S & \longrightarrow & S \\ \downarrow & \lrcorner & \downarrow \\ Y & \xrightarrow{f} & X \end{array}$$

is again a covering sieve.

Grothendieck topologies

- Let $S \Rightarrow yX$ be a covering sieve on X , and let T be any sieve on X . If for each object Y of \mathcal{C} and each arrow $f : Y \rightarrow X$ in SY the pullback sieve $f^* T$ is a covering sieve on Y , then T is a covering sieve on X .

Grothendieck topologies

- Let $S \Rightarrow yX$ be a covering sieve on X , and let T be any sieve on X . If for each object Y of \mathcal{C} and each arrow $f : Y \rightarrow X$ in SY the pullback sieve $f^* T$ is a covering sieve on Y , then T is a covering sieve on X .
- the identity $1 : yX \Rightarrow yX$ is a covering sieve.

Grothendieck topologies

- Let $S \Rightarrow yX$ be a covering sieve on X , and let T be any sieve on X . If for each object Y of \mathcal{C} and each arrow $f : Y \rightarrow X$ in SY the pullback sieve $f^* T$ is a covering sieve on Y , then T is a covering sieve on X .
- the identity $1 : yX \Rightarrow yX$ is a covering sieve.
 - if $\{U_i\}$ covers U , then for every $V \subseteq U$ $V \cap U_i$ covers V ;

Grothendieck topologies

- Let $S \Rightarrow yX$ be a covering sieve on X , and let T be any sieve on X . If for each object Y of \mathcal{C} and each arrow $f : Y \rightarrow X$ in SY the pullback sieve $f^* T$ is a covering sieve on Y , then T is a covering sieve on X .
- the identity $1 : yX \Rightarrow yX$ is a covering sieve.
 - if $\{U_i\}$ covers U , then for every $V \subseteq U$ $V \cap U_i$ covers V ;
 - if $\{U_i\}$ covers U and $\{V_{ij}\}$ covers U_i , then V_{ij} covers U ;

Grothendieck topologies

- Let $S \Rightarrow yX$ be a covering sieve on X , and let T be any sieve on X . If for each object Y of \mathcal{C} and each arrow $f : Y \rightarrow X$ in SY the pullback sieve $f^* T$ is a covering sieve on Y , then T is a covering sieve on X .
- the identity $1 : yX \Rightarrow yX$ is a covering sieve.
 - if $\{U_i\}$ covers U , then for every $V \subseteq U$ $V \cap U_i$ covers V ;
 - if $\{U_i\}$ covers U and $\{V_{ij}\}$ covers U_i , then V_{ij} covers U ;
 - $\{U\}$ covers U .

Grothendieck topologies

- Let $S \Rightarrow yX$ be a covering sieve on X , and let T be any sieve on X . If for each object Y of \mathcal{C} and each arrow $f : Y \rightarrow X$ in SY the pullback sieve $f^* T$ is a covering sieve on Y , then T is a covering sieve on X .
- the identity $1 : yX \Rightarrow yX$ is a covering sieve.
 - if $\{U_i\}$ covers U , then for every $V \subseteq U$ $V \cap U_i$ covers V ;
 - if $\{U_i\}$ covers U and $\{V_{ij}\}$ covers U_i , then V_{ij} covers U ;
 - $\{U\}$ covers U .

A Grothendieck site is a category with a Grothendieck topology, i.e. a function j that assigns to every object a family of covering sieves.

Grothendieck topologies

- Let $S \Rightarrow yX$ be a covering sieve on X , and let T be any sieve on X . If for each object Y of \mathcal{C} and each arrow $f : Y \rightarrow X$ in SY the pullback sieve $f^* T$ is a covering sieve on Y , then T is a covering sieve on X .
- the identity $1 : yX \Rightarrow yX$ is a covering sieve.
 - if $\{U_i\}$ covers U , then for every $V \subseteq U$ $V \cap U_i$ covers V ;
 - if $\{U_i\}$ covers U and $\{V_{ij}\}$ covers U_i , then V_{ij} covers U ;
 - $\{U\}$ covers U .

A Grothendieck site is a category with a Grothendieck topology, i.e. a function j that assigns to every object a family of covering sieves.

We denote a site as the pair (\mathcal{C}, j) .

Sheaves on a site

A *sheaf* on a small site \mathcal{C} is a functor $F : \mathcal{C}^{\text{op}} \rightarrow \underline{\text{Set}}$ such that for every covering sieve $R \rightarrow yU$ and every diagram

$$\begin{array}{ccc} R & \xrightarrow{f} & F \\ m \downarrow & \nearrow & \\ yU & & \end{array}$$

there is a unique dotted extension $yU \Rightarrow F$

Sheaves on a site

A *sheaf* on a small site \mathcal{C} is a functor $F : \mathcal{C}^{\text{op}} \rightarrow \underline{\text{Set}}$ such that for every covering sieve $R \rightarrow yU$ and every diagram

$$\begin{array}{ccc} R & \xrightarrow{f} & F \\ m \downarrow & \nearrow & \\ yU & & \end{array}$$

there is a unique dotted extension $yU \Rightarrow F$ (by the Yoneda lemma, this consists of a unique element $s \in FU$, **exercise**).

Sheaves on a site

A *sheaf* on a small site \mathcal{C} is a functor $F : \mathcal{C}^{\text{op}} \rightarrow \underline{\text{Set}}$ such that for every covering sieve $R \rightarrow yU$ and every diagram

$$\begin{array}{ccc} R & \xrightarrow{f} & F \\ m \downarrow & \nearrow & \\ yU & & \end{array}$$

there is a unique dotted extension $yU \Rightarrow F$ (by the Yoneda lemma, this consists of a unique element $s \in FU$, **exercise**).

The full subcategory of sheaves on a site (\mathcal{C}, j) is denoted $\text{Sh}(\mathcal{C}, j)$.

Giraud Theorem

By general facts on locally presentable categories, the subcategory of sheaves on a site is reflective via a functor

$$r : \text{Cat}(\mathcal{C}^{\text{op}}, \underline{\text{Set}}) \rightarrow \text{Sh}(\mathcal{C}, j)$$

called *sheafification* of a presheaf $F : \mathcal{C}^{\text{op}} \rightarrow \underline{\text{Set}}$.

Giraud Theorem

By general facts on locally presentable categories, the subcategory of sheaves on a site is reflective via a functor

$$r : \text{Cat}(\mathcal{C}^{\text{op}}, \underline{\text{Set}}) \rightarrow \text{Sh}(\mathcal{C}, j)$$

called *sheafification* of a presheaf $F : \mathcal{C}^{\text{op}} \rightarrow \underline{\text{Set}}$.

Historical note

Grothendieck was the first to note that in every topos of sheaves the **internal language** is sufficiently expressive to concoct **higher-order logic** and he strived to advertise his intuitions to an audience of logicians.

But it wasn't until Lawvere devised the notion of **elementary topos** that the community agreed on the potential of this theory.

Elementary toposes

An *elementary topos* is a category \mathcal{E} that

- it has finite limits;

Elementary toposes

An *elementary topos* is a category \mathcal{E} that

- it has finite limits;
- is cartesian closed;

Elementary toposes

An *elementary topos* is a category \mathcal{E} that

- it has finite limits;
- is cartesian closed;
- has a *subobject classifier*, i.e. an object $\Omega \in \mathcal{E}$ such that the functor $\text{Sub} : \mathcal{E}^{\text{op}} \rightarrow \underline{\text{Set}}$ sending A into the set of isomorphism classes of monomorphisms $\begin{smallmatrix} U \\ \downarrow \\ A \end{smallmatrix}$ is representable by the object Ω .

Elementary toposes

The natural bijection $\mathcal{E}(A, \Omega) \cong \text{Sub}(A)$ is obtained pulling back a “characteristic arrow” $\chi_U : A \rightarrow \Omega$ along a *universal arrow* $t : 1 \rightarrow \Omega$ to obtain the monic U , as in the diagram

$$\begin{array}{ccc} U & \longrightarrow & 1 \\ m \downarrow & \lrcorner & \downarrow t \\ A & \xrightarrow{\chi_m} & \Omega \end{array}$$

so, the bijection is induced by the maps

- $\chi_- : \left[\begin{smallmatrix} U \\ \downarrow \\ A \end{smallmatrix} \right] \mapsto \chi_m$ and

Elementary toposes

The natural bijection $\mathcal{E}(A, \Omega) \cong \text{Sub}(A)$ is obtained pulling back a “characteristic arrow” $\chi_U : A \rightarrow \Omega$ along a *universal arrow* $t : 1 \rightarrow \Omega$ to obtain the monic U , as in the diagram

$$\begin{array}{ccc} U & \longrightarrow & 1 \\ m \downarrow & \lrcorner & \downarrow t \\ A & \xrightarrow{\chi_m} & \Omega \end{array}$$

so, the bijection is induced by the maps

- $\chi_- : \begin{bmatrix} U \\ \downarrow \\ A \end{bmatrix} \mapsto \chi_m$ and
- $- \times_{\Omega} t : \chi_U \mapsto \chi_U \times_{\Omega} t.$

Grothendieck \subset elementary

En los libros herméticos está escrito que lo que hay abajo es igual a lo que hay arriba, y lo que hay arriba, igual a lo que hay abajo; en el Zohar, que el mundo inferior es reflejo del superior.[†]

JLB

- Every Grothendieck topos is elementary;

Grothendieck \subset elementary

En los libros herméticos está escrito que lo que hay abajo es igual a lo que hay arriba, y lo que hay arriba, igual a lo que hay abajo; en el Zohar, que el mundo inferior es reflejo del superior.[†]

JLB

- Every Grothendieck topos is elementary;
- An elementary topos is Grothendieck if and only if it is a locally finitely presentable category.

En los libros herméticos está escrito que lo que hay abajo es igual a lo que hay arriba, y lo que hay arriba, igual a lo que hay abajo; en el Zohar, que el mundo inferior es reflejo del superior.[†]

JLB

- Every Grothendieck topos is elementary;
- An elementary topos is Grothendieck if and only if it is a locally finitely presentable category.

Giraud theorem characterises Grothendieck toposes as such elementary toposes.

En los libros herméticos está escrito que lo que hay abajo es igual a lo que hay arriba, y lo que hay arriba, igual a lo que hay abajo; en el Zohar, que el mundo inferior es reflejo del superior.[†]

JLB

- Every Grothendieck topos is elementary;
- An elementary topos is Grothendieck if and only if it is a locally finitely presentable category.

Giraud theorem characterises Grothendieck toposes as such elementary toposes.

[†]Microcosm principle: a topos, i.e. a place where subobjects are well-behaved, is but a well-behaved subobject in the 2-category of presheaf categories.

Logic of categories

What is the internal logic of a category?

This would have deserved a dedicated seminar but:

- Every category \mathcal{C} is a universe in which we can interpret type theory;

This can be made precise in various ways:

<https://ncatlab.org/nlab/show/relation+between+type+theory+and+category+theory>

Logic of categories

What is the internal logic of a category?

This would have deserved a dedicated seminar but:

- Every category \mathcal{C} is a universe in which we can interpret type theory;
- every object $A \in \mathcal{C}$ is a type – a term in the type of types:

This can be made precise in various ways:

<https://ncatlab.org/nlab/show/relation+between+type+theory+and+category+theory>

Logic of categories

What is the internal logic of a category?

This would have deserved a dedicated seminar but:

- Every category \mathcal{C} is a universe in which we can interpret type theory;
- every object $A \in \mathcal{C}$ is a type – a term in the type of types;
- every morphism $X \rightarrow A$ is a (generalised) term of type A , in a context X .

This can be made precise in various ways:

<https://ncatlab.org/nlab/show/relation+between+type+theory+and+category+theory>

Logic of toposes

flavor of type theory	equivalent to	flavor of category theory	
intuitionistic propositional logic/simply-typed lambda calculus		cartesian closed category	
multiplicative intuitionistic linear logic		symmetric closed monoidal category	(various authors since ~68)
first-order logic		hyperdoctrine	(Seely 1984a)
classical linear logic		star-autonomous category	(Seely 89)
extensional dependent type theory		locally cartesian closed category	(Seely 1984b)
homotopy type theory without univalence (intensional M-L dependent type theory)		locally cartesian closed $(\infty, 1)$-category	(Cisinski 12- (Shulman 12))
homotopy type theory with higher inductive types and univalence		elementary $(\infty, 1)$-topos	see here
dependent linear type theory		indexed monoidal category (with comprehension)	(Vákár 14)

Axiomatic Cohesion

What is cohesion

Cohesion is the mutual attraction of molecules sticking together to form *droplets*, caused by mild electrical attraction between them.

What is cohesion

Cohesion is the mutual attraction of molecules sticking together to form *droplets*, caused by mild electrical attraction between them.

Figure 1: Droplets of mercury “exhibiting cohesion”

What is cohesion

Classes of geometric spaces exhibit similar coagulation properties,

What is cohesion

Classes of geometric spaces exhibit similar coagulation properties, similar to internal forces leading them to adhere and form **coherent conglomerates**.

What is cohesion

Classes of geometric spaces exhibit similar coagulation properties, similar to internal forces leading them to adhere and form **coherent conglomerates**. This behaviour is typical of **smooth spaces**.

What is cohesion

Classes of geometric spaces exhibit similar coagulation properties, similar to internal forces leading them to adhere and form **coherent conglomerates**. This behaviour is typical of **smooth spaces**.

Example

Smooth manifolds can be probed via smooth open balls and every smooth space is a “coherent conglomerate” of *cohesive pieces*.

What is cohesion

Classes of geometric spaces exhibit similar coagulation properties, similar to internal forces leading them to adhere and form **coherent conglomerates**. This behaviour is typical of **smooth spaces**.

Example

Smooth manifolds can be probed via smooth open balls and every smooth space is a “coherent conglomerate” of *cohesive pieces*.

Question

Which formal axioms describe the mathematics behind this intuition? What is *axiomatic cohesion*?

Axioms to answer this question have been devised by Lawvere [Law1] (worth reading, but quite mystical!).

Desiderata

We would like to operate in a *category* (a **topos**) of “cohesive spaces”, such that

Desiderata

We would like to operate in a *category* (a **topos**) of “cohesive spaces”, such that

- there is a functor $\Pi: \mathcal{H} \rightarrow \underline{\text{Set}}$ that sends every cohesive space $X \in \mathcal{H}$ into its set of **connected components**.

Desiderata

We would like to operate in a *category* (a **topos**) of “cohesive spaces”, such that

- there is a functor $\Pi: \mathcal{H} \rightarrow \underline{\text{Set}}$ that sends every cohesive space $X \in \mathcal{H}$ into its set of **connected components**.
- Every set $S \in \underline{\text{Set}}$ can be regarded as a cohesive space in two complementary ways:

Desiderata

We would like to operate in a *category* (a **topos**) of “cohesive spaces”, such that

- there is a functor $\Pi: \mathcal{H} \rightarrow \underline{\text{Set}}$ that sends every cohesive space $X \in \mathcal{H}$ into its set of **connected components**.
- Every set $S \in \underline{\text{Set}}$ can be regarded as a cohesive space in two complementary ways:
 - *discretely*, with a functor $\underline{\text{Set}} \rightarrow \mathcal{H}$ that regards every singleton of S as a cohesive droplet;

Desiderata

We would like to operate in a *category* (a **topos**) of “cohesive spaces”, such that

- there is a functor $\Pi: \mathcal{H} \rightarrow \underline{\text{Set}}$ that sends every cohesive space $X \in \mathcal{H}$ into its set of **connected components**.
- Every set $S \in \underline{\text{Set}}$ can be regarded as a cohesive space in two complementary ways:
 - *discretely*, with a functor $\underline{\text{Set}} \rightarrow \mathcal{H}$ that regards every singleton of S as a cohesive droplet;
 - *codiscretely*, with a functor $\underline{\text{Set}} \rightarrow \mathcal{H}$ that regards the whole S as an unseparable cohesive droplet.

Desiderata

We would like to operate in a *category* (a **topos**) of “cohesive spaces”, such that

- there is a functor $\Pi: \mathcal{H} \rightarrow \underline{\text{Set}}$ that sends every cohesive space $X \in \mathcal{H}$ into its set of **connected components**.
- Every set $S \in \underline{\text{Set}}$ can be regarded as a cohesive space in two complementary ways:
 - *discretely*, with a functor $\underline{\text{Set}} \rightarrow \mathcal{H}$ that regards every singleton of S as a cohesive droplet;
 - *codiscretely*, with a functor $\underline{\text{Set}} \rightarrow \mathcal{H}$ that regards the whole S as an unseparable cohesive droplet.
- Discretely and codiscretely cohesive spaces embed in \mathcal{H} , with fully faithful functors: in that

$$\mathcal{H}(\text{disc}(S), \text{disc}(T)) \cong \underline{\text{Set}}(S, T)$$

$$\mathcal{H}(\text{codisc}(S), \text{codisc}(T)) \cong \underline{\text{Set}}(S, T)$$

Axiomatic cohesion

An adjunction

$$\Pi \dashv \text{disc} \dashv \Gamma \dashv \text{codisc} : \mathcal{H}$$
$$\begin{array}{ccccc} & & \xrightarrow{\Pi} & & \\ & \xleftarrow{\text{disc}} & \perp & \xrightarrow{\perp} & \text{Set} \\ \xleftarrow{\Gamma} & & \perp & & \perp \\ & \xleftarrow{\text{codisc}} & & & \end{array}$$

exhibits the cohesion of \mathcal{H} over Set if

Axiomatic cohesion

An adjunction

$$\Pi \dashv \text{disc} \dashv \Gamma \dashv \text{codisc} : \mathcal{H} \begin{array}{c} \xrightarrow{\Pi} \\ \xleftarrow{\text{disc}} \quad \xrightarrow{\perp} \\ \xleftarrow{\Gamma} \quad \xrightarrow{\perp} \\ \xleftarrow{\text{codisc}} \end{array} \underline{\text{Set}}$$

exhibits the cohesion of \mathcal{H} over Set if

- disc and codisc are fully faithful;

Axiomatic cohesion

An adjunction

$$\Pi \dashv \text{disc} \dashv \Gamma \dashv \text{codisc} : \mathcal{H} \begin{array}{c} \xrightarrow{\quad \Pi \quad} \\ \xleftarrow{\quad \text{disc} \quad} \quad \perp \quad \xrightarrow{\quad \perp \quad} \\ \xleftarrow{\quad \Gamma \quad} \quad \perp \quad \xrightarrow{\quad \perp \quad} \\ \xleftarrow{\quad \text{codisc} \quad} \end{array} \underline{\text{Set}}$$

exhibits the cohesion of \mathcal{H} over Set if

- disc and codisc are fully faithful;
- the leftmost adjoint Π preserves finite products.

Axiomatic cohesion

An adjunction

$$\Pi \dashv \text{disc} \dashv \Gamma \dashv \text{codisc} : \mathcal{H} \begin{array}{c} \xrightarrow{\Pi} \\ \xleftarrow{\text{disc}} \quad \xrightarrow{\perp} \\ \xleftarrow{\Gamma} \quad \xrightarrow{\perp} \\ \xleftarrow{\text{codisc}} \end{array} \underline{\text{Set}}$$

exhibits the cohesion of \mathcal{H} over Set if

- disc and codisc are fully faithful;
- the leftmost adjoint Π preserves finite products.

(Γ “forgets cohesion”: it sends a space to its underlying set of points)

Formal fact. Every quadruple of adjoints induces a triple of adjoints.


Formal fact. Every quadruple of adjoints induces a triple of adjoints.

- There is an adjoint triple of idempotent co/monads on \mathcal{H} , induced by the cohesion:

$$\begin{array}{ccccc} & \xrightarrow{\Pi} & & \xrightarrow{\text{disc}} & \\ \mathcal{H} & \xleftarrow{\text{disc}} & \perp & \perp & \mathcal{H} \\ & \xleftarrow{\perp} & \text{Set} & \xleftarrow{\Gamma} & \perp \\ & \xrightarrow{\Gamma} & & \xrightarrow{\perp} & \perp \\ & & & \xrightarrow{\text{codisc}} & \end{array}$$

Formal fact. Every quadruple of adjoints induces a triple of adjoints.

- There is an adjoint triple of idempotent co/monads on \mathcal{H} , induced by the cohesion:



monad	comonad	monad
\int	$(-)^{\flat}$	$(-)^{\sharp}$
$\text{disc} \circ \Pi$	$\text{disc} \circ \Gamma$	$\text{codisc} \circ \Gamma$
pron.: <i>shape</i>	pron.: <i>flat</i>	pron.: <i>sharp</i>

Modalities, pieces

The triple of adjoints

$$\begin{array}{ccc} & \lrcorner & \\ \mathcal{H} & \xrightleftharpoons{\quad} & \mathcal{H} \\ & \sharp & \end{array}$$

is called the **shape, flat, sharp** string of “co/modalities”
(idempotent co/monads) for the cohesive topos \mathcal{H} .

The **shape** of $X \in \mathcal{H}$ is the discrete object on the “fundamental groupoid” of X . The adjunction $\Pi \dashv \text{disc}$ has something to do with (topological) Galois theory.

Modalities, pieces

1. The **flat** functor corresponds to the **object of flat connections** on $X \in \mathcal{H}$: if G is a group,

$$\left\{ \begin{array}{c} \text{principal} \\ \text{bundles on } X \end{array} \right\} \cong [X \rightarrow BG] \quad \left\{ \begin{array}{c} \text{flat con-} \\ \text{nections on } X \end{array} \right\} \cong \left\{ \begin{array}{c} \nearrow \text{flat} \\ X \xrightarrow{\quad} BG \end{array} \right\}$$

(keep in mind these equivalences: they will reappear later)

Modalities, pieces

1. The **flat** functor corresponds to the **object of flat connections** on $X \in \mathcal{H}$: if G is a group,

$$\left\{ \begin{array}{c} \text{principal} \\ \text{bundles on } X \end{array} \right\} \cong [X \rightarrow BG] \quad \left\{ \begin{array}{c} \text{flat con-} \\ \text{nections on } X \end{array} \right\} \cong \left\{ \begin{array}{c} \nearrow bBG \\ X \longrightarrow BG \\ \downarrow \end{array} \right\}$$

(keep in mind these equivalences: they will reappear later)

2. **sharp** of X , $\sharp X$, corresponds to the codiscrete object on the sets of **points** ΓX of X .

Modalities, pieces

1. The **flat** functor corresponds to the **object of flat connections** on $X \in \mathcal{H}$: if G is a group,

$$\left\{ \begin{smallmatrix} \text{principal} \\ \text{bundles on } X \end{smallmatrix} \right\} \cong [X \rightarrow BG] \quad \left\{ \begin{smallmatrix} \text{flat con-} \\ \text{nections on } X \end{smallmatrix} \right\} \cong \left\{ \begin{array}{c} \nearrow bBG \\ X \xrightarrow{\quad} BG \\ \downarrow \end{array} \right\}$$

(keep in mind these equivalences: they will reappear later)

2. **sharp** of X , $\sharp X$, corresponds to the codiscrete object on the sets of **points** ΓX of X .
3. Co/discrete objects are precisely the objects for which $bX \cong X$, resp. $\sharp Y \cong Y$.

Every object fits in a “complex”:

Definition

There is a canonical natural transformation

$$\sharp X \xrightarrow{\epsilon_{(\text{disc} \dashv \Gamma), X}} X \xrightarrow{\eta_{(\Pi \dashv \text{disc}), X}} \int X$$

called the “points to pieces” map;

Every object fits in a “complex”:

Definition

There is a canonical natural transformation

$$\sharp X \xrightarrow{\epsilon_{(\text{disc} \dashv \Gamma), X}} X \xrightarrow{\eta_{(\Pi \dashv \text{disc}), X}} \int X$$

called the “**points to pieces**” map; this map comes from a natural transformation

$$\alpha : \Gamma \Rightarrow \Pi$$

$$\alpha_X : \Gamma X \rightarrow \Pi X$$

Every object fits in a “complex”:

Definition

There is a canonical natural transformation

$$\sharp X \xrightarrow{\epsilon_{(\text{disc} \dashv \Gamma), X}} X \xrightarrow{\eta_{(\Pi \dashv \text{disc}), X}} \int X$$

called the “**points to pieces**” map; this map comes from a natural transformation

$$\alpha : \Gamma \Rightarrow \Pi$$

$$\alpha_X : \Gamma X \rightarrow \Pi X$$

It is a “comparison” between the action of Γ (send X into its “sections” or “set of points”) and Π (send X into its “pieces” or “components”).

- We say that **pieces have points** in the cohesive topos \mathcal{H} (or that “ \mathcal{H} satisfies *Nullstellensatz*”) if the points-to-pieces transformation $\alpha_X : \Gamma X \rightarrow \Pi X$ is surjective for all $X \in \mathcal{H}$.

- We say that **pieces have points** in the cohesive topos \mathcal{H} (or that “ \mathcal{H} satisfies *Nullstellensatz*”) if the points-to-pieces transformation $\alpha_X : \Gamma X \rightarrow \Pi X$ is surjective for all $X \in \mathcal{H}$.
- We say that **discrete is concrete** in \mathcal{H} if natural transformation whose components are

$$\text{disc}(S) \rightarrow \text{codisc}(\Gamma(\text{disc}(S))) \cong \text{codisc}(S)$$

is a monomorphism (discrete cohesion sits into codiscrete cohesion).

- We say that **pieces have points** in the cohesive topos \mathcal{H} (or that “ \mathcal{H} satisfies *Nullstellensatz*”) if the points-to-pieces transformation $\alpha_X : \Gamma X \rightarrow \Pi X$ is surjective for all $X \in \mathcal{H}$.
- We say that **discrete is concrete** in \mathcal{H} if natural transformation whose components are

$$\text{disc}(S) \rightarrow \text{codisc}(\Gamma(\text{disc}(S))) \cong \text{codisc}(S)$$

is a monomorphism (discrete cohesion sits into codiscrete cohesion).

- We say that \mathcal{H} has **contractible subobjects** or has **sufficient cohesion** if $\Pi(\Omega) \cong *$. This implies that for all $X \in \mathcal{H}$ also $\Pi(\Omega^X) \cong *$.

- We say that **pieces have points** in the cohesive topos \mathcal{H} (or that “ \mathcal{H} satisfies *Nullstellensatz*”) if the points-to-pieces transformation $\alpha_X : \Gamma X \rightarrow \Pi X$ is surjective for all $X \in \mathcal{H}$.
- We say that **discrete is concrete** in \mathcal{H} if natural transformation whose components are

$$\text{disc}(S) \rightarrow \text{codisc}(\Gamma(\text{disc}(S))) \cong \text{codisc}(S)$$

is a monomorphism (discrete cohesion sits into codiscrete cohesion).

- We say that \mathcal{H} has **contractible subobjects** or has **sufficient cohesion** if $\Pi(\Omega) \cong *$. This implies that for all $X \in \mathcal{H}$ also $\Pi(\Omega^X) \cong *$.
- ... and many others (see [Law]).

Proposition

- The adjunctions $\Pi \dashv \text{disc}$ and $\Gamma \dashv \text{codisc}$ exhibit the subcategories of discrete and codiscrete objects as reflective subcategories of \mathcal{H} ; these subcategories form **exponential ideals** in \mathcal{H} .
- If \mathcal{H} exhibits cohesion, then Set is equivalent to the full subcategory of \mathcal{H} whose objects are the X such that $\eta_{(\Gamma \dashv \text{codisc}), X} : X \rightarrow \text{codisc}(\Gamma(X))$ is an isomorphism.

Equivalently: every cohesive topos “contains the trivial cohesion of disconnected pieces ($= \underline{\text{Set}}$)”.

Non-trivial fact

Whenever a topos arises, there's an interaction between logic and geometry.

Definition

A monomorphism $\psi: S \rightarrow A$ in a cohesive topos \mathcal{H} is a proposition of type A in the internal logic of \mathcal{H} . We say that ψ is *discretely true* if the pullback $\psi^*(S) \rightarrow A$

$$\begin{array}{ccc} \psi^*(S) & \xrightarrow{\quad} & \flat S \\ \downarrow & \lrcorner & \downarrow \flat \psi \\ A & \xrightarrow[\eta]{} & \flat A \end{array}$$

is an isomorphism in \mathcal{H} , where $\eta: A \rightarrow \flat A$ is the \flat -unit of the flat monad.

- Discrete truth specifies a mode/modality in which a proposition can be true. Propositions true over all discrete objects (i.e., such that $b\psi$ is an iso) are discretely true.

- Discrete truth specifies a mode/modality in which a proposition can be true. Propositions true over all discrete objects (i.e., such that $b\psi$ is an iso) are discretely true.
- Let $\mathcal{H} = \text{Sh}(\text{Cart}, J)$ be the topos of sheaves over cartesian spaces ($\text{hom}(m, n) = \text{smooth maps } \mathbb{R}^n \rightarrow \mathbb{R}^m$) is cohesive.

- Discrete truth specifies a mode/modality in which a proposition can be true. Propositions true over all discrete objects (i.e., such that $b\psi$ is an iso) are discretely true.
- Let $\mathcal{H} = \text{Sh}(\text{Cart}, J)$ be the topos of sheaves over cartesian spaces ($\text{hom}(m, n) = \text{smooth maps } \mathbb{R}^n \rightarrow \mathbb{R}^m$) is cohesive.
- Let $\psi: Z^p(U) \hookrightarrow \Omega^p(U)$ be the proposition in \mathcal{H} given by “the p -form ω is closed on a neighbourhood U_x of a point”. Then ψ is discretely true (“every form is closed over a discrete space”).

Examples

The Sierpiński topos

Let $\mathcal{C} = \{0 \rightarrow 1\}$ be the interval category with a unique non-identity arrow.

The category of **presheaves** on \mathcal{C} forms a topos $\mathcal{H} = \underline{\text{Set}}^{\mathcal{C}}$ = arrows in Set, that exhibits cohesion:

The Sierpiński topos

Let $\mathcal{C} = \{0 \rightarrow 1\}$ be the interval category with a unique non-identity arrow.

The category of presheaves on \mathcal{C} forms a topos $\mathcal{H} = \underline{\text{Set}}^{\mathcal{C}}$ = arrows in Set, that exhibits cohesion:

- the functor \sqcap sends an object $S \rightarrow I$ to its codomain I ;

The Sierpiński topos

Let $\mathcal{C} = \{0 \rightarrow 1\}$ be the interval category with a unique non-identity arrow.

The category of presheaves on \mathcal{C} forms a topos $\mathcal{H} = \underline{\text{Set}}^{\mathcal{C}}$ = arrows in Set, that exhibits cohesion:

- the functor Π sends an object $S \rightarrow I$ to its codomain I ;
- the functor Γ sends an object $S \rightarrow I$ to its domain S ;

The Sierpiński topos

Let $\mathcal{C} = \{0 \rightarrow 1\}$ be the interval category with a unique non-identity arrow.

The category of presheaves on \mathcal{C} forms a topos $\mathcal{H} = \underline{\text{Set}}^{\mathcal{C}}$ = arrows in Set, that exhibits cohesion:

- the functor Π sends an object $S \rightarrow I$ to its codomain I ;
- the functor Γ sends an object $S \rightarrow I$ to its domain S ;
- the functor disc sends a set K into the identity $1: K \rightarrow K$;

The Sierpiński topos

Let $\mathcal{C} = \{0 \rightarrow 1\}$ be the interval category with a unique non-identity arrow.

The category of presheaves on \mathcal{C} forms a topos $\mathcal{H} = \underline{\text{Set}}^{\mathcal{C}}$ = arrows in Set, that exhibits cohesion:

- the functor Π sends an object $S \rightarrow I$ to its codomain I ;
- the functor Γ sends an object $S \rightarrow I$ to its domain S ;
- the functor disc sends a set K into the identity $1: K \rightarrow K$;
- the functor codisc sends a set K into its terminal morphism $K \rightarrow *$.

The Sierpiński topos

Evidently these functors form an adjunction ($\Pi \dashv \text{disc} \dashv \Gamma \dashv \text{codisc}$) so that \mathcal{H} exhibits cohesion; this matches our intuition, in that

- The “points to pieces” transformation sends $f : S \rightarrow I$ into $S = \Gamma(f) \rightarrow \Pi(f) = I$;
- $\text{disc}(K)$ “keeps all the pieces of K maximally distinguished” and
- $\text{codisc}(K)$ “lumps all the pieces of K together”.

Pointed categories exhibit cohesion

Let \mathcal{C} small and with a terminal object. Then there exists a triple

$$\begin{array}{ccc} & \xrightarrow{\lim\limits_{\longrightarrow}} & \\ [\mathcal{C}^{\text{op}}, \underline{\text{Set}}] & \xleftarrow{\text{const}} & \underline{\text{Set}} \\ & \xrightarrow{\lim\limits_{\longleftarrow}} & \end{array}$$

that extends to $\lim\limits_{\longleftarrow} \dashv \mathcal{K}$:

$$S \xrightarrow{\mathcal{K}} \left(c \mapsto \underline{\text{Set}}(\mathcal{C}(*, c), S) \right)$$

(Dually, if \mathcal{C} has an initial object...)

Pointed categories exhibit cohesion

Let \mathcal{C} small and with a terminal object. Then there exists a triple

$$\begin{array}{ccc} & \xrightarrow{\lim\limits_{\longrightarrow}} & \\ [\mathcal{C}^{\text{op}}, \underline{\text{Set}}] & \xleftarrow{\text{const}} & \underline{\text{Set}} \\ & \xrightarrow{\lim\limits_{\longleftarrow}} & \end{array}$$

that extends to $\lim\limits_{\longleftarrow} \dashv \aleph$:

$$S \xrightarrow{\aleph} \left(c \mapsto \underline{\text{Set}}(\mathcal{C}(*, c), S) \right)$$

(Dually, if \mathcal{C} has an initial object...)

Proposition

If \mathcal{C} has both an initial and a terminal object (e.g. it is pointed) then $[\mathcal{C}^{\text{op}}, \underline{\text{Set}}]$ exhibits cohesion with

$$(\lim\limits_{\longrightarrow} \dashv \text{const} \dashv \lim\limits_{\longleftarrow} \dashv \aleph) : [\mathcal{C}^{\text{op}}, \underline{\text{Set}}] \xrightleftharpoons{\text{const}} \underline{\text{Set}}$$

Reflexive directed graphs

Consider the category \mathcal{C}

$$\begin{array}{ccc} & s & \\ 0 & \xrightleftharpoons[q]{\quad} & 1 \\ & t & \end{array}$$

such that $0 \rightrightarrows 1 \rightarrow 0$ is a reflexive coequalizer;

The category $[\mathcal{C}^{\text{op}}, \underline{\text{Set}}]$ is the category of **reflexive directed graphs** RDGph, and it exhibits cohesion, since the terminal geometric morphism

$$(\text{disc} \dashv \Gamma) : \text{RDGph} \leftrightarrows \underline{\text{Set}}$$

extends on the left with $\Pi : X \mapsto \text{coeq}(X_1 \rightrightarrows X_0)$ (simply the connected components of the graph).

Since this is a reflexive coequalizer, it preserves products.

(exercise: define disc, codisc)

Simplicial sets

Proposition

Let Δ be the simplex category having objects nonempty finite ordinals and morphisms monotone maps. The topos

$\mathcal{H} = [\Delta^{\text{op}}, \underline{\text{Set}}]$ exhibits cohesion, and in \mathcal{H} pieces have points.

- $\Gamma = (-)_0$ sends a simplicial set X into its set of 0-simplices X_0

Simplicial sets

Proposition

Let Δ be the simplex category having objects nonempty finite ordinals and morphisms monotone maps. The topos

$\mathcal{H} = [\Delta^{\text{op}}, \underline{\text{Set}}]$ exhibits cohesion, and in \mathcal{H} pieces have points.

- $\Gamma = (-)_0$ sends a simplicial set X into its set of 0-simplices X_0
- $\Pi = \pi_0$ sends a simplicial set X into its set of connected components $\text{coeq}(X_1 \rightrightarrows X_0)$.

Simplicial sets

Proposition

Let Δ be the simplex category having objects nonempty finite ordinals and morphisms monotone maps. The topos

$\mathcal{H} = [\Delta^{\text{op}}, \underline{\text{Set}}]$ exhibits cohesion, and in \mathcal{H} pieces have points.

- $\Gamma = (-)_0$ sends a simplicial set X into its set of 0-simplices X_0
- $\Pi = \pi_0$ sends a simplicial set X into its set of connected components $\text{coeq}(X_1 \rightrightarrows X_0)$.
- disc sends a set S into the constant simplicial set in S having constant set of simplices and identities as faces and degeneracies.

Simplicial sets

Proposition

Let Δ be the simplex category having objects nonempty finite ordinals and morphisms monotone maps. The topos

$\mathcal{H} = [\Delta^{\text{op}}, \underline{\text{Set}}]$ exhibits cohesion, and in \mathcal{H} pieces have points.

- $\Gamma = (-)_0$ sends a simplicial set X into its set of 0-simplices X_0
- $\Pi = \pi_0$ sends a simplicial set X into its set of connected components $\text{coeq}(X_1 \rightrightarrows X_0)$.
- disc sends a set S into the constant simplicial set in S having constant set of simplices and identities as faces and degeneracies.
- codisc sends a set S into the simplicial set whose n -simplices are $(n + 1)$ -tuples of elements of S (and faces and degeneracies forget and add elements accordingly).

Tangent cohesion

Consider the codomain fibration

$$\mathcal{C}^{\rightarrow} \xrightarrow{p} \mathcal{C}$$

of a finitely complete category \mathcal{C} , sending an arrow $f: X \rightarrow Y$ to its codomain. The fiber $p^{-1}(Y)$ is canonically isomorphic to the category \mathcal{C}/Y of arrows over Y .

There exists a fibration $T\mathcal{C} \rightarrow \mathcal{C}$ having typical fiber the fiberwise abelianization of \mathcal{C}/Y , i.e. the category $\text{Ab}(\mathcal{C}/Y)$ of abelian groups in \mathcal{C}/Y .

Tangent cohesion

Consider the codomain fibration

$$\mathcal{C}^{\rightarrow} \xrightarrow{p} \mathcal{C}$$

of a finitely complete category \mathcal{C} , sending an arrow $f: X \rightarrow Y$ to its codomain. The fiber $p^{\leftarrow}(Y)$ is canonically isomorphic to the category \mathcal{C}/Y of arrows over Y .

There exists a fibration $T\mathcal{C} \rightarrow \mathcal{C}$ having typical fiber the fiberwise abelianization of \mathcal{C}/Y , i.e. the category $\text{Ab}(\mathcal{C}/Y)$ of abelian groups in \mathcal{C}/Y .

(hint: un/straighten the prestack $\mathcal{C} \rightarrow \text{Cat}: Y \mapsto \text{Ab}(\mathcal{C}/Y)$).

Tangent cohesion

Consider the codomain fibration

$$\mathcal{C}^{\rightarrow} \xrightarrow{p} \mathcal{C}$$

of a finitely complete category \mathcal{C} , sending an arrow $f: X \rightarrow Y$ to its codomain. The fiber $p^{-1}(Y)$ is canonically isomorphic to the category \mathcal{C}/Y of arrows over Y .

There exists a fibration $T\mathcal{C} \rightarrow \mathcal{C}$ having typical fiber the fiberwise abelianization of \mathcal{C}/Y , i.e. the category $\text{Ab}(\mathcal{C}/Y)$ of abelian groups in \mathcal{C}/Y .

(hint: un/straighten the prestack $\mathcal{C} \rightarrow \text{Cat}: Y \mapsto \text{Ab}(\mathcal{C}/Y)$).

Proposition

If \mathcal{C} is locally presentable, then so is $T\mathcal{C}$; moreover, the projection $q: T\mathcal{C} \rightarrow \mathcal{C}$ creates co/limits.

Tangent cohesion

Consider the codomain fibration

$$\mathcal{C}^{\rightarrow} \xrightarrow{p} \mathcal{C}$$

of a finitely complete category \mathcal{C} , sending an arrow $f: X \rightarrow Y$ to its codomain. The fiber $p^{\leftarrow}(Y)$ is canonically isomorphic to the category \mathcal{C}/Y of arrows over Y .

There exists a fibration $T\mathcal{C} \rightarrow \mathcal{C}$ having typical fiber the fiberwise abelianization of \mathcal{C}/Y , i.e. the category $\text{Ab}(\mathcal{C}/Y)$ of abelian groups in \mathcal{C}/Y .

(hint: un/straighten the prestack $\mathcal{C} \rightarrow \text{Cat}: Y \mapsto \text{Ab}(\mathcal{C}/Y)$).

Proposition

If \mathcal{C} is a topos over \mathcal{S} , then so is $T\mathcal{C}$; moreover, the projection $q: T\mathcal{C} \rightarrow \mathcal{C}$ creates co/limits.

Tangent cohesion

Proposition

There is a functor $\delta: T\mathcal{C} \rightarrow \mathcal{C}$ giving for each morphism in $T\mathcal{C}$ its domain. This functor is a right adjoint to a functor $\Omega: \mathcal{C} \rightarrow T\mathcal{C}$ that is also a *section* for q .

The object $\Omega(A)$ can be thought as the complex of **differential forms** on an internal abelian group $A \in \text{Ab}(\mathcal{C}/X)$.

Proposition

There is a functor $\delta: T\mathcal{C} \rightarrow \mathcal{C}$ giving for each morphism in $T\mathcal{C}$ its domain. This functor is a right adjoint to a functor $\Omega: \mathcal{C} \rightarrow T\mathcal{C}$ that is also a *section* for q .

The object $\Omega(A)$ can be thought as the complex of **differential forms** on an internal abelian group $A \in \text{Ab}(\mathcal{C}/X)$.

In classical differential geometry a leading theorem is that the co/tangent bundle to a smooth manifold is itself a smooth manifold. Here we can prove that

Tangent cohesion

Proposition

There is a functor $\delta: T\mathcal{C} \rightarrow \mathcal{C}$ giving for each morphism in $T\mathcal{C}$ its domain. This functor is a right adjoint to a functor $\Omega: \mathcal{C} \rightarrow T\mathcal{C}$ that is also a *section* for q .

The object $\Omega(A)$ can be thought as the complex of **differential forms** on an internal abelian group $A \in \text{Ab}(\mathcal{C}/X)$.

In classical differential geometry a leading theorem is that the co/tangent bundle to a smooth manifold is itself a smooth manifold. Here we can prove that

Proposition

If \mathcal{H} is a cohesive topos with cohesion $(\Pi \dashv \text{disc} \dashv \Gamma \dashv \text{codisc})$, then the tangent category is itself a cohesive topos.

Infinitesimal cohesion

Neighbourhoods of some spaces are “infintesimally extended around a single (global) point”. Cohesive structure can be refined to capture this phenomenon.

Infinitesimal cohesion

Let \mathcal{H} be cohesive. An **infinitesimal thickening** of \mathcal{H} is a new cohesive topos $\tilde{\mathcal{H}}$ linked to the previous by a quadruple of adjoints

$$\begin{array}{ccc} & \mathcal{H} & \\ (i_! \dashv i^* \dashv i_! \dashv i^!) & \downarrow & \\ & \tilde{\mathcal{H}} & \end{array}$$

Infinitesimal cohesion

Let \mathcal{H} be cohesive. An **infinitesimal thickening** of \mathcal{H} is a new cohesive topos $\tilde{\mathcal{H}}$ linked to the previous by a quadruple of adjoints

$$\begin{array}{ccc} & \mathcal{H} & \\ (i_! \dashv i^* \dashv i_! \dashv i^!) & \swarrow \quad \uparrow & \\ & \tilde{\mathcal{H}} & \end{array}$$

Infinitesimal cohesion

Let \mathcal{H} be cohesive. An **infinitesimal thickening** of \mathcal{H} is a new cohesive topos $\tilde{\mathcal{H}}$ linked to the previous by a quadruple of adjoints

$$\begin{array}{ccc} & \mathcal{H} & \\ (i_! \dashv i^* \dashv i_! \dashv i^!) & \Downarrow & \Updownarrow \\ & \tilde{\mathcal{H}} & \end{array}$$

Infinitesimal cohesion

Let \mathcal{H} be cohesive. An **infinitesimal thickening** of \mathcal{H} is a new cohesive topos $\tilde{\mathcal{H}}$ linked to the previous by a quadruple of adjoints

$$\begin{array}{ccc} & \mathcal{H} & \\ (i_! \dashv i^* \dashv i_! \dashv i^!) & \Downarrow & \Downarrow \\ & \tilde{\mathcal{H}} & \end{array}$$

Infinitesimal cohesion

Let \mathcal{H} be cohesive. An **infinitesimal thickening** of \mathcal{H} is a new cohesive topos $\tilde{\mathcal{H}}$ linked to the previous by a quadruple of adjoints

$$\begin{array}{ccc} & \mathcal{H} & \\ (i_! \dashv i^* \dashv i_! \dashv i^!) & \Downarrow & \Updownarrow \\ & \tilde{\mathcal{H}} & \end{array}$$

such that i_* , $i_!$ are fully faithful and $i_!$ commutes with finite products.

If such a structure exists, \mathcal{H} “exhibits **infinitesimal cohesion**”.

Infinitesimal cohesion

Let \mathcal{H} be cohesive. An **infinitesimal thickening** of \mathcal{H} is a new cohesive topos $\tilde{\mathcal{H}}$ linked to the previous by a quadruple of adjoints

$$\begin{array}{ccccc} & & \mathcal{H} & & \\ & \swarrow & \uparrow & \searrow & \uparrow \\ (i_! \dashv i^* \dashv i_* \dashv i^!) & \downarrow & & \downarrow & \uparrow \\ & & \tilde{\mathcal{H}} & & \end{array}$$

such that i_* , $i_!$ are fully faithful and $i_!$ commutes with finite products.

If such a structure exists, \mathcal{H} “exhibits **infinitesimal cohesion**”.

The functor i_* is fully faithful as well, and then we can consider \mathcal{H} “sitting nicely” inside its thickening $\tilde{\mathcal{H}}$.

Infinitesimal cohesion

- The cohesion exhibited by $\tilde{\mathcal{H}}$ factors through that of \mathcal{H} , in that

$$(\Pi_{\tilde{\mathcal{H}}} \dashv \text{disc}_{\tilde{\mathcal{H}}} \dashv \Gamma_{\tilde{\mathcal{H}}}) : \quad \begin{array}{ccccc} \tilde{\mathcal{H}} & \xrightleftharpoons[i^*]{\hspace{-1cm}} & \mathcal{H} & \xrightleftharpoons[\hspace{-1cm}]{} & \underline{\text{Set}} \\ & \xleftarrow[i_*]{\hspace{-1cm}} & & \xleftarrow[\text{disc}]{\hspace{-1cm}} & \\ & \xrightarrow[i^!]{\hspace{-1cm}} & & \xrightarrow[\Gamma]{\hspace{-1cm}} & \end{array}$$

Infinitesimal cohesion

- The cohesion exhibited by $\tilde{\mathcal{H}}$ factors through that of \mathcal{H} , in that

$$(\Pi_{\tilde{\mathcal{H}}} \dashv \text{disc}_{\tilde{\mathcal{H}}} \dashv \Gamma_{\tilde{\mathcal{H}}}) : \tilde{\mathcal{H}} \begin{array}{c} \xrightarrow{i^*} \\[-1ex] \xleftarrow{i_*} \\[-1ex] \xrightarrow{i^!} \end{array} \mathcal{H} \begin{array}{c} \xrightarrow{\Pi} \\[-1ex] \xleftarrow{\text{disc}} \\[-1ex] \xrightarrow{\Gamma} \end{array} \underline{\text{Set}}$$

- Infinitesimal cohesion describes formally infinitesimally extended neighbourhoods: if the functor i^* is interpreted as a contraction of a fat point onto its singleton, then $X \in \tilde{\mathcal{H}}$ is infinitesimal if $i^*(X) \cong *$.

Infinitesimal cohesion

- The cohesion exhibited by $\tilde{\mathcal{H}}$ factors through that of \mathcal{H} , in that

$$(\Pi_{\tilde{\mathcal{H}}} \dashv \text{disc}_{\tilde{\mathcal{H}}} \dashv \Gamma_{\tilde{\mathcal{H}}}) : \tilde{\mathcal{H}} \begin{array}{c} \xrightarrow{i^*} \\[-1ex] \xleftarrow{i_*} \\[-1ex] \xrightarrow{i^!} \end{array} \mathcal{H} \begin{array}{c} \xrightarrow{\Pi} \\[-1ex] \xleftarrow{\text{disc}} \\[-1ex] \xrightarrow{\Gamma} \end{array} \underline{\text{Set}}$$

- Infinitesimal cohesion describes formally infinitesimally extended neighbourhoods: if the functor i^* is interpreted as a contraction of a fat point onto its singleton, then $X \in \tilde{\mathcal{H}}$ is infinitesimal if $i^*(X) \cong *$. This motivates the fact that

$$\tilde{\mathcal{H}}(*, X) \cong \tilde{\mathcal{H}}(i_!(*), X) \cong \mathcal{H}(*, i^*(X)) \cong \mathcal{H}(*, *) \cong *$$

so that \mathcal{H} sees X as a “small neighbourhood concentrated around a single point $*_X$ ”.

Higher order cohesion: jet spaces

Most examples of infinitesimal cohesion come equipped with an infinite chain of thickening approximations.

Consider the **infinitesimal shape modality** $\Im := i_* i^*$
(it comes equipped with other two adjoints, $\Re \dashv \Im \dashv \&$)²

²This is the same general fact inducing $\int \dashv b \dashv \sharp$ adjunction.

Higher order cohesion: jet spaces

Most examples of infinitesimal cohensions come equipped with an infinite chain of thickening approximations.

Consider the **infinitesimal shape modality** $\Im := i_* i^*$
(it comes equipped with other two adjoints, $\Re \dashv \Im \dashv \&$)²

In several cases (like **smooth manifolds**) we have a **chain** of infinitesimal thickenings

$$\begin{array}{ccccccc} \widetilde{\mathcal{H}}_0 & \xrightarrow{i^{*(0)}} & \widetilde{\mathcal{H}}_1 & \xrightarrow{i^{*(1)}} & \widetilde{\mathcal{H}}_2 & \xrightarrow{i^{*(2)}} & \dots \\ \xleftarrow{i_{*,(0)}} & & \xleftarrow{i_{*,(1)}} & & \xleftarrow{i_{*,(2)}} & & \\ & & & & & & \\ \widetilde{\mathcal{H}}_\infty & \xrightarrow{i^{*(\infty)}} & \mathcal{H} & & \mathcal{H} & & \end{array}$$

²This is the same general fact inducing $\int \dashv \flat \dashv \sharp$ adjunction.

Higher order cohesion: jet spaces

Most examples of infinitesimal cohensions come equipped with an infinite chain of thickening approximations.

Consider the **infinitesimal shape modality** $\Im := i_* i^*$
(it comes equipped with other two adjoints, $\Re \dashv \Im \dashv \&$)²

In several cases (like **smooth manifolds**) we have a **chain** of infinitesimal thickenings

$$\begin{array}{ccccccc} \widetilde{\mathcal{H}}_0 & \xrightarrow{i^{*(0)}} & \widetilde{\mathcal{H}}_1 & \xrightarrow{i^{*(1)}} & \widetilde{\mathcal{H}}_2 & \xrightarrow{i^{*(2)}} & \dots \\ \xleftarrow{i_{*,(0)}} & & \xleftarrow{i_{*,(1)}} & & \xleftarrow{i_{*,(2)}} & & \\ & & & & & & \\ \widetilde{\mathcal{H}}_\infty & \xrightarrow{i^{*(\infty)}} & \mathcal{H} & & & & \end{array}$$

here we speak of a **sequence of orders of differential structures**.

²This is the same general fact inducing $\int \dashv \flat \dashv \sharp$ adjunction.

Higher order cohesion: jet spaces

Each of these approximations comes equipped with an *order k infinitesimal shape modality* $\mathfrak{S}^{(k)}X$ in a sequence

$$X \rightarrow \mathfrak{S}X = \mathfrak{S}^{(0)}X \rightarrow \mathfrak{S}^{(1)}X \rightarrow \mathfrak{S}^{(2)}X \rightarrow \dots$$

Example: Every cohesive topos exhibits infinitesimal cohesion via its **tangent** cohesive topos. This cohesion extends to any order of differential structure (“cohesive jet spaces”).

One can go **way** further, but the terminology becomes pretty dire:

Remark 2.2.13. The perspective of def. 2.2.12 has been highlighted in [Law91], where it is proposed (p. 7) that adjunctions of this form usefully formalize “many instances of the *Unity and Identity of Opposites*” that control Hegelian metaphysics [He1841].

[DCCT170811], 1040 pages of Hegel-ish mathematics

uses axiomatic cohesion of ∞ -toposes to axiomatise string theory.

One can go **way** further, but the terminology becomes pretty dire:

Remark 2.2.13. The perspective of def. 2.2.12 has been highlighted in [Law91], where it is proposed (p. 7) that adjunctions of this form usefully formalize “many instances of the *Unity and Identity of Opposites*” that control Hegelian metaphysics [He1841].

[DCCT170811], 1040 pages of Hegel-ish mathematics

uses axiomatic cohesion of ∞ -toposes to axiomatise string theory.

With Aufhebung.

Supergroupoids: rheonomy

We can speak of **supergroupoids** and show that certain categories of supersmooth manifolds exhibit cohesion (but not over Set. . .):

$$\begin{array}{c} \text{SuperSmoothS} \\ \uparrow d \quad \uparrow \Gamma \quad \uparrow c \\ \Pi \downarrow \quad \downarrow \quad \downarrow \\ \text{SuperS} \end{array}$$

The quadruple of adjoints generates the triple

$$\Rightarrow \dashv \rightsquigarrow \dashv \text{Rh}$$

(in some sense “fermions” \dashv “bosons”)

Supergroup: rheonomy

There is a “quadruple-to-triple” pattern here:



de Rham cohomology in cohesion

- Let \mathcal{H} be a cohesive topos, and $0 \rightarrow A$ a pointed object (e.g. an internal abelian group); then, A fits into a pullback square

$$\begin{array}{ccc} b_{dR}A & \longrightarrow & bA \\ \downarrow & \lrcorner & \downarrow \\ 0 & \longrightarrow & A \end{array}$$

where $b_{dR}A$ is the **object of coefficients** for de Rham cohomology.

- Let $X \in \mathcal{H}$ any object; we define $\int_{dR} X$ to be the pushout

$$\begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ \int X & \longrightarrow & \int_{dR} X \end{array}$$

where $\int_{dR} X$ is the **de Rham object** associated to X .

de Rham cohomology in cohesion

There is an adjunction

$$\begin{array}{ccc} */\mathcal{H} & \xrightarrow{\flat_{dR}} & \mathcal{H} \\ & \xleftarrow{\sharp_{dR}} & \end{array}$$

The mapping space $*/\mathcal{H}(\sharp_{dR} X, A) \cong \mathcal{H}(X, \flat_{dR} A)$ is called the de Rham space of X with coefficients in A and denoted $\mathbf{H}_{dR}^0(X, A)$.

de Rham cohomology in cohesion

Consider the pullback defining $\flat_{dR} A$ and apply the limit-preserving functor $\mathcal{H}(X, -)$: the square

$$\begin{array}{ccc} \mathcal{H}(X, \flat_{dR} A) & \longrightarrow & \mathcal{H}(X, \flat A) \\ \downarrow & \lrcorner & \downarrow \\ 0 & \longrightarrow & \mathcal{H}(X, A) \end{array}$$

remains a pullback and the object $\mathcal{H}(X, \flat A)$ identifies to **A-valued differential forms**, and the maps $X \rightarrow \flat_{dR} A$ are the **flat** ones: under the $\int \dashv \flat$ adjunction, a map $X \rightarrow \flat A$ mates to a smooth map

$$\int X \rightarrow A, \text{ corresponding to a square } \begin{array}{ccc} X & \xrightarrow{\quad} & 0 \\ \downarrow & & \downarrow \\ \int X & \rightarrow & A \end{array}$$

(minimal) Bibliography

- Law1 Lawvere, F. William. *Axiomatic cohesion*. Theory Appl. Categ 19.3 (2007): 41-49.
- Law2 Lawvere, F. William. *Categorical dynamics*. Topos theoretic methods in geometry 30 (1979): 1-28.
- Law3 Lawvere, F. William. *Toposes of laws of motion*. American Mathematical Society, Transcript from video, Montreal-September 27 (1997): 1997.
- Urs Schreiber, Urs. *Differential cohomology in a cohesive infinity-topos*. arXiv:1310.7930 (2013).
- Men Menni, Matias. *Continuous cohesion over sets*. Theory and Applications of Categories 29.20 (2014): 542-568.
- MLM MacLane, Saunders, and Ieke Moerdijk. *Sheaves in geometry and logic: A first introduction to topos theory*. Springer Science & Business Media, 2012.