
Fosco Loregian **TAL
TECH**

May 8, 2020

- Ph.D. at SISSA - Trieste 




Stable homotopy theory, ∞ -categories, derived AG





- Ph.D. at SISSA - Trieste 






Stable homotopy theory, ∞ -categories, derived AG







- University of Western Ontario 

∞ -categories, derivators

- Ph.D. at SISSA - Trieste 
Stable homotopy theory, ∞ -categories, derived AG
- University of Western Ontario 
 ∞ -categories, derivators
- Masaryk University 
Accessible categories, derivators, 2-categories

- Ph.D. at SISSA - Trieste 
Stable homotopy theory, ∞ -categories, derived AG
- University of Western Ontario 
 ∞ -categories, derivators
- Masaryk University 
Accessible categories, derivators, 2-categories
- Max Planck Inst. für Math. 
2-categories, derivators, applied category theory

- Ph.D. at SISSA - Trieste 
Stable homotopy theory, ∞ -categories, derived AG
- University of Western Ontario 
 ∞ -categories, derivators
- Masaryk University 
Accessible categories, derivators, 2-categories
- Max Planck Inst. für Math. 
2-categories, derivators, applied category theory
- Centro de Matemática da Universidade de Coimbra 
2-categories; finishing my first book

- Ph.D. at SISSA - Trieste 
Stable homotopy theory, ∞ -categories, derived AG
- University of Western Ontario 
 ∞ -categories, derivators
- Masaryk University 
Accessible categories, derivators, 2-categories
- Max Planck Inst. für Math. 
2-categories, derivators, applied category theory
- Centro de Matemática da Universidade de Coimbra 
2-categories; finishing my first book
- Tallinna Tehnikaülikooli 
2-categories; functorial semantics; categorical probability theory and its applications

STABLE HOMOTOPY THEORY

∞ -categories: a thickening of the notion of category, suitable for homotopy-invariant mathematics (algebraic geometry, algebraic topology type theory).

Turns out some parts of Mathematics are easier if stated in these terms:

- homological algebra : the scary part of algebraic topology
higher algebra: the linear algebra of ∞ -categories
- 1-topos theory : a synthetic type theory,
 ∞ -topos theory: a synthetic homotopy theory of homotopy types

A stable ∞ -category is an ∞ -category

- with all finite limits and colimits
- such that a square is cartesian iff cocartesian

The homotopy category of a stable ∞ -cat is always triangulated.

A stable ∞ -category is an ∞ -category

- with all finite limits and colimits
- such that a square is cartesian iff cocartesian

The homotopy category of a stable ∞ -cat is always triangulated.

The correspondence sending an abelian category \mathcal{A} into its derived category has a nice and clear universal property stated in terms of the heart of a canonical t -structure.

A stable ∞ -category is an ∞ -category

- with all finite limits and colimits
- such that a square is cartesian iff cocartesian

The homotopy category of a stable ∞ -cat is always triangulated.

The correspondence sending an abelian category \mathcal{A} into its derived category has a nice and clear universal property stated in terms of the heart of a canonical t -structure.

Stable, rational, p -adic, ... homotopy theory become pieces of the commutative algebra of ∞ -categories

A t-structure on a triangulated \mathcal{D} is a pair of triangulated subcategories of \mathcal{D} such that every object X lies in a sequence

$$X_{\leq} \rightarrow X \rightarrow X_{\geq} \rightarrow X_{\leq}[1]$$

[FL14] : On stable ∞ -categories a t-structure is a factorization system (E, M)

- such that E and M are 3-for-2 classes

A t-structure on a triangulated \mathcal{D} is a pair of triangulated subcategories of \mathcal{D} such that every object X lies in a sequence

$$X_{\leq} \rightarrow X \rightarrow X_{\geq} \rightarrow X_{\leq}[1]$$

[FL14] : On stable ∞ -categories a t-structure is a factorization system (E, M)

- such that E and M are 3-for-2 classes
- thus the category of E -cofibrant objects is coreflective

A t-structure on a triangulated \mathcal{D} is a pair of triangulated subcategories of \mathcal{D} such that every object X lies in a sequence

$$X_{\leq} \rightarrow X \rightarrow X_{\geq} \rightarrow X_{\leq}[1]$$

[FL14] : On stable ∞ -categories a t-structure is a factorization system (E, M)

- such that E and M are 3-for-2 classes
- thus the category of E -cofibrant objects is coreflective
- and the category of M -fibrant objects is reflective

A t-structure on a triangulated \mathcal{D} is a pair of triangulated subcategories of \mathcal{D} such that every object X lies in a sequence

$$X_{\leq} \rightarrow X \rightarrow X_{\geq} \rightarrow X_{\leq}[1]$$

[FL14] : On stable ∞ -categories a t-structure is a factorization system (E, M)

- such that E and M are 3-for-2 classes
- thus the category of E -cofibrant objects is coreflective
- and the category of M -fibrant objects is reflective
- cof/fib replacement = neg/pos truncation

A t-structure on a triangulated \mathcal{D} is a pair of triangulated subcategories of \mathcal{D} such that every object X lies in a sequence

$$X_{\leq} \rightarrow X \rightarrow X_{\geq} \rightarrow X_{\leq}[1]$$

[FL14] : On stable ∞ -categories a t-structure is a factorization system (E, M)

- such that E and M are 3-for-2 classes
- thus the category of E -cofibrant objects is coreflective
- and the category of M -fibrant objects is reflective
- cof/fib replacement = neg/pos truncation

(Algebraic) geometry reduces to a piece of categorical algebra.

Plan: redo t -structures

- [FL15] The set of t -structures has a natural choice of \mathbb{Z} -action (\mathbb{Z} = the integers); so, study \mathbb{Z} -equivariant monotone maps from a poset J to $TS(C)$.

This describes Bridgeland stability manifolds, and Postnikov towers on ∞ -toposes.

Plan: redo t -structures

- [FL15] The set of t -structures has a natural choice of \mathbb{Z} -action (\mathbb{Z} = the integers); so, study \mathbb{Z} -equivariant monotone maps from a poset J to $TS(C)$.

This describes Bridgeland stability manifolds, and Postnikov towers on ∞ -toposes.

- [FL15b] Every stratified manifold (X, \mathfrak{s}) generates a pair of t -structure that can be glued together
recollements, stratified schemes, representation of algebras

Plan: redo t -structures

- The space of slicings on a stable ∞ -category is a metric space and

Plan: redo t -structures

- The space of **slicings** on a stable ∞ -category is a metric space and
- The space

$$\{J: \mathbb{R} \rightarrow TS(\mathcal{C}) \mid J \text{ is Sorgenfrey continuous}\}$$

is an interesting set [**L-PhD**]

Plan: redo t -structures

- The space of slicings on a stable ∞ -category is a metric space and
- The space

$$\{J: \mathbb{R} \rightarrow TS(\mathcal{C}) \mid J \text{ is Sorgenfrey continuous}\}$$

is an interesting set [L-PhD]

Conjecture

Study

$$\{J: \operatorname{Spec}(\mathbb{Z}) \rightarrow TS(\mathcal{C}) \mid J \text{ is Zariski continuous}\}$$

to get something about motivic t -structure.

Plan: redo t -structures

- The space of **slicings** on a stable ∞ -category is a metric space and
- The space

$$\{J: \mathbb{R} \rightarrow TS(\mathcal{C}) \mid J \text{ is Sorgenfrey continuous}\}$$

is an interesting set [**L-PhD**]

Conjecture

Study


$$\{J: \operatorname{Spec}(\mathbb{Z}) \rightarrow TS(\mathcal{C}) \mid J \text{ is Zariski continuous}\}$$

to get something about motivic t -structure.

Conjecture

Blakers-Massey in positive characteristic is a theorem about factorization systems.

Morse theory as a theory of FS



Fact: $\text{Bord}(n)$ is the free (∞, n) -symmoncat on the point.

Monoidal
functors $Z : \text{Bord}(n) \rightarrow \text{Vect}$
are completely classified.

Morse theory is the theory of suitable factorization systems on $\text{Bord}(n)$.

critical points of a Morse function correspond to

critical values [L-PhD, Ch.7] of a certain slicing $J : \mathbb{R} \rightarrow \text{FS}(\text{Bord}(n))$.

DERIVATORS

A derivator is a strict 2-functor

$$\mathbb{D} : \mathbf{Cat}^{\mathrm{op}} \rightarrow \mathbf{CAT}$$

satisfying stacky conditions. They form the 2-category \mathbf{Der} .

They subsume most of ∞ -category theory; in particular, their stable homotopy.

A derivator is a strict 2-functor

$$\mathbb{D} : \mathbf{Cat}^{\mathrm{op}} \rightarrow \mathbf{CAT}$$

satisfying stacky conditions. They form the 2-category \mathbf{Der} .

They subsume most of ∞ -category theory; in particular, their stable homotopy.

[LV17] : A t-structure on a stable derivator is still a certain kind of factorization system; FS are still strict 2-algebras for the "squaring" 2-monad $(_)^2 : \mathbf{A} \rightarrow \mathbf{A}^2$ (see [KT93])

A derivator is a strict 2-functor

$$\mathbb{D} : \mathbf{Cat}^{\mathrm{op}} \rightarrow \mathbf{CAT}$$

satisfying stacky conditions. They form the 2-category \mathbf{Der} .

They subsume most of ∞ -category theory; in particular, their stable homotopy.

[LV17] : A t-structure on a stable derivator is still a certain kind of factorization system; FS are still strict 2-algebras for the "squaring" 2-monad $(_)^2 : \mathbf{A} \mapsto \mathbf{A}^2$ (see [KT93])

[Lor18] : reflective subderivators correspond to reflective factorization systems, and to algebras for idempotent monads (the "formal theory of monads" [S80] still holds in \mathbf{Der} , a monad $T : D \rightarrow D$ is just defined objectwise)

There is a **Yoneda structure**¹ on the 2-category of derivators

If this is true

¹A 2-categorical device to encode the calculus of pointwise Kan extensions.

There is a **Yoneda structure**¹ on the 2-category of derivators

If this is true

- comprehensive account of various notions of **accessible** and **locally presentable** derivator using the theory of locally presentable objects in a Yoneda structure, done in [DLL18]; categorical logic for derivators (see Prest's treatment of **definability** for module categories); derivator topos theory?

¹A 2-categorical device to encode the calculus of pointwise Kan extensions.

There is a **Yoneda structure**¹ on the 2-category of derivators

If this is true

- comprehensive account of various notions of **accessible** and **locally presentable** derivator using the theory of locally presentable objects in a Yoneda structure, done in [DLL18]; categorical logic for derivators (see Prest's treatment of **definability** for module categories); derivator topos theory?
- a convincing form of **adjoint functor theorem** for derivators; existence of a six-operation calculus for mappings of derivators. 2-categorical account of Grothendieck duality complicated diagrams...

¹A 2-categorical device to encode the calculus of pointwise Kan extensions.

There is a **Yoneda structure**¹ on the 2-category of derivators


If this is true

- comprehensive account of various notions of **accessible** and **locally presentable** derivator using the theory of locally presentable objects in a Yoneda structure, done in [DLL18]; categorical logic for derivators (see Prest's treatment of **definability** for module categories); derivator topos theory?
- a convincing form of **adjoint functor theorem** for derivators; existence of a six-operation calculus for mappings of derivators. 2-categorical account of Grothendieck duality complicated diagrams...
- **profunctors** between derivators; fibered derivators; **operads** in derivator theory; applications in representation theory of algebras, stable homotopy, ...

¹A 2-categorical device to encode the calculus of pointwise Kan extensions.


COENDS AND DG-STUFF

I have written a book on **coend calculus**, soon to appear under Cambridge University Press LNMs:




- Coends $\int_C T$ are universal objects associated to $T : C^{op} \times C \rightarrow D$, treated as integrals (a “Fubini rule” is valid).

I have written a book on **coend calculus**, soon to appear under Cambridge University Press LNMs:



- Coends $\int_C T$ are universal objects associated to $T : \mathcal{C}^{op} \times \mathcal{C} \rightarrow \mathcal{D}$, treated as integrals (a “Fubini rule” is valid).
- They find application in (monoidal) category theory, algebraic topology, algebraic geometry, categorical logic, representation theory (see ch.7 for an application to **DG-categories**), functional programming...

I have written a book on **coend calculus**, soon to appear under Cambridge University Press LNMs:



- Coends $\int_C T$ are universal objects associated to $T : \mathcal{C}^{op} \times \mathcal{C} \rightarrow \mathcal{D}$, treated as integrals (a “Fubini rule” is valid).
- They find application in (monoidal) category theory, algebraic topology, algebraic geometry, categorical logic, representation theory (see ch.7 for an application to **DG-categories**), functional programming...
- The book is being extensively cited (45 citations on Scholar May 8, 2020)

In [L20, 7.2.2]

For example: if \mathcal{A} is any dg-category its identity profunctor $\mathcal{A} \rightsquigarrow \mathcal{A}$ is a functor $\mathcal{A}^{\text{op}} \boxtimes \mathcal{A} \rightarrow \text{Ch}(\mathbb{Z})$, so that the coherent end

$$\oint_{\mathcal{A}} \mathcal{A}(A, A) \tag{7.82}$$

i.e. the object of derived natural transformations of the identity functor $\text{id}_{\mathcal{A}}$, recovers the *Hochschild complex* of \mathcal{A} . Then, if \mathcal{A} is an associative algebra regarded as a one-object dg-category concentrated in degree zero, the object $H^n(\int_* A)$ is the *Hochschild cohomology* of A , understood in the classical sense of, say, [Pie82, Ch. 11].

In [L20, 7.2.2]

For example: if \mathcal{A} is any dg-category its identity profunctor $\mathcal{A} \rightsquigarrow \mathcal{A}$ is a functor $\mathcal{A}^{\text{op}} \boxtimes \mathcal{A} \rightarrow \text{Ch}(\mathbb{Z})$, so that the coherent end

$$\oint_{\mathcal{A}} \mathcal{A}(A, A) \quad (7.82)$$

i.e. the object of derived natural transformations of the identity functor $\text{id}_{\mathcal{A}}$, recovers the *Hochschild complex* of \mathcal{A} . Then, if \mathcal{A} is an associative algebra regarded as a one-object dg-category concentrated in degree zero, the object $H^n(\int_* \mathcal{A})$ is the *Hochschild cohomology* of A , understood in the classical sense of, say, [Pie82, Ch. 11].

Applications to Kuznetsov-Lunts **categorical resolutions of singularities**: a smooth DG-category is a \mathcal{D} such that its identity profunctor $h : \mathcal{D} \rightsquigarrow \mathcal{D}$ is a perfect object (read as: a variety is smooth if the diagonal map $\Delta : X \rightarrow X \times X$ is smooth)

TEACHING AND ORGANIZATIONAL ACTIVITIES

- 2015 A short course on **model categories** at the University of Pavia

- 2015 A short course on **model categories** at the University of Pavia
- 2015 and 2019 Attendee and speaker at the **Kan Seminar I** (a webinar on category theory), and **Applied Category Theory 2019** (a webinar on applied category theory, from which the paper [MLR⁺20] stemmed)

- 2015 A short course on **model categories** at the University of Pavia
- 2015 and 2019 Attendee and speaker at the **Kan Seminar I** (a webinar on category theory), and **Applied Category Theory** 2019 (a webinar on applied category theory, from which the paper [MLR⁺20] stemmed)
- 2016 “**Elements of Finite Mathematics**” at UWO (mostly statistics to kinesiologists)

- 2015 A short course on **model categories** at the University of Pavia
- 2015 and 2019 Attendee and speaker at the **Kan Seminar I** (a webinar on category theory), and **Applied Category Theory** 2019 (a webinar on applied category theory, from which the paper [MLR⁺20] stemmed)
- 2016 “**Elements of Finite Mathematics**” at UWO (mostly statistics to kinesiologists)
- 2016 **Advisor** of a BSc thesis @unibo, “Elementary aspects of adjoint functors”

- 2015 A short course on **model categories** at the University of Pavia
- 2015 and 2019 Attendee and speaker at the **Kan Seminar I** (a webinar on category theory), and **Applied Category Theory** 2019 (a webinar on applied category theory, from which the paper [MLR⁺20] stemmed)
- 2016 “**Elements of Finite Mathematics**” at UWO (mostly statistics to kinesiologists)
- 2016 **Advisor** of a BSc thesis @unibo, “Elementary aspects of adjoint functors”
- 2018 **Organiser** of the 103rd Peripathetic Seminar on Sheaves and Logic, Brno

- 2015 A short course on **model categories** at the University of Pavia
- 2015 and 2019 Attendee and speaker at the **Kan Seminar I** (a webinar on category theory), and **Applied Category Theory** 2019 (a webinar on applied category theory, from which the paper [MLR⁺20] stemmed)
- 2016 “**Elements of Finite Mathematics**” at UWO (mostly statistics to kinesiologists)
- 2016 **Advisor** of a BSc thesis @unibo, “Elementary aspects of adjoint functors”
- 2018 **Organiser** of the 103rd Peripathetic Seminar on Sheaves and Logic, Brno
- 2020 **Category theory** course Teacher @taltech

- 2015 A short course on **model categories** at the University of Pavia
- 2015 and 2019 Attendee and speaker at the **Kan Seminar I** (a webinar on category theory), and **Applied Category Theory** 2019 (a webinar on applied category theory, from which the paper [MLR⁺20] stemmed)
- 2016 “**Elements of Finite Mathematics**” at UWO (mostly statistics to kinesiologists)
- 2016 **Advisor** of a BSc thesis @unibo, “Elementary aspects of adjoint functors”
- 2018 **Organiser** of the 103rd Peripathetic Seminar on Sheaves and Logic, Brno
- 2020 **Category theory** course Teacher @taltech
- 2019 and 2020 **Organiser** of ItaCa (**Italian Category** theorists) in Milan and on zoom, due to COVID19.

- 2015 A short course on **model categories** at the University of Pavia
- 2015 and 2019 Attendee and speaker at the **Kan Seminar I** (a webinar on category theory), and **Applied Category Theory** 2019 (a webinar on applied category theory, from which the paper [MLR⁺20] stemmed)
- 2016 “**Elements of Finite Mathematics**” at UWO (mostly statistics to kinesiologists)
- 2016 **Advisor** of a BSc thesis @unibo, “Elementary aspects of adjoint functors”
- 2018 **Organiser** of the 103rd Peripathetic Seminar on Sheaves and Logic, Brno
- 2020 **Category theory** course Teacher @taltech
- 2019 and 2020 **Organiser** of ItaCa (**Italian Category** theorists) in Milan and on zoom, due to COVID19.
- Reviewer for zbMath and AMS.

Miscellaneous projects

- Homotopy theory and set theory: [DL18] no homotopy category of a model category is “concrete”; what about ∞ -categories?

Miscellaneous projects

- Homotopy theory and set theory: [DL18] no homotopy category of a model category is “concrete”; what about ∞ -categories?
- Functional programming and type theory: HoTT, linear types, proof-checkers, categorical algebra in relational database architecture; natural language processing using category theory...

Miscellaneous projects

- Homotopy theory and set theory: [DL18] no homotopy category of a model category is “concrete”; what about ∞ -categories?
- Functional programming and type theory: HoTT, linear types, proof-checkers, categorical algebra in relational database architecture; natural language processing using category theory...
- Categorical logic and foundations of mathematics; functorial semantics à la Lawvere, but sprinkled with operads

Miscellaneous projects

- Homotopy theory and set theory: [DL18] no homotopy category of a model category is “concrete”; what about ∞ -categories?
- Functional programming and type theory: HoTT, linear types, proof-checkers, categorical algebra in relational database architecture; natural language processing using category theory...
- Categorical logic and foundations of mathematics; functorial semantics à la Lawvere, but sprinkled with operads
- more in detail, “2-semantics” of algebraic theories: profunctorial PROPs and theories, categorical algebra of cartesian bicategories...

