

# An Introduction to Proofs

Hassium, Fung San Gaan, Tingyu Wu

1 Basic Logic	7 Groups
2 Some Axioms of Sets	8 Rings
3 Functions and Well-Ordering	9 Real Numbers
4 Integers and Cardinality	10 Topological Spaces
5 Vector Spaces and Linear Maps	11 Categorical Reformulations
6 Functions as Morphisms	Alphabetical Index

In higher-level mathematics, students need a certain level of “mathematical maturity” to understand and apply abstract ideas. However, there is no clear way to measure this maturity, nor a definitive method to teach someone how to write a proof. This note is intended to be a transition from elementary math to proof-based formal mathematics. The beauty of mathematics is hidden behind the craft of abstraction and the search to common forms between entities. It is not merely about techniques of solving problems, but about a way of thinking. As Poincaré famously said, “Mathematics is the art of giving the same name to different things.”

The first section is a quick review of basic logic, the language we used in proofs. Sections 2 to 4 are set-theoretic constructions that are essential in any kind of math. Those sections discuss the axioms of sets in ZFC and their consequences. This part largely follows from [?] and [?]. In the later sections, those properties will be frequently used.

Section 5 is an introduction to linear algebra. The next two sections are introductory abstract algebra. We include discussions on groups, rings, and relations between rings and matrices. Section 8 is an algebraic construction of the real numbers  $\mathbb{R}$ . This section is entirely adopted from professor Lerman’s Math 347 lectures given in . We will also use the construction to verify some analytic properties of  $\mathbb{R}$ . Section 9 talks about general topology. This section includes several examples of topologies and the Tikhonov theorem. In the last two sections, we offer a more general theory to think about every entity in mathematics, the theory of categories. Our goal is to reformulate some results we proved before in a more elegant and general way by using categorical definitions. You may skip those sections if you think they are too “formal”.

Many people have contributed to this note. Special thanks to my collaborators. Fung San Gaan wrote the entire section 8 and part of section 7. ? Tingyu Wu collaborated with me on section 4. I shall also thank Bryce for his suggestions on the contents.

## 1 Basic Logic

Logic is the formal framework and rules of inference that ensure the validity and coherence of arguments in math.

**Remark.** We shall accept that sentences can be either true or false. Moreover, we assume that every English sentence can be stated in symbolic logic form.

A *proposition* is a sentence that is either true or false in a mathematical system. The label “true” or “false” assigned to a proposition is called its *truth value*. We use the letters  $T$  and  $F$  to represent “true” and “false”, respectively. An *axiom* is a proposition that is assumed to be true within a mathematical system without requiring proof. Axioms serve as the foundational building blocks of a mathematical theory, from which other propositions can be derived. A *theorem* is a proposition that has been proven to be true using logical reasoning and the accepted axioms and

previously established theorems of the mathematical system. The proof demonstrates why the theorem must hold based on these foundations.

Consider the proposition “ $\pi$  is not a rational number”, which is trivially true. However, we could always find some false companion of this proposition, such as “ $\pi$  is a rational number”. Similarly, we can find a true companion of a false proposition. Let  $P$  be a proposition, such a companion of  $P$  is called the *negation* of  $P$ , denoted  $\neg P$ .

Let  $P$  and  $Q$  be propositions. Those sentences can be combined using the word “and”, denoted  $P \wedge Q$ , and called the *conjunction* of  $P$  and  $Q$ . The proposition  $P \wedge Q$  is true if both  $P$  and  $Q$  are true. We can combine the propositions by the word “or”, denoted  $P \vee Q$ , and called the *disjunction* of  $P$  and  $Q$ . The proposition  $P \vee Q$  is true if at least one of  $P$  or  $Q$  is true. A *truth table* is shown below.

$P$	$Q$	$\neg P$	$P \wedge Q$	$P \vee Q$
$T$	$T$	$F$	$T$	$T$
$T$	$F$	$F$	$F$	$T$
$F$	$T$	$T$	$F$	$T$
$F$	$F$	$T$	$F$	$F$

Two propositions  $P$  and  $Q$  are *logically equivalent* if they have the same truth value in every possible combination of truth values for the variables in the statements, denoted  $P \equiv Q$ .

**Example.** Let  $P$  be a proposition, then  $P \equiv \neg(\neg P)$  is logically equivalent. To prove this statement, consider  $\neg P$  as a proposition  $Q$ , then we obtain the following truth table.

$P$	$Q \equiv \neg P$	$\neg Q \equiv \neg(\neg P)$
$T$	$F$	$T$
$F$	$T$	$F$

Here  $P$  and  $\neg Q$  has the same truth value in each case, so  $P \equiv \neg(\neg P)$ .

**Problem 1.1.** Let  $P$ ,  $Q$ , and  $R$  be propositions. Consider the following statements:

1.  $P \vee (Q \vee R) \equiv (P \vee Q) \vee R$ ;
2.  $P \wedge (Q \wedge R) \equiv (P \wedge Q) \wedge R$ .

Try to prove or disprove the statements.

**Problem 1.2.** Let  $P$ ,  $Q$ , and  $R$  be propositions. Consider the following statements:

1.  $\neg(P \vee Q) \equiv (\neg P) \wedge (\neg Q)$ ;
2.  $\neg(P \wedge Q) \equiv (\neg P) \vee (\neg Q)$ ;
3.  $P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)$ .

Try to prove or disprove the statements. Based on your results, can you find more properties?

Let  $P$  and  $Q$  be propositions. Consider the proposition “if  $n$  is a natural number, then  $2n$  is an even number”. Let  $P$  denotes “ $n$  is a natural number” and let  $Q$  denotes “ $2n$  is an even number”, then the sentence becomes “if  $P$ , then  $Q$ ”, denoted  $P \implies Q$ . This implication called a *conditional proposition*,  $P$  is called the *antecedent* and  $Q$  is called the *consequent*. The proposition  $P \implies Q$  is true if  $P$  is true and  $Q$  is true. What if  $P$  is false? The answer arises from one’s intuition.

Imagine your high school teacher say “if you didn’t submit your homework, then you haven’t completed it”. How would you argue against this sentence? The most likely response would be, “I did the homework but I didn’t submit it”. Whether or not you submitted your homework does not affect the truth value of the implication.

You should be convinced by your own intuition. This case is called a *vacuous truth*. In the proposition  $P \implies Q$ , when  $P$  is false,  $P \implies Q$  is true. The truth table of  $P \implies Q$  is shown below.

$P$	$Q$	$P \implies Q$
$T$	$T$	$T$
$T$	$F$	$F$
$F$	$T$	$T$
$F$	$F$	$T$

Let  $P$  and  $Q$  be propositions,  $(P \implies Q) \wedge (Q \implies P)$  is called a *biconditional proposition*, denoted  $P \iff Q$ . We will write this by “ $P$  is true if and only if  $Q$  is true”.

**Problem 1.3.** Let  $P$  and  $Q$  be propositions, show  $(\neg P \equiv \neg Q) \iff (P \equiv Q)$ .

**Example.** Let  $P$  and  $Q$  be propositions. Consider the conditional proposition  $P \implies Q$ . It is false only if  $P$  is true and  $Q$  is false, that is,  $\neg(P \implies Q) \equiv P \wedge (\neg Q)$ . Now we take the negation of the right side,  $\neg(P \wedge (\neg Q)) \equiv (\neg P) \vee (\neg(\neg Q)) \equiv (\neg P) \vee Q$ .

**Problem 1.4.** Write down the truth table of a biconditional proposition. Based on your truth table and the previous example, try to find a proposition  $R$  by “ $\vee$ ”, “ $\wedge$ ”, and “ $\neg$ ” such that  $R \equiv (P \iff Q)$ . If  $P \iff Q$  is true, does  $P \equiv Q$ ?

**Problem 1.5.** Let  $P$ ,  $Q$ ,  $R$ , and  $S$  be propositions. Rewrite  $P \implies (Q \implies (R \implies S))$  by “ $\vee$ ”, “ $\wedge$ ”, and “ $\neg$ ”. What is the negation of this sentence?

**Problem 1.6.** Let  $P$ ,  $Q$ , and  $R$  be propositions. Try to prove or disprove  $P \implies (Q \vee R) \equiv (\neg P) \vee Q \vee R$ . What about  $P \implies (Q \wedge R)$ ?

Given a proposition  $P \implies Q$ , the *converse* is defined as  $Q \implies P$  and the *contrapositive* is defined as  $(\neg Q) \implies (\neg P)$ . The truth table is shown below, and it suffices to conclude that  $(P \implies Q) \equiv (\neg Q \implies \neg P)$ .

$P$	$Q$	$P \implies Q$	$Q \implies P$	$\neg Q \implies \neg P$
$T$	$T$	$T$	$T$	$T$
$T$	$F$	$F$	$T$	$F$
$F$	$T$	$T$	$F$	$T$
$F$	$F$	$T$	$T$	$T$

**Problem 1.7.** Let  $P$  and  $Q$  be propositions, when does  $(P \implies Q) \equiv (Q \implies P)$ ?

Let  $P$  be the proposition “ $x$  is a natural number”. Here  $x$  is a *variable*, and the truth value of this proposition depends on  $x$ . For instance, if  $x = 1$ , then  $P$  is true; if  $x = 0.86$ , then  $P$  is false. A *propositional function* is a family of propositions depending on one or more variables. The collection of permitted variables is the *domain*. Now we write  $P(x)$  instead of  $P$ , so  $P(1)$  is true and  $P(0.86)$  is false.

**Problem 1.8.** Let  $x$  be a variable and let  $x$  be a natural number. Give a proposition  $P(x)$  such that  $P(x)$  is true when  $x \leq 2024$  and false when  $x \geq 2025$ .

Propositional functions are often quantified. The *universal quantifier* is denoted by “ $\forall$ ”, and the proposition  $\forall x(P(x))$  is true if and only if  $P(x)$  is true for every  $x$  in its domain. The *existential quantifier* is denoted by “ $\exists$ ”, and the proposition  $\exists x(P(x))$  is true if and only if  $P(x)$  is true for at least one  $x$  in its domain. Consider the proposition  $\forall x(P(x))$ , this means all  $x$  make  $P(x)$  true, so there does not exist some  $x$  such that  $P(x)$  is false, which is  $\neg(\exists x(\neg P(x)))$ .

**Example.** Let  $P(x)$  be a proposition, then  $\neg(\forall x(P(x))) \iff \neg(\neg(\exists x(\neg P(x)))) \iff \exists x(\neg P(x))$ .

**Problem 1.9.** Let  $P(x)$  be a proposition, show that  $\neg(\exists x(P(x))) \iff \forall x(\neg P(x))$ .

The order of quantifiers does matter the meaning of a proposition. Consider the proposition “for all natural number  $x$ , there exists a natural number  $y$  such that  $y > x$ ”. Pick some  $x$ , let  $y = x + 1$ , then  $y > x$  and  $y$  is a natural number, so the proposition is true. However, switching the order of quantifiers gives “there exists a natural number  $y$ , for all natural number  $x$ ,  $y > x$ ”. Suppose there exists such  $y$ , then  $y + 1$  is a natural number, so let  $x = y + 1$ , it is trivial that  $y < x$ , hence the proposition is false.

**Example.** Let  $P(x)$  and  $Q(y)$  be propositions. Consider the proposition  $\forall x(\exists y(P(x) \vee Q(y)))$ . To find its negation, let  $R(x) \equiv \exists y(P(x) \vee Q(y))$ , now the negation becomes  $\exists x(\neg R(x))$ . Since  $P$  only depends on  $x$ , let  $S(y) \equiv (P(x) \vee Q(y))$ , then we have  $\exists x(\neg(\exists y(S(y)))) \equiv \exists x(\forall y(\neg S(y))) \equiv \exists x(\forall y(\neg(P(x) \vee Q(y)))) \equiv \exists x(\forall y((\neg P(x)) \wedge (\neg Q(y))))$ .

**Problem 1.10.** Let  $P(x, y, z)$  be a proposition, consider the following propositions.

1.  $Q(x, y, z) \equiv \exists x(\forall y(\forall z(P(x, y, z))))$
2.  $R(x, y, z) \equiv \forall x(\exists y(\forall z(P(x, y, z))))$
3.  $S(x, y, z) \equiv \forall x(\forall y(\exists z(P(x, y, z))))$

What are the negations of those propositions? What is the negation of  $Q \vee (R \wedge S)$ ?

**Example.** Let  $P(x)$  and  $Q(x)$  be propositions. Consider the negation of  $P(x) \implies Q(x)$ ,  $\neg(P(x) \implies Q(x)) \equiv \neg((\neg P(x)) \vee Q(x)) \equiv P(x) \wedge (\neg Q(x)) \equiv \forall x(P(x) \wedge (\exists x(\neg Q(x)))) \equiv \exists x(P(x) \wedge (\neg Q(x)))$ . Notice that taking the negation brings an existential quantifier.

In the following sections, we shall assume readers are familiar with basic logic and use it as a tool to understand or prove propositions. Several expressions and their “translations” are shown below.

$P \implies Q$	$P \iff Q$
$P$ implies $Q$ ; if $P$ , then $Q$	$P$ if and only if $Q$
$P$ is sufficient for $Q$ ; $Q$ is necessary for $P$	$P$ is necessary and sufficient for $Q$

**Problem 1.11.** Given the following propositions, analyze their structures.

1. The number  $\sqrt{2}$  is not a rational number.
2. If  $x$  is a natural number, then  $x$  is an integer.
3. For all natural number  $x$ , for all rational number  $y$  with  $x < y < x + 1$ , there exists a real number  $z$  such that  $y < z < y + 1$  and  $z$  is irrational.
4. Given a sequence  $(x_n)$  of real numbers, we say  $(x_n)$  converges to a real number  $L$  if, for all real number  $\epsilon > 0$ , there exists a real number  $N$  such that, for all natural number  $n$ ,  $n > N$  implies  $|x_n - L| < \epsilon$ .

Find the negation of each proposition.

## 2 Some Axioms of Sets

In this section, we begin investigating sets, the most basic entities in mathematics. It is natural to ask: What is a set? There is no precise definition of sets. Intuitively, a *set* is a collection of objects that satisfy some property, and the objects are called *elements*.

**Remark.** This note is based on the ZFC set theory. In this system, every object is a set, so there are sets of sets. From now on, assume that there exists a set.

If  $S$  is a set and  $x$  is an element in  $S$ , then we say  $x$  belongs to  $S$ , denoted  $x \in S$ . If  $x$  does not belong to  $S$ , then we write  $x \notin S$ . If  $S$  has no element, then we call it an *empty set*, denoted  $\emptyset$ .

**Axiom of empty set.** There exists an empty set.

**Axiom of extensionality.** Two sets  $A$  and  $B$  are equal if and only if they have the same elements.

**Axiom schema of separation.** If  $P$  is a property, then for any set  $X$  there exists a set  $Y = \{x \in X \mid P(x)\}$ .

Elements determine a set. One way to describe a set is to explicitly list the elements. For example, we can write a set  $S = \{6, 7, 8\}$ . Another way is to express the elements by the properties they satisfy.

**Example.** Here are several examples of sets.

1. the set  $S = \{\emptyset, \{\emptyset\}, \{\{\emptyset\}\}$  has three elements;
2. the set  $\{2n \mid n \in \mathbb{N}\}$  is the set of all even numbers, where  $\mathbb{N}$  is the set of natural numbers;
3. the set  $\mathbb{Q} = \{a/b \mid a, b \in \mathbb{Z}, b \neq 0\}$  is the set of rational numbers, where  $\mathbb{Z}$  is the set of integers.

We shall provide constructions for  $\mathbb{N}$  and  $\mathbb{Z}$  later.

**Problem 2.1.** Write out the set of all positive integers and the set of all prime numbers.

**Definition 2.1.** Let  $S$  be a set. A set  $R$  is a *subset* of  $S$ , denoted  $R \subset S$ , if for all  $x \in R$ ,  $x \in S$ . If there exists some  $x \in S$  such that  $x \notin R$ , then  $R$  is called a *proper subset* of  $S$ , denoted  $R \subsetneq S$ .

It suffices to check that axiom schema of separation guarantees that subsets are sets.

**Remark.** Some textbooks use “ $\subseteq$ ” for subsets and “ $\subset$ ” for proper subsets.

**Remark.** From now on, try to verify whether those constructions are actually sets.

**Proposition.** Let  $A$  be a set, then  $A \subset A$ .

*Proof.* For all  $x \in A$ ,  $x \in A$ , so  $A \subset A$ . □

**Proposition.** Let  $X$  and  $Y$  be sets, then  $X = Y$  if and only  $X \subset Y$  and  $Y \subset X$ .

**Remark.** For a biconditional proposition  $P \iff Q$ , we use the notation “ $(\Rightarrow)$ ” in the proof to show  $P \implies Q$  and “ $(\Leftarrow)$ ” for  $Q \implies P$ .

*Proof.* Let  $X$  and  $Y$  be sets.  $(\Rightarrow)$  For all  $x \in X$ , since  $X = Y$ ,  $x \in Y$ , so  $X \subset Y$ . For all  $y \in Y$ , since  $X = Y$ ,  $y \in X$ , so  $Y \subset X$ .  $(\Leftarrow)$  Suppose  $X \neq Y$ . If  $X \subset Y$ , then there exists  $a \in Y$  such that  $a \notin X$ , so  $X \not\subset Y$ , a contradiction. □

**Proposition.** Let  $A$  be any set, then  $\emptyset \subset A$ .

*Proof.* Suppose  $\emptyset \not\subset A$ , then there exists  $x \in \emptyset$  such that  $x \notin A$ , since  $x \in \emptyset$  is false, a contradiction. □

**Problem 2.2.** Prove that a set is independent of the order of its elements. For example,  $\{1, 2, 3\} = \{3, 2, 1\}$ .

**Problem 2.3.** If  $X$ ,  $Y$ , and  $Z$  are sets such that  $X \subset Y$  and  $Y \subset Z$ , prove that  $X \subset Z$ .

**Axiom of pairing.** For two objects  $a$  and  $b$ , there exists a set  $\{a, b\}$  containing exactly  $a$  and  $b$ .

**Definition 2.2.** Let  $a$  and  $b$  be some objects. An *ordered pair*  $(a, b)$  is defined as the set  $\{\{a\}, \{a, b\}\}$ .

**Problem 2.4.** Show that an ordered pair is indeed a set.

**Proposition.** Let  $(a, b)$  and  $(c, d)$  be ordered pairs, then  $(a, b) = (c, d)$  if and only if  $a = c$  and  $b = d$ .

*Proof.* We have  $(a, b) = \{\{a\}, \{a, b\}\}$  and  $(c, d) = \{\{c\}, \{c, d\}\}$ .  $(\Rightarrow)$  Suppose  $a \neq c$ , then  $\{a\} \neq \{c\}$ . If  $\{a\} = \{c, d\}$ , then  $c = d = a$ , a contradiction. Suppose  $b \neq d$ . If  $a = c$ , then  $\{a\} = \{c\}$  and  $\{a, b\} \neq \{c, d\}$ , a contradiction.  $(\Leftarrow)$  If  $a = c$  and  $b = d$ , then  $\{a, b\} = \{c, d\}$  and  $\{a\} = \{c\}$ , hence  $(a, b) = (c, d)$ . □

The definition of ordered pairs can be extended to multiple elements. We call  $(a_1, \dots, a_n)$  a  *$n$ -tuple*.

**Problem 2.5.** Prove that  $\{a\} = \{a, a\}$ . A set with one element is called a *singleton*.

**Problem 2.6.** Write out the definition to  $n$ -tuples, where  $n$  is a positive integer.

**Axiom of union.** For all set  $X$ , there exists a set  $Y = \bigcup X$ , the union of all elements of  $X$ .

**Definition 2.3.** Let  $A$  and  $B$  be sets. The *union* of  $A$  and  $B$  is the set  $\{x \mid x \in A \text{ or } x \in B\}$ , denoted  $A \cup B$ . The *intersection* of  $A$  and  $B$  is the set  $\{x \mid x \in A, x \in B\}$ . We say  $A$  and  $B$  are *disjoint* if  $A \cap B = \emptyset$ . The *complement* of  $A$  in  $B$  is the set  $\{x \mid x \in B, x \notin A\}$ , denoted  $B \setminus A$ .

**Problem 2.7.** Let  $A$  and  $B$  be sets. Prove that  $A \cup B$ ,  $A \cap B$ , and  $A \setminus B$  are sets based on the axioms.

**Proposition.** Let  $A$  and  $B$  be sets, then  $A \cup B = B \cup A$ .

*Proof.* For all  $x \in A \cup B$ , if  $x \in A$ , then  $x \in B \cup A$ . If  $x \in B$ , then  $x \in B$ , hence  $A \cup B = B \cup A$ .  $\square$

**Problem 2.8.** Let  $A$ ,  $B$ , and  $C$  be sets. Prove the following propositions.

1.  $A \cap B = B \cap A$ .
2.  $A \cup (B \cap C) = (A \cup B) \cap C$ .
3.  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .
4.  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .
5.  $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$ .

**Theorem 2.1** (De Morgan's law). Let  $A$ ,  $B$ , and  $C$  be sets, then  $C \setminus (A \cap B) = (C \setminus A) \cup (C \setminus B)$  and  $C \setminus (A \cup B) = (C \setminus A) \cap (C \setminus B)$ .

*Proof.* Let  $x \in C \setminus (A \cap B)$ , then  $x \in C$  and  $x \notin A \cap B$ , that is,  $x \notin A$  and  $x \notin B$ . If  $x \notin C \setminus A$ , then  $x \notin A$ , so  $x \notin B$  and  $x \in C \setminus B$ . Hence  $C \setminus (A \cap B) \subset (C \setminus A) \cup (C \setminus B)$ . Now let  $x \in (C \setminus A) \cup (C \setminus B)$ , then  $x \in C$  and  $x \notin A$  or  $x \notin B$ , so  $x \notin A \cap B$ , that is,  $x \in C \setminus (A \cap B)$ , hence  $(C \setminus A) \cup (C \setminus B) \subset C \setminus (A \cap B)$ . The proof of the second part is left as an exercise.  $\square$

Some texts assume the existence of an “universal set”, denoted  $U$ , which has all objects as elements including itself, so we can define complements of any set  $S$  as the set  $U \setminus S$ . However, this assumption leads to a paradox. Consider the set  $S$ , defined as the set of all sets that are not members of themselves, that is,  $S = \{X \mid X \notin X\}$ . Does  $S$  belong to  $S$ ? This is known as Russell's Paradox. Assume  $S \in S$ , then by the definition of  $S$ ,  $S \in S$  implies  $S \notin S$ , a contradiction. Assume  $S \notin S$ , then  $S \in S$ . This is also a contradiction.

**Axiom of regularity.** For all nonempty sets, there is an element of the set that shares no element with the set.

**Proposition.** A set is not an element of itself.

*Proof.* Suppose  $X$  is a set such that  $X \in X$  and  $X \in \{X\}$ . By contradiction,  $X \neq \emptyset$ . We have  $X \cap \{X\} = \{x \mid x \in X, x \in \{X\}\} = X$ . By the axiom of regularity,  $X \cap \{X\} = \emptyset$ , then  $X = \emptyset$ , a contradiction.  $\square$

**Definition 2.4.** Let  $X$  be a set. A *partition* of  $X$  is a set  $\{X_i\}$  of non-empty subsets of  $X$  such that every elements  $x \in X$  lies in exactly one of there subsets.

**Problem 2.9.** Prove that a partition is a set.

**Definition 2.5.** Let  $X$  be a set, and let the *successor* of  $X$  be  $X^+ = X \cup \{X\}$ . A set  $S$  is called an *inductive set* if  $\emptyset \in S$  and for all  $X \in S$ ,  $X^+ \in S$ .

**Axiom of infinity.** There exists an inductive set.

**Proposition.** The intersection of two inductive sets is an inductive set.

*Proof.* Let  $A$  and  $B$  be inductive sets, then  $\emptyset \in A \cap B$ . For all  $S \in A \cap B$ ,  $S \in A$  and  $S \in B$ . Since  $A$  and  $B$  are inductive,  $S^+ \in A$  and  $S^+ \in B$ , hence  $A \cap B$  is inductive.  $\square$

**Definition 2.6.** The set of all *natural numbers*, denoted  $\mathbb{N}$ , is the intersection of all inductive sets.

We denote  $0 = \emptyset$ ,  $1 = 0^+$ ,  $2 = 1^+$ ,  $\dots$

**Problem 2.10.** Prove that the set of all natural numbers is a subset of any inductive set.

**Axiom of power set.** For any  $X$  there exists a set consisting of all subsets of  $X$ .

**Definition 2.7.** Given a set  $X$ , the set of all subsets of  $X$  is called its *power set*, denoted  $\mathcal{P}(X)$ .

**Example.** Let  $X = \{a, b\}$ , the power set  $\mathcal{P}(X) = \{\emptyset, \{a\}, \{b\}, \{a, b\}\}$ .

**Definition 2.8.** Let  $X$  and  $Y$  be sets. The *Cartesian product*  $X \times Y$  is the set of all ordered pairs  $(a, b)$ , where  $a \in X$  and  $b \in Y$ .

**Remark.** Let  $S$  be a set. We denote  $S \times \dots \times S$ , the Cartesian product of  $n$  times of  $S$ , by  $S^n$ .

**Problem 2.11.** Let  $X$  and  $Y$  be sets. Write out the set  $\mathcal{P}(\mathcal{P}(X \cup Y))$ . Prove that  $X \times Y = \{z \in \mathcal{P}(\mathcal{P}(X \cup Y)) \mid \text{there exists } x \in X, y \in Y \text{ such that } z = (x, y)\} \subset \mathcal{P}(\mathcal{P}(X \cup Y))$ , hence  $X \times Y$  is a set.

**Problem 2.12.** Let  $S$  be a set, prove that  $S \subsetneq \mathcal{P}(S)$ .

**Problem 2.13.** Let  $A$ ,  $B$ , and  $C$  be sets. Prove the following propositions.

1.  $A \times B = B \times A$  if and only if  $A = B$ .
2.  $A \times (B \times C) = (A \times B) \times C$ .
3.  $(A \cup B) \times C = (A \times C) \cup (B \times C)$ .
4.  $(A \cap B) \times C = (A \times C) \cap (B \times C)$ .
5.  $(A \setminus B) \times C = (A \times C) \setminus (B \times C)$ .

**Definition 2.9.** The *disjoint union* of two sets  $A$  and  $B$ , denoted  $A \amalg B$ , is the set  $A \amalg B = (A \times \{0\}) \cup (B \times \{1\})$ .

**Problem 2.14.** Let  $A$  and  $B$  be sets, prove that  $A \amalg B$  is a set.

**Problem 2.15.** Let  $X$  and  $Y$  be sets. The *symmetric difference*  $X \triangle Y$  is defined to be  $(X \setminus Y) \cup (Y \setminus X)$ . Prove that  $X \triangle Y$  is a set.

**Definition 2.10.** A *binary operation*  $R$  is a set of ordered pairs. If  $(x, y) \in R$ , we write  $xRy$ . The *domain* of  $R$  is the set  $\text{dom}(R) = \{u \mid \text{there exists } v \text{ such that } (u, v) \in R\}$ . The *range* of  $R$  is the set  $\text{ran}(R) = \{v \mid \text{there exists } u \text{ such that } (u, v) \in R\}$ .

It suffices to show a binary operation is indeed a set. Let  $R$  be a binary operation, then  $R \subset X \times Y$  for some sets  $X$  and  $Y$ . By the axiom schema of separation,  $R$  is a set.

**Problem 2.16.** Let  $R$  be a binary operation. Prove that  $\text{dom}(R), \text{ran}(R) \subset \bigcup(\bigcup R)$ , hence, by the axiom of union,  $\text{dom}(R)$  and  $\text{ran}(R)$  are sets.

**Definition 2.11.** Let  $R$  be a binary operation on a set  $S$ , that is,  $R \subset S \times S$ . We say  $R$  is an *equivalence relation* if the following properties hold.

1. For all  $a \in X$ ,  $aRa$ . (reflexive)
2. If  $aRb$ , then  $bRa$ . (symmetric)
3. If  $aRb$  and  $bRc$ , then  $aRc$ . (transitive)

For all  $a \in A$ , the set  $S_a = \{b \mid aRb\}$  is the *equivalence class* of  $a$ .

**Problem 2.17.** Prove that an equivalence class is a set.

**Problem 2.18.** Prove that “=” is an equivalence relation in  $\mathbb{N}$ .

**Problem 2.19.** Let  $R$  be a binary operation on a set  $X$ . For all  $a, b \in A$ , prove that  $S_a \cap S_b$  is either  $\emptyset$  or  $S_a$ . Prove that the collection of  $S_a$  forms a partition of  $X$ .

**Definition 2.12.** Let  $\leq$  be a binary relation on a set  $X$ . We say  $\leq$  is a *partial ordering* if the following conditions hold.

1. For all  $x \in X$ ,  $x \leq x$ .
2. For all  $x, y \in X$ ,  $x \leq y$  and  $y \leq x$  implies  $x = y$ .
3. For all  $a, b, c \in X$ , if  $a \leq b$  and  $b \leq c$ , then  $a \leq c$ .

The set with a partial ordering is called a *partially ordered set*.

Let  $(X, \leq)$  be a partially ordered set. Let  $Y \subset X$  and  $Y \neq \emptyset$ . We say  $a \in Y$  is the greatest element of  $Y$  if for all  $y \in Y$ ,  $a \geq y$ . Similarly, we could define the smallest element of a nonempty subset of  $X$ .

**Definition 2.13.** Let  $(X, \leq)$  be a partially ordered set and let  $Y$  be a nonempty subset of  $X$ . An element  $a \in X$  is said to be an *upper bound* of  $Y$  if for all  $y \in Y$ ,  $y \leq a$ . An element  $a \in X$  is said to be a *lower bound* of  $Y$  if for all  $y \in Y$ ,  $a \leq y$ . We call the least upper bound of  $Y$  the *supremum* and we call the greatest lower bound of  $Y$  the *infimum*.

**Definition 2.14.** A partially ordered set  $(X, \leq)$  is *linearly ordered* if for all  $p, q \in X$ , either  $p \leq q$  or  $q \leq p$ .

**Example.** The set of natural numbers  $\mathbb{N}$  forms a linearly ordered set in set inclusions.

**Proposition.** Let  $(X, \leq)$  be a partially ordered set and let  $Y \subset X$ , then  $Y$  is partially ordered.

*Proof.* For all elements  $a, b, c \in Y$ ,  $a, b, c \in X$ , so  $Y$  inherits the partial ordering of  $X$ . □

**Problem 2.20.** Let  $(X, \leq)$  be a linearly ordered set and let  $Y \subset X$ , prove that  $Y$  is linearly ordered.

**Problem 2.21.** Let  $X$  be a set. If  $(\mathcal{P}(X), \subset)$  is a linearly ordered set, prove that  $X$  is either a singleton or the empty set.

**Definition 2.15.** Let  $(X, \leq)$  be a partially ordered set. The set is *well-ordered* if for every nonempty subset  $S$  of  $X$ , there exists  $a \in S$  such that for all  $s \in S$ ,  $a \leq s$ .

**Problem 2.22.** Prove that every well-ordered set has a smallest element.

Let  $a, b \in X$ , where  $X$  is well-ordered. If  $a \geq b$  and  $a \neq b$ , then we write the strict ordering as  $a > b$ .

**Definition 2.16.** Let  $(X, \leq)$  be a well-ordered set. Let  $x \in X$ . Then the set  $\{a \mid a \leq x\}$  is called an *initial segment* of  $X$  given by  $x$ .

It is trivial that any initial segment of a well-ordered set is well-ordered.

**Theorem 2.2** (well-ordering theorem). Every set is well-orderable.

The well-ordering theorem is equivalent to the axiom of choice, which will be discussed later in the next section. You may assume it is correct for now.

**Theorem 2.3** (finite induction). Given a subset  $S \subset \mathbb{N}$  of the natural numbers with  $0 \in S$  and  $n \in S$  implies  $n + 1 \in S$ , then  $S = \mathbb{N}$ .

*Proof.* Suppose  $S \neq \mathbb{N}$ , then  $X = \mathbb{N} \setminus S$  is a nonempty set. By the well-ordering principle,  $X$  has a smallest element. Since  $0 \in S$ ,  $0 \notin X$ , so the minimal element of  $X$  can be written in the form  $k + 1$ , where  $k \in \mathbb{N}$ . Recall that  $k + 1 = k^+ = k \cup \{k\}$  is the successor of  $k \in \mathbb{N}$ , since  $k + 1$  is the smallest element,  $k \notin X$ , so  $k \in S$ . Now we have  $k \in S$  and  $k + 1 \notin S$ , a contradiction. □

**Example.** Consider the statement: let  $n \in \mathbb{N}$ , show that  $\sum_{i=0}^n = (n(n+1))/2$ . If  $n = 0$ , then the equation trivially holds. Assume  $\sum_{i=0}^k = (k(k+1))/2$  holds for some  $k \in \mathbb{N}$ , then  $\sum_{i=0}^{k+1} = (k(k+1))/2 + (k+1) = ((k+1)(k+2))/2$ . Hence, by induction,  $\sum_{i=0}^n = (n(n+1))/2$  for all  $n \in \mathbb{N}$ .



**Problem 2.23.** Prove that given a subset  $S \subset \mathbb{N}$  of the natural numbers with  $0 \in S$  and  $\{0, 1, \dots, n\} \subset S$  implies  $n + 1 \in S$ , then  $S = \mathbb{N}$ . This is known as the *complete finite induction*. Prove that the complete finite induction implies the finite induction, conclude that they are equivalent.

### 3 Functions and Well-Ordering

**Definition 3.1.** Let  $X$  and  $Y$  be sets. A *function*  $f$  is a binary operation  $f \subset X \times Y$  such that for all  $x \in X$ , there exists a unique  $y \in Y$  such that  $(x, y) \in f$ . We say  $f$  is a function from  $X$  to  $Y$ , denoted  $f : X \rightarrow Y$ . The set  $Y$  is called the *codomain* of  $f$ , denoted  $\text{cod}(f)$ .

**Problem 3.1.** Given two sets  $A$  and  $B$ , prove that the collection of functions from  $A$  to  $B$  is a subset of  $\mathcal{P}(A \times B)$ , hence it is a set.

**Definition 3.2.** Let  $f : X \rightarrow Y$  be a function, the *image* of  $X$  under  $f$ , denoted  $\text{im}(f)$ , is the range of  $f$ . For all  $(x, y) \in f$ , we write  $f(x) = y$ . The *preimage* of  $Y$  under  $f$ , denoted  $f^{-1}(Y)$ , is the set  $\{x \mid x \in X \text{ and } f(x) \in Y\}$ . Two functions  $f$  and  $g$  are the same if  $\text{dom}(f) = \text{dom}(g)$ ,  $\text{cod}(f) = \text{cod}(g)$ , and for all  $x \in \text{dom}(f)$ ,  $f(x) = g(x)$ .

**Example.** The operation  $f : \mathbb{N} \rightarrow \mathbb{N}$  defined by  $f(n) = n^+$ , where  $n \in \mathbb{N}$ , is a function.

**Example.** The operation  $f : \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$  defined by  $f(x) = (x, x)$ , where  $x \in \mathbb{R}$ , is a function.

**Problem 3.2.** Let  $f$  be a function, prove that  $\text{ran}(f) \subset \text{cod}(f)$ .

**Problem 3.3.** Consider a function  $f : \emptyset \rightarrow A$ , prove that  $f = \emptyset$ . Prove that such  $f$  is unique for every set  $A$ .

**Problem 3.4.** Verify whether the following binary operations are functions.

1.  $f : \mathbb{R} \rightarrow \mathbb{R}$  with  $f(x) = \sqrt{x}$  for all  $x \in \mathbb{R}$ .
2.  $f : \{1, 2, 3\} \rightarrow \{2, 3\}$  with  $f = \{\{1, 2\}, \{2, 2\}, \{3, 2\}\}$ .
3.  $f : \mathbb{N} \rightarrow \mathbb{Q}$  with  $f(x) = x^4 - x^2$  for all  $x \in \mathbb{N}$ .
4.  $f : \mathbb{Q} \rightarrow \mathbb{Z}$  with  $f(x) = |x|$  for all  $x \in \mathbb{Q}$ .

**Definition 3.3.** Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  be functions. The *composition* of  $f$  and  $g$ , denoted  $g \circ f$ , is defined as  $g \circ f : X \rightarrow Z$  with  $(g \circ f)(x) = g(f(x))$  for all  $x \in X$ .

**Problem 3.5.** Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  be functions. Prove that  $g \circ f$  is a well-defined function.

**Proposition.** The composition of functions is associative, that is, for all  $f : X \rightarrow Y$ ,  $g : Y \rightarrow Z$ , and  $h : Z \rightarrow S$ ,  $h \circ (g \circ f) = (h \circ g) \circ f$ .

*Proof.* For all  $x \in X$ ,  $(h \circ g \circ f)(x) = h(g(f(x))) = (h \circ g)(f(x)) = ((h \circ g) \circ f)(x)$ . □

Consider a diagram, where every vertex is an object and every arrow preserves the structure of those objects. Such a diagram is said to be *commutative* if all paths between two vertices are equivalent. In this section, every vertex is a set and every arrow is a function.

**Example.** Consider the following diagrams.



The left diagram is commutative if  $g \circ f = h$ . The right diagram is commutative if  $g \circ f = h$  and  $\psi \circ \varphi = h$ .

**Problem 3.6.** Let  $f : S_1 \times S_2 \rightarrow S_3$  be a function. We say  $f$  is commutative as a composition if  $f(a, b) = f(b, a)$ . Similarly,  $f$  is associative as a composition if  $f(f(a, b), c) = f(a, f(b, c))$ . Prove that if  $f$  is commutative, then  $S_1 = S_2$ . Prove that if  $f$  is associative, then the following diagram commutes.

$$\begin{array}{ccc} S_1 \times S_2 \times S_3 & \xrightarrow{f} & S_4 \times S_3 \\ \downarrow g: S_2 \times S_3 \rightarrow S_5 & & \downarrow \\ S_5 & \longrightarrow & S_6 \end{array}$$

**Definition 3.4.** A function  $f : X \rightarrow Y$ , where  $X, Y \subset \mathbb{R}$ , is said to be an *odd function* if for all  $x \in X$ ,  $f(x) = -f(-x)$ . The function  $f$  is said to be an *even function* if for all  $x \in X$ ,  $f(x) = f(-x)$ .

**Example.** Here are some examples of odd and even functions.

1. The function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f(x) = x^2$  for all  $x \in \mathbb{R}$  is even.
2. The function  $g : \mathbb{R} \rightarrow \mathbb{R}$  defined by  $g(x) = x$  for all  $x \in \mathbb{R}$  is odd.
3. The function  $0 : \mathbb{R} \rightarrow \mathbb{R}$  defined by  $0(x) = 0$  for all  $x \in \mathbb{R}$  is both odd and even.

**Proposition.** Any function  $f : X \rightarrow Y$ , where  $X, Y \subset \mathbb{R}$ , can be written as  $f = \varphi + \psi$ , where  $\varphi$  is an odd function and  $\psi$  is an even function.

*Proof.* For all  $x \in X$ , define  $\varphi = (f(x) - f(-x))/2$  and  $\psi = (f(x) + f(-x))/2$ , then  $\varphi(-x) = (f(-x) - f(x))/2 = -\varphi(x)$  and  $\psi(-x) = (f(-x) + f(x))/2 = \psi(x)$ . Moreover,  $\varphi(x) + \psi(x) = (f(x) - f(-x) + f(x) + f(-x))/2 = f(x)$ .  $\square$

**Problem 3.7.** Given an example of a function that is neither odd nor even. Decompose it as a sum of an odd function and an even function.

**Problem 3.8.** Prove that such decomposition for each  $f : X \rightarrow Y$ , where  $X, Y \subset \mathbb{R}$ , is unique.

**Definition 3.5.** Let  $f : A \rightarrow B$  be a function. We say  $f$  is *injective* if for all  $x, y \in B$  and  $x = y$ , then  $f^{-1}(x) = f^{-1}(y)$ . We say  $f$  is *surjective* if  $\text{ran}(f) = \text{cod}(f)$ . The function  $f$  is said to be *bijective* if it is both injective and surjective.

**Problem 3.9.** State examples if such functions exist.

1.  $f : \{1, 2, 3\} \rightarrow \{4, 5, 6\}$  that is injective but not surjective.
2.  $f : \{1, 2, 3\} \rightarrow \{4, 5, 6\}$  that is surjective but not injective.
3.  $f : \mathbb{R} \rightarrow \mathbb{R}$  that is injective but not surjective.
4.  $f : \mathbb{R} \rightarrow \mathbb{R}$  that is surjective but not injective.
5.  $f : \mathbb{R} \rightarrow \{1, 2, 3\}$  that is injective but not surjective.
6.  $f : \mathbb{R} \rightarrow \{1, 2, 3\}$  that is surjective but not injective.

**Proposition.** The composition of two surjective functions is surjective.

*Proof.* Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be surjective functions, then  $\text{ran}(f) = \text{cod}(f) = B$  and  $\text{ran}(g) = \text{cod}(g) = C$ . For all  $x \in C$ ,  $g^{-1}(x) \in B$  and  $f^{-1}(g^{-1}(x)) \in A$ , so  $C \subset \text{ran}(g \circ f)$ , hence  $g \circ f$  is surjective.  $\square$

**Problem 3.10.** Let  $f : X \rightarrow Y$  be a surjective function. Prove that there exists an injective function  $g : Y \rightarrow X$ .

**Problem 3.11.** Let  $f$  and  $g$  be functions. Prove or disprove the following statements.

1. If  $f$  and  $g$  are injective, then  $f \circ g$  is injective.
2. If  $f$  and  $g$  are bijective, then  $f \circ g$  is bijective.
3. If  $f$  is surjective and  $g$  is injective, then  $f \circ g$  is injective.

**Definition 3.6.** A set  $S$  is said to be *finite* if there exists a bijective function  $f : S \rightarrow n$ , where  $n \in \mathbb{N}$ . The *cardinality* of  $S$ , denoted  $|S|$ , is the number  $n$ . If  $S$  is not finite, we say  $S$  is *infinite*.

**Problem 3.12.** Let  $S$  and  $X$  be finite sets with  $|S| = |X|$ . Let  $f : S \rightarrow X$  be a function. Prove that if  $f$  is injective, then  $f$  is surjective. Does the converse hold?

**Problem 3.13.** Let  $S$  be a finite set. Prove that there does not exist a surjective function  $f : S \rightarrow \mathcal{P}(S)$ .

**Definition 3.7.** Let  $f : X \rightarrow Y$  be a function. Let  $Z \subset X$ , then the *restriction* of  $f$  onto  $Z$  is the map  $f|_Z : Z \rightarrow Y$  defined by  $f|_Z(z) = f(z)$  for all  $z \in Z$ .

**Definition 3.8.** Let  $A$  be a set. The *identity function*, denoted  $\text{id}_A$ , is the function  $\text{id}_A(x) = x$  for all  $x \in A$ .

**Problem 3.14.** Let  $f : A \rightarrow B$  be a function, prove that  $f \circ \text{id}_A = f = \text{id}_B \circ f$ .

**Definition 3.9.** Let  $f : A \rightarrow B$  be a function. The function  $g : B \rightarrow A$  is a *left inverse* of  $f$  if  $g \circ f = \text{id}_A$ . The function  $h : B \rightarrow A$  is a *right inverse* of  $f$  if  $f \circ h = \text{id}_B$ . A function  $\varphi$  is called an *inverse* of  $f$  if it is both a left inverse and a right inverse of  $f$ .

**Proposition.** Let  $f$  be a function. If  $g$  is an inverse of  $f$ , then  $g$  is unique.

*Proof.* Suppose  $g$  and  $h$  are inverses of  $f$ , then  $h = h \circ f \circ g = (h \circ f) \circ g = g$ . □

**Problem 3.15.** Let  $f$  be a function with a left inverse  $g$ . Prove that if  $f$  has a right inverse, then the right inverse is  $g$ , conclude that  $g$  is the inverse of  $f$ .

**Problem 3.16.** Prove that a function has a left inverse if and only if it is injective. Prove that a function has a right inverse if and only if it is surjective, conclude that a function is bijective if and only if it has an inverse.

**Definition 3.10.** Let  $\sim$  be an equivalence relation on a set  $X$ . The *quotient set*, denoted  $X_{\sim}$ , is the set  $\{[x] \mid x \in X\}$ .

**Example.** Let  $S = \{a, b, c, d\}$ . Let  $\sim$  be an equivalence relation on  $S$  such that  $a \sim b$  and  $c \sim d$ . The quotient set  $S_{\sim}$  is  $\{a, c\}$ .

**Problem 3.17.** Consider the set of all integers  $\mathbb{Z}$ . Fix  $n \in \mathbb{Z}$ , Let  $\sim_n$  be an equivalence relation on  $\mathbb{Z}$  such that  $a \sim_n b$  if and only if  $a - b = kn$  for some  $k \in \mathbb{Z}$ . Prove that  $\sim_n$  is indeed an equivalence relation. Find the quotient set  $\mathbb{Z}_{\sim_2}$ . Find the quotient set  $\mathbb{Z}_{\sim_n}$  for all  $n$ .

**Definition 3.11.** Let  $X$  and  $Y$  be sets. Let  $\sim$  be an equivalence relation on  $X$ . A function  $f : X \rightarrow Y$  is *invariant* under the equivalence relation such that,  $x \sim y$  if and only if  $f(x) = f(y)$ .

**Problem 3.18.** Let  $X$  be a set and let  $\sim$  be an equivalence relation on  $X$ . Prove that  $\pi : X \rightarrow X_{\sim}$  defined by  $x \mapsto [x]$  is a well-defined surjective function. Prove it is invariant under  $\sim$ .

**Proposition.** Let  $f : X \rightarrow Y$  be a function, then  $f$  is an invariant function under some equivalence relation on  $X$ .

*Proof.* Define  $\sim$  on  $X$  by  $x \sim y$ , where  $x, y \in X$ , if and only if  $f(x) \sim f(y)$ . For all  $x = y$ ,  $f(x) = f(y)$ . For all  $x = y = z$ ,  $f(x) = f(y) = f(z)$ . Hence  $\sim$  is a well-defined equivalence relation on  $X$ , and  $f$  is trivially invariant under  $\sim$ . □

Recall that we have defined partial ordering by “ $\leq$ ”. From now on, we require that every partial-ordering to be a strict ordering.

**Definition 3.12.** Let  $(P, <)$  and  $(Q, <)$  be partially ordered set and let  $f : P \rightarrow Q$  be a function. We say  $f$  is *order-preserving* if  $x < y$  implies  $f(x) < f(y)$ . If  $f$  is also bijective, then we say  $P$  and  $Q$  have the same *order-type*, denoted  $P \approx Q$ .

**Problem 3.19.** Prove that if  $f : P \rightarrow Q$  is order-preserving and bijective, then  $f^{-1}$  is also order-preserving.

**Problem 3.20.** Let  $f : X \rightarrow Y$  be order-preserving. Prove that if  $X$  is well-ordered, then  $Y$  is well ordered.

**Proposition.** Let  $X$  be well-ordered and let  $f : X \rightarrow X$  be order-preserving. Then  $f(x) \geq x$  for all  $x \in X$ .

*Proof.* Suppose  $\{x \in X \mid f(x) < x\} \neq \emptyset$ , then it is well-ordered. Let  $z$  be the least element of this set and let  $f(z) = w$ , then  $f(w) < w$ , a contradiction.  $\square$

**Problem 3.21.** Prove that any well-ordered set cannot have the same order-type as any of its initial segment.

**Proposition.** If  $W_1$  and  $W_2$  are well-ordered, then one of the following conditions hold.

1.  $W_1$  and  $W_2$  have the same order-type.
2.  $W_1$  has the same order-type as an initial segment of  $W_2$ .
3.  $W_2$  has the same order-type as an initial segment of  $W_1$ .

*Proof.* Denote the initial segment of  $W_i$  by  $x$  by  $W_i(x)$ . Let  $f : W_1 \rightarrow W_2$  be a function defined by  $f(x) = y$  if  $W_1(x)$  has the same order-type as  $W_2(y)$ . It is trivial that this map is well-defined and order-preserving. Let  $y_1 = y_2$ , then there exist  $x_1, x_2 \in W_1$  such that  $W_1(x_1) \approx W_2(y_1) = W_2(y_2) \approx W_1(x_2)$ , so  $W_1(x_1) \approx W_1(x_2)$ , which implies  $x_1 = x_2$ . Hence  $f$  is injective. Let  $h : W_1(x) \rightarrow W_2(y)$  be an order-preserving bijection. If  $x' < x$ , since  $f$  is order-preserving,  $W_1(x') \approx W_2(h(x'))$ . If  $\text{dom}(f) = W_1$  and  $\text{ran}(f) = W_2$ , then it is exactly case (i). If  $\text{ran}(f) \neq W_2$ . Let  $y_0$  be the least element in  $W_2 \setminus \text{ran}(f)$ , then  $W_2(y_0) = \text{ran}(f)$ . Suppose  $\text{dom}(f) \neq W_1$ , then pick the least element of  $W_1 \setminus \text{dom}(f)$ , denoted  $x_0$ . It is trivial that  $f(x_0) = y_0$ , a contradiction. This is case (ii). Similarly, if  $\text{dom}(f) \neq W_1$ , define the same  $x_0$ . By contradiction, we have case (iii).  $\square$

**Definition 3.13.** If  $\varphi(x, p_1, p_2, \dots)$  is some formula, then we say  $C = \{x \mid \varphi\}$  is a *class*, that is,  $x$  is a member of  $C$  if and only if  $x$  satisfies  $\varphi$ .

**Remark.** Although a class might not be a set, we extend the notion “ $\in$ ” to classes. Let  $C$  is a class, we write  $x \in C$  if  $x$  satisfies the formula  $\varphi$  correspond to  $C$ .

**Example.** The collection of all sets is a class.

**Problem 3.22.** Prove that every set is a class.

**Axiom schema of replacement.** If a class  $F$  is a function, then there exists a set  $Y = F(X) = \{F(x) \mid x \in X\}$  for all  $X$ .

**Problem 3.23.** Prove that axiom schema of replacement implies axiom schema of separation.

Notice that the axiom schema of separation guarantees the existence of functions, so we need both axioms.

**Definition 3.14.** A set  $S$  is said to be *transitive* if for all  $s \in S$ ,  $s \subset S$ . A set is an *ordinal* if it is transitive and well-ordered by  $\in$ . The *successor ordinal* of an ordinal  $\alpha$  is defined to be  $\alpha \cup \{\alpha\}$ , and we denote it by  $\alpha + 1$ .

**Problem 3.24.** Prove that the element of an ordinal is an ordinal.

**Proposition.** The successor ordinal of an ordinal is an ordinal.

*Proof.* Let  $\alpha$  be an ordinal. For all  $s \in \alpha + 1$ , if  $s \in \alpha$ , since  $\alpha$  is an ordinal,  $s \subset \alpha \subset \alpha + 1$ ; if  $s \in \{\alpha\}$ , then  $s = \alpha \subset \alpha + 1$ . Since  $\alpha$  is well-ordered by  $\in$ ,  $\alpha$  is the  $\in$ -maximal element in  $\alpha + 1$ , hence  $\alpha + 1$  is also well-ordered.  $\square$

Let  $\alpha$  and  $\beta$  be ordinals. We define  $\alpha < \beta$  if and only if  $\alpha \in \beta$ .

**Example.** The set  $0 \in \mathbb{N}$  is an ordinal.

**Problem 3.25.** The intersection of ordinals is an ordinal.

**Definition 3.15.** An ordinal  $\lambda$  is said to be a *limit ordinal* if  $\lambda \neq \emptyset$  and for all  $\alpha < \lambda$ ,  $\alpha + 1 < \lambda$ .

**Problem 3.26.** Prove that every limit ordinal is an ordinal.

Suppose there exists a set of all ordinals, denoted  $\text{On}$ . Since every element of an ordinal is an ordinal,  $\text{On}$  is transitive. Let “ $<$ ” be the ordering on  $\text{On}$ , then it is trivially a partial ordering. Let  $S$  be a nonempty subset of  $\text{On}$ . Consider  $\bigcap s_i \subset s_i$ , where  $s_i \in S$ , if  $T \notin S$ , then  $T \subset s_i$ , so  $T \subset s$ . Now we have  $T \in \bigcap s_i = T$ , a contradiction. Hence  $T \in S$  and  $S$  is  $T$  is trivially the desired infimum. This proves  $\text{On}$  is an ordinal, so  $\text{On} \in \text{On}$ . Recall that  $\text{On}$  is a set, so  $\text{On} \notin \text{On}$ , a contradiction. This is known as the *Burali-Forti paradox*.

**Problem 3.27.** Prove that two different ordinals do not have the same order-type.

**Proposition.** If a set is well-ordered, then it has the same order-type as an ordinal.

This proof is slightly technical, you may skip the proof and remember the result.

*Proof.* Let  $W$  be well-ordered. For all  $x \in W$ , consider the initial segment  $I_x$  given by  $x$ . If  $I_x \approx \alpha_x$ , where  $\alpha_x$  is some ordinal, then define  $F(x) = \alpha_x$ . Suppose we have a set  $M \subset W$  consisting of elements  $x \in W$  such that  $I_x$  does not exist. Let  $m$  be the least element in  $M$ . Consider the set  $\{\alpha_y \mid y < m\}$ , this is a set of ordinals. Define  $\alpha = \sup\{\alpha_y \mid y < m\}$ . It is trivial that  $I_m \approx \alpha$ , a contradiction. Hence  $F : W \rightarrow \text{On}$  is well-defined, by the axiom of replacement,  $F(W) = \{\alpha_x\}$  is a set. For all  $x, y \in W$  and  $x < y$ ,  $I_x \subsetneq I_y$ , so  $\alpha_x \in \alpha_y$ , which is  $F(x) < F(y)$ . This also shows  $F$  is injective. Let  $\gamma = \min(\text{On} \setminus F(W))$ , then  $F(W) \subset \gamma$ . If  $\beta < \gamma$  is an ordinal, then  $\beta \in F(W)$ , so  $F(W) = \{\beta \mid \beta < \gamma\} = \gamma$ . Hence  $F : W \rightarrow \gamma$  is an order-preserving bijection.  $\square$

We can conclude that every well-ordered set has the same order-type as a unique ordinal.

**Proposition.** Every non-empty ordinal is either a successor ordinal or a limit ordinal.

*Proof.* Let  $\lambda$  be an ordinal and  $\lambda \neq \emptyset$ . Assume  $\lambda$  is not a successor ordinal, then  $\lambda \neq \alpha + 1$  for any ordinal  $\alpha$ . Now suppose  $\lambda$  is not a limit ordinal, then there exists  $\beta$  such that  $\beta < \lambda \not< \beta + 1$ . Since  $\lambda \neq \alpha + 1$  for all  $\alpha$ ,  $\lambda < \beta + 1$ , then either  $\lambda \in \beta$  or  $\lambda = \beta$ , a contradiction.  $\square$

**Problem 3.28.** Let  $C$  be a class of ordinals that satisfies:

1.  $0 \in C$ ;
2. if  $\alpha \in C$ , then  $\alpha + 1 \in C$ ;
3. if  $\alpha$  is a limit ordinal and  $\beta \in C$  for all  $\beta < \alpha$ , then  $\alpha \in C$ .

Use the previous proposition, prove that  $C$  is the class of all ordinals. This is called the *transfinite induction*.

**Axiom of choice.** Let  $X$  be a set and  $X \neq \emptyset$ . Then there exists a map  $f : \mathcal{P}(X) \rightarrow X$  such that for every  $x \subsetneq X$ , where  $x \neq \emptyset$ ,  $f(x) \in x$ .

We call such a function  $f$  a *choice function*. Therefore, we can reformulate the axiom of choice as: for every nonempty set  $X$ , there exists a choice function.

**Theorem 3.1** (well-ordering theorem). Every set is well-orderable.

*Proof.* Let  $S$  be a set, by the axiom of choice, there exists a choice function  $f : \mathcal{P}(S) \setminus \{\emptyset\} \rightarrow S$ . Let an ordinal  $\alpha = 0$  and let  $s_0 = f(S)$ , this is the basis for the transfinite induction. Assume  $s_\beta$  has been defined, where  $\beta < \alpha$ . If  $S \setminus \{s_\beta\} = \emptyset$ , then the map  $f : \{s_\beta\} \rightarrow \alpha$  defined by  $f(\beta) = s_\beta$  is a bijection, so  $S$  is well-ordered. If  $S \setminus \{s_\beta\} \neq \emptyset$ , define  $s_\alpha = f(S \setminus \{s_\beta\})$ . Suppose the process does not stop, then we have an injection from the class of all ordinal to the set  $S$ , by Burali-Forti paradox, this is a contradiction. Hence we have a bijection between  $\alpha$  and  $S$ , so  $S$  is well-ordered.  $\square$

By the well-ordering theorem,  $\mathbb{N}$  is well-ordered, this is called the *well-ordering principle*.

**Remark.** Well-ordering principle is based on ZF system, the system without axiom of choice. The well-ordering theorem is an equivalent form of axiom of choice. For more information, we recommend you to check [3].

## 4 Integers and Cardinality

Recall that we have constructed  $\mathbb{N}$  by axioms. By the convention of quotient sets, we could further construct the set of integers and rational numbers.

**Theorem 4.1** (recursion). Given a set  $X$  and  $x \in X$ . Let  $f : X \rightarrow X$  be a function, then there exists a unique function  $F : \mathbb{N} \rightarrow X$  such that  $F(0) = x$  and  $F(n^+) = f(F(n))$  for all  $n \in \mathbb{N}$ .

*Proof.* Construct such  $F$  inductively. Suppose  $F$  is not well-defined and  $F(n^+) = f(F(n)) = f(F(m))$  for some  $n, m \in \mathbb{N}$ , since  $f$  is well-defined, a contradiction. Suppose  $G$  is another function satisfies the property, then  $F(0) = x = G(0)$ . Assume  $F(n) = G(n)$  for some  $n \in \mathbb{N}$ , then  $F(n^+) = f(F(n)) = f(G(n)) = G(n^+)$ . Hence, by induction,  $F = G$ .  $\square$

**Definition 4.1.** The *addition* on  $\mathbb{N}$  is defined to be  $\{+_n\}_{n \in \mathbb{N}}$  such that  $+_n(0) = n$  and  $+_n(m^+) = (+_n(m))^+$  for a fixed  $n$  and every  $m \in \mathbb{N}$ . We denote  $+_n(m)$  by  $n + m$ .

**Proposition.** For all  $n \in \mathbb{N}$ ,  $n + 0 = n = 0 + n$ .

*Proof.* It is trivial that  $n + 0 = n$ . Let  $n = 0$ , then  $0 + 0 = 0$ . Suppose  $0 + k = k$  for some  $k \in \mathbb{N}$ , then  $0 + (k^+) = (0 + k)^+ = k^+$ . Hence, by induction,  $n + 0 = n = 0 + n$ .  $\square$

**Problem 4.1.** Prove that the following properties hold.

1.  $n + 1 = n^+$  for all  $n \in \mathbb{N}$ .
2.  $m + n = n + m$  for all  $m, n \in \mathbb{N}$ .
3.  $a + (b + c) = (a + b) + c$  for all  $a, b, c \in \mathbb{N}$ .

**Problem 4.2.** Consider a relation  $\sim_{\mathbb{N}^+} \subset \mathbb{N} \times \mathbb{N}$  defined by  $(a, b) \sim (c, d)$  if and only if  $a + d = b + c$ . Prove that  $\sim_{\mathbb{N}^+}$  is an equivalence relation.

**Definition 4.2.** The set of all *integers*, denoted  $\mathbb{Z}$ , is defined to be  $(\mathbb{N} \times \mathbb{N})_{\sim_{\mathbb{N}^+}}$ . Let  $[(a, b)] \in \mathbb{Z}$ , the *inverse* of  $[(a, b)]$  is defined to be  $[(b, a)]$ .

**Problem 4.3.** Prove that there exists a bijection between  $\{[(a, b)] \mid a, b \in \mathbb{N}, a \subsetneq b\}$  and  $\{[(b, a)] \mid a, b \in \mathbb{N}, b \subsetneq a\}$ , conclude that the inverse of an integer is unique.

Let  $n \in \mathbb{Z}$ , the inverse of  $n$  is denoted by  $-n$ .

**Definition 4.3.** Let  $[(a, b)], [(c, d)] \in \mathbb{Z}$ . The *addition* on  $\mathbb{Z}$  is defined to be  $[(a, b)] + [(c, d)] = [(a + c, b + d)]$ .

**Proposition.** The addition on  $\mathbb{Z}$  is well-defined.

*Proof.* Suppose  $(a, b) \sim (a', b')$  and  $(c, d) \sim (c', d')$ , then  $a + b' = b + a'$  and  $c + d' = d + c'$ . We have  $(a + c) + (b' + d') = a + b' + c + d' = b + a' + d + c' = (b + d) + (a' + c')$ , hence  $(a + c, b + d) \sim (a' + c', b' + d')$ .  $\square$

**Definition 4.4.** Let  $[(a, b)], [(c, d)] \in \mathbb{Z}$ , the *multiplication* on  $\mathbb{Z}$  is defined to be  $[(a, b)] \cdot [(c, d)] = [(ac + bd, ad + bc)]$ .

**Problem 4.4.** Prove that the multiplication on  $\mathbb{Z}$  is well-defined.

Recall that we have defined  $\leq$  on  $\mathbb{N}$  by the natural inclusion, that is, for all  $a, b \in \mathbb{N}$ ,  $a \leq b$  if and only if  $a \subset b$ . Let  $[(a, b)], [(c, d)] \in \mathbb{Z}$ , we say  $[(a, b)] \leq [(c, d)]$  if and only if  $a + d < b + c$ .

**Definition 4.5.** Let  $p \in \mathbb{Z}$ . We say  $p$  is a *prime* if  $p \neq a \cdot b$  for all  $a, b \in \mathbb{Z}$  and  $a, b \neq p$ .

**Problem 4.5.** Let  $n \in \mathbb{N}$  and  $n \neq 0$ . Prove that if there exists  $a, b \in \mathbb{N} \setminus \{0\}$  such that  $a \cdot b = n$ , then  $a, b \leq n$ .

**Theorem 4.2** (fundamental theorem of arithmetic). For all  $n \in \mathbb{N}$  and  $n \geq 2$ ,  $n$  can be factored as a product of prime numbers and the factorization is unique.

*Proof.* It is trivial that  $2 = 2$ ,  $3 = 3$ ,  $4 = 2 \cdot 2$ ,  $\dots$ . Suppose every number  $n \leq k$  for a given  $k$  admits a prime factorization. If  $k + 1$  is prime, then  $k + 1 = k + 1$ . If  $k + 1$  is not prime, then there exists  $n_1, n_2 \in \mathbb{Z}$  such that  $n_1 n_2 = k + 1$ . Since  $n_1$  and  $n_2$  admits prime factorization,  $k + 1$  admits prime factorization. We have the trivial case  $2 = 2$ . Assume the factorization is unique for all  $m$  with  $2 \leq m < n$ . Suppose  $\prod p_i$  and  $\prod q_j$  are prime factorizations of  $n$ . Rewrite those factorizations such that  $p_i \leq p_{i+1}$  and  $q_j \leq q_{j+1}$ . Since  $p_1$  divides  $\prod q_j$  and  $p_1$  is prime, it must divide one of the  $q_j$ , so  $p_1 \leq q_1 \leq q_j$ . Apply the same argument,  $q_1 \leq p_1$ , so  $q_1 = p_1$ . Now  $p_2 \cdots p_r = q_2 \cdots q_s < n$ . Hence, by induction,  $p_i = q_i$ , the factorization is unique.  $\square$

**Theorem 4.3** (Euclid's theorem). There are infinitely many primes.

*Proof.* Suppose there are finitely many primes, denoted  $p_1, \dots, p_n$ , where  $p_i \leq p_{i+1}$ . Define  $x = 1 + \prod_{i=1}^n p_i$ , then  $N$  is not prime. Let  $p_x$  be a prime in the factorization, then  $p_i$  lies in the factorization of  $N - \prod p_i = 1$ , a contradiction.  $\square$

**Problem 4.6.** Consider the relation  $\sim_{\mathbb{Z} \times} \subset \mathbb{Z} \times (\mathbb{Z} \setminus \{0\})$  defined by  $(a, b) \sim (c, d)$  if and only if  $ad = bc$ . Prove that  $\sim_{\mathbb{Z} \times}$  is an equivalence relation.

**Definition 4.6.** The *rational numbers*, denoted  $\mathbb{Q}$ , is defined to be the set  $(\mathbb{Z} \times (\mathbb{Z} \setminus \{0\}))_{\sim_{\mathbb{Z} \times}}$ . Let  $[(a, b)], [(c, d)] \in \mathbb{Q}$ , the *addition* is defined to be  $[(a, b)] + [(c, d)] = [(ad + bc, bd)]$  and the *multiplication* on  $\mathbb{Q}$  is defined to be  $[(a, b)] \cdot [(c, d)] = [(ac, bd)]$ .

**Problem 4.7.** Prove that addition and multiplication on  $\mathbb{Q}$  are well-defined.

**Problem 4.8.** Let  $0$  be the element  $[(0, x)] \in \mathbb{Q}$  and let  $1$  be the element  $[(x, x)] \in \mathbb{Q}$  for any  $x$ . Verify the following algebraic properties of  $\mathbb{Q}$ .

1.  $a + b = b + a$  and  $ab = ba$  for all  $a, b \in \mathbb{Q}$ . (commutativity)
2.  $a + b + c = a + (b + c)$  and  $abc = a(bc)$  for all  $a, b, c \in \mathbb{Q}$ . (associativity)
3.  $a + 0 = a$  for all  $a \in \mathbb{Q}$ . (additive identity)
4.  $a \cdot 1 = a$  for all  $a \in \mathbb{Q}$ . (multiplicative identity)
5. For all  $a \in \mathbb{Q}$ , there exists a unique  $-a \in \mathbb{Q}$  such that  $a + (-a) = 0$ . (additive inverse)
6. For all  $a \in \mathbb{Q}$  and  $a \neq 0$ , there exists a unique  $a^{-1} \in \mathbb{Q}$  such that  $aa^{-1} = 1$ . (multiplicative inverse)
7.  $a(b + c) = ab + ac$ . (distributivity)

**Definition 4.7.** Let  $a, b \in \mathbb{Q}$  and  $b \neq 0$ . The *subtraction* on  $\mathbb{Q}$  is defined to be  $a - b = a + (-b)$ . The *division* on  $\mathbb{Q}$  is defined to be  $a/b = a \cdot b^{-1}$ .

Let  $[(a, b)], [(c, d)] \in \mathbb{Q}$  with  $b, d \geq 0$  and  $b, d \neq 0$ , then  $[(a, b)] \leq [(c, d)]$  if and only if  $ad \leq bc$ .

**Proposition.** For all  $a, b, c \in \mathbb{Q}$ , the following properties hold.

1.  $a + c = b + c$  implies  $a = b$ .
2.  $a \cdot 0 = 0$ .
3.  $(-a)b = -ab$ .
4.  $(-a)(-b) = ab$ .
5.  $ac = bc$  and  $c \neq 0$  imply  $a = b$ .
6.  $ab = 0$  implies either  $a = 0$  or  $b = 0$ .

*Proof.* (i) We have  $a = a + (c + (-c)) = a + c + (-c) = b + c + (-c) = b + (c + (-c)) = b$ . (ii) We have  $a \cdot 0 = a \cdot (0 + 0) = a \cdot 0 + a \cdot 0$ , then  $a \cdot 0 + 0 = a \cdot 0 + a \cdot 0$ , so  $a \cdot 0 = 0$ . (iii) We have  $ab + (-a)b = (a + (-a))b = 0b = 0$ . Hence  $(-a)b = -ab$ . (iv) We have  $(-a)(-b) + (-ab) = (-a)(-b) + (-a)b = (-a)(-b + b) = 0$ . Hence  $(-a)(-b) = ab$ . (v) We have  $a = a(cc^{-1}) = acc^{-1} = bcc^{-1} = b(cc^{-1}) = b$ . (vi) Let  $b \neq 0$ , then  $ab = 0 = 0b$ , so  $a = 0$ .  $\square$

**Problem 4.9.** Prove that for all  $a, b, c \in \mathbb{Q}$ , the following properties hold.

1. If  $a \leq b$ , then  $-b \leq -a$ .
2. If  $a \leq b$  and  $c \leq 0$ , then  $bc \leq ac$ .
3. If  $0 \leq a$  and  $0 \leq b$ , then  $0 \leq ab$ .
4.  $0 \leq a^2$ .
5. If  $0 < a$ , then  $0 < a^{-1}$ .
6. If  $0 < a < b$ , then  $0 < b^{-1} < a^{-1}$ .

Now we consider the cardinality of infinite sets. We have constructed  $\mathbb{Z}$  and  $\mathbb{Q}$  based on  $\mathbb{N}$ , and all of those sets are considered to have the same cardinality.

**Definition 4.8.** A set  $S$  is said to be *countable* if  $S$  is finite or there exists a bijective function  $f : S \rightarrow \mathbb{N}$ . If a set is not countable, then we say the set is *uncountable*.

The function  $\text{id}_{\mathbb{N}} : \mathbb{N} \rightarrow \mathbb{N}$  is trivially bijective, so  $\mathbb{N}$  is countable.

**Definition 4.9.** Let  $X$  and  $Y$  be sets. We say  $X$  has smaller cardinality than  $Y$ , denoted  $|X| \leq |Y|$ , if there exists an injection  $f : X \rightarrow Y$ .

**Remark.** Since  $|X|$  can be infinite, the symbol “ $\leq$ ” is not the typical ordering in  $\mathbb{N}$ .

Similarly, one could define  $|X| \leq |Y|$  by surjections, that is, if there exists a surjection  $f : Y \rightarrow X$ , then  $|X| \leq |Y|$ .

**Theorem 4.4** (Cantor’s theorem). Let  $S$  be a set. Then any function  $f : S \rightarrow \mathcal{P}(S)$  is not surjective.

*Proof.* Suppose there exists a bijection  $f : S \rightarrow \mathcal{P}(S)$ . Define  $X = \{s \in S \mid s \notin f(s)\}$ , this is a subset of  $S$ , so  $X \in \mathcal{P}(S)$  and  $X = f(x)$  for some  $x \in S$ . If  $x \in X$ , then  $x \notin f(x) = X$ , so  $x \notin X$ , a contradiction. If  $x \notin X$ , then  $x \in f(x) = X$ , so  $x \in S$ , a contradiction.  $\square$

**Problem 4.10.** Use Cantor’s theorem, prove that for all set  $S$ , there exists some element  $x \notin S$ . Moreover, if  $S$  is finite, let  $x \notin S$ , prove that  $|S \cup \{x\}| = |S| + 1$ .

**Proposition.** Let  $X$  be a finite set and let  $Y \subset X$ , then  $Y$  is finite and  $|Y| \leq |X|$ .

*Proof.* Let  $f : Y \rightarrow X$  defined by  $f(y) = y \in Y \subset X$ , this map is trivially injective.  $\square$

**Problem 4.11.** Let  $X$  be a finite set and let  $f : X \rightarrow Y$  be a function. Prove that  $\text{im}(f)$  is finite and  $|f(X)| \leq |X|$ .

**Problem 4.12.** Let  $X$  and  $Y$  be finite sets. Prove that  $|X \cup Y| \leq |X| + |Y|$ .

**Proposition.** The finite Cartesian product of countable set is countable.

*Proof.* Let  $A_1, A_2, \dots, A_n$  be countable sets, then each  $A_i$  itself is in bijection with  $\mathbb{N}$ . Suppose we pick  $n$  distinct prime numbers  $p_1, p_2, \dots, p_n$  and define  $f : \mathbb{N}^n \rightarrow \mathbb{N}$  such that  $f(a_1, a_2, \dots, a_n) = p_1^{a_1+1} \cdot p_2^{a_2+2} \cdot \dots \cdot p_n^{a_n+n}$ . Notice  $f$  is injective as the prime factorization of a number is unique. Hence  $\mathbb{N}^n$  is countable then  $\prod_{i=1}^n A_i$  is countable.  $\square$

**Proposition.** The sets  $\mathbb{Z}$  and  $\mathbb{Q}$  are countable.

*Proof.* Since  $\mathbb{N}$  is countable,  $\mathbb{N} \times \mathbb{N}$  is countable. The natural projection  $\mathbb{N} \times \mathbb{N} \rightarrow \mathbb{Z}$  is surjective, so  $\mathbb{Z}$  is countable. Since  $\mathbb{Z}$  is countable,  $\mathbb{Z} \setminus \{0\} \subset \mathbb{Z}$ , so  $\mathbb{Z}$  is countable. Now  $\mathbb{Z} \times (\mathbb{Z} \setminus \{0\})$  is countable, since  $\mathbb{Q}$  is a quotient set of  $\mathbb{Z} \times (\mathbb{Z} \setminus \{0\})$ ,  $\mathbb{Q}$  is countable.  $\square$

**Theorem 4.5** (Cantor-Schröder-Bernstein theorem). Let  $f : A \rightarrow B$  and  $g : B \rightarrow A$  be injective functions. Then there exists a bijective function  $h : A \rightarrow B$ .



**Remark.** Cantor-Schröder-Bernstein theorem was originally proved as a consequence of the axiom of choice. However, it is provable in ZF, the system without the axiom of choice. In the next proof, assume we have all axioms except for the axiom of choice. This proof is technical and you may skip it.

*Proof.* Let  $f : A \rightarrow B$  and  $g : B \rightarrow A$  be injections. Define  $A_0 = A \setminus g(B)$ . For any  $n \geq 0$ , let  $B_n = f(A_n)$  and let  $A_{n+1} = g(B_n)$ . Since  $g$  is injective,  $g(B_n)$  are pairwise disjoint. It is clear that by induction,  $A_n$  are pairwise disjoint. Similarly,  $B_n$  are pairwise disjoint. Let  $U = \bigcup A_n$  and  $V = A \setminus U$ . We have  $B \setminus B_0 = B \setminus f(A_0)$ ,  $V \subset A \setminus A_1 = g(B \setminus f(A_0))$ . Here there exists a bijection between  $B \setminus f(A_0)$  and  $g(B \setminus f(A_0))$ . The restriction  $g|_V : V \rightarrow A$  is a bijection between  $B \setminus B_0$  and  $V$ . Now let  $h : A \rightarrow B$  be a function defined by  $h(x) = f(x)$  if  $x \in U$  and  $h(x) = g^{-1}(x)$  if  $x \in V$ . Suppose  $h(x_1) = h(x_2)$ . If  $x_1, x_2 \in U$ , then  $h(x_i) = f(x_i) \in B_n$  for some  $n$ . The restriction of  $f$  on every  $A_n$  is injective, so  $x_1 = x_2$ . Similarly, if  $x_1, x_2 \in V$ ,  $x_1 = x_2$ . If  $x_1 \in U$  and  $x_2 \in V$ , then  $h(x_1) \in B_n$  for some  $n$ , where  $B_n \cap (B \setminus B_0) = \emptyset$ . Hence  $h$  is injective. Pick some  $y \in B$ . If  $y \in B_n$  for some  $n$ , then there is a unique  $x \in A_n$  such that  $f(x) = y$ , so  $h(x) = f(x) = y$ . If  $y \notin B_n$ , then  $y \in B \setminus B_0$ . Consider  $x = g(y) \in V$ , then  $h(x) = h(g(y)) = g^{-1}(g(y)) = y$ . Hence  $h$  is surjective, conclude that  $h$  is a bijection.  $\square$

**Problem 4.13.** Prove the Cantor-Schröder-Bernstein theorem using the axiom of choice.

By Cantor-Schröder-Bernstein theorem, we conclude that if  $|A| \leq |B|$  and  $|B| \leq |A|$ , then  $|A| = |B|$ .

## 5 Vector Spaces and Linear Maps

## 6 Functions as Morphisms

Functions are the bridges to “connect” sets. According to the definition, a function takes a set to another while preserving the structure of sets. This property allows us to consider functions as morphisms between sets.

**Definition 6.1.** A *category*  $C$  consists of:

1. a class, denoted  $\text{Ob}(C)$ , of *objects*;
2. for each pair of objects  $X$  and  $Y$ , there exists a class of *morphisms*  $f : X \rightarrow Y$ , where  $X$  is called the *domain* and  $Y$  is called the *codomain*;
3. a *composition operation*, which gives, for each pair of morphisms  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$ , a morphism  $g \circ f : X \rightarrow Z$ ,

such that


1. given any  $f : X \rightarrow Y$ ,  $g : Y \rightarrow Z$ , and  $h : Z \rightarrow W$ , we have the identity  $(h \circ g) \circ f = h \circ (g \circ f)$ ;
2. for each object  $X$ , there exists an *identity morphism*  $\text{id}_X : X \rightarrow X$  with the property that  $f \circ \text{id}_X = f$  and  $\text{id}_Y \circ g = g$  for any  $f : X \rightarrow Y$  and  $g : Z \rightarrow Y$ .

The class of morphisms from  $X$  to  $Y$  is denoted by  $\text{Hom}_C(X, Y)$ . A category  $C$  is a *locally small category* if  $\text{Hom}_C(A, B)$  is a set for all objects  $A$  and  $B$  in  $C$ .

Let sets be objects, let functions be morphisms, and let the composition of morphisms be the composition of functions. By our previous observations, it suffices to check this defines a category of sets, denoted  $\mathbf{Set}$ . Moreover,  $\mathbf{Set}$  is locally small.

**Example.** Let  $X$  be a set, then  $\text{Hom}_{\mathbf{Set}}(X, X)$  is a category.

**Example.** In the following diagram, let each vertex be an object and let each arrow be a morphism. This defines a category.



**Problem 6.1.** Let the objects be monoids and let the morphisms be monoid homomorphisms. Prove that this defines a category, denoted  $\mathbf{Mon}$ . Let the objects be groups and let the morphisms be group homomorphisms. Prove that this defines a category, denoted  $\mathbf{Grp}$ . Let the objects be topological spaces and let the morphisms be continuous maps. Prove that this defines a category, denoted  $\mathbf{Top}$ .

**Problem 6.2.** Morphisms are not guaranteed to be functions. Let  $(S, \leq)$  be a partially ordered set. Let  $\text{Ob}(C) = S$  and  $x \rightarrow y$  be a morphism if  $x \leq y$ . Prove that  $C$  is a category.

**Problem 6.3.** Let  $C$  be a category. Let a system  $C^{op}$  consists of all objects in  $C$  and all morphisms  $f : A \rightarrow B$  if  $B \rightarrow A$  is a morphism in  $C$ . Prove that  $C^{op}$  is indeed a category. This is called the *dual category* of  $C$ .

**Problem 6.4.** Prove that the identity morphism is unique for each object  $A$  in a category  $C$ .

**Definition 6.2.** Let  $C$  be a category. A category  $D$  is a *subcategory* of  $C$  if the  $\text{Ob}(D)$  and the  $\text{Hom}_D(X, Y)$  are subcollections of  $\text{Ob}(C)$  and  $\text{Hom}_C(X, Y)$ , respectively, for all objects  $X$  and  $Y$  in  $C$ . The subcategory  $D$  is said to be *full* if for all objects  $X$  and  $Y$  in  $D$ ,  $\text{Hom}_D(X, Y)$  is exactly  $\text{Hom}_C(X, Y)$ .

**Example.** The category  $\mathbf{Grp}$  is a subcategory of  $\mathbf{Set}$ .

**Problem 6.5.** Let the objects be abelian groups, let the morphisms be group homomorphisms, and let the composition of morphisms be the composition of functions. Prove that this defines a category, denoted  $\mathbf{Ab}$ . Prove that  $\mathbf{Ab}$  is a full subcategory of  $\mathbf{Grp}$ .

**Definition 6.3.** Let  $C$  be a category. A morphism  $f : A \rightarrow B$  is a *monomorphism* if for all  $g, h : C \rightarrow A$ ,  $f \circ g = f \circ h$  implies  $g = h$ . A morphism  $f : A \rightarrow B$  is called an *epimorphism* if for all  $i, j : B \rightarrow D$ ,  $i \circ f = j \circ f$  implies  $i = j$ .

**Proposition.** A function is injective if and only if it is a monomorphism in **Set**.

*Proof.* ( $\Rightarrow$ ) Let  $f : A \rightarrow B$  be injective. If  $A = \emptyset$ , then  $f$  is the unique empty function. If  $A \neq \emptyset$ , let  $g : B \rightarrow A$  be the left inverse of  $f$ , ( $\Leftarrow$ ) Let  $f : A \rightarrow B$  be a monomorphism. If  $A = \emptyset$ , then  $f : \emptyset \rightarrow B$  is vacuously injective. If  $A \neq \emptyset$ ,  $\square$

**Problem 6.6.** Prove that a function is surjective if and only if it is an epimorphism in **Set**.

**Definition 6.4.** Let  $C$  be a category. A morphism  $f : A \rightarrow B$  is an *isomorphism* if there exists  $g \in \text{Hom}_C(B, A)$  such that  $f \circ g = \text{id}_B$  and  $g \circ f = \text{id}_A$ . If there exists a morphism between two objects  $A$  and  $B$ , then we say they are *isomorphic*, denoted  $A \approx B$ .

**Problem 6.7.** Prove that a morphism is an isomorphism if and only if it is a monomorphism and an epimorphism.

**Problem 6.8.** Let  $A$  and  $B$  be well-ordered sets that are isomorphic to each other. Prove that the isomorphism between them is unique.

**Definition 6.5.** An *initial object*  $0$  of a category  $C$  is an object in  $C$  such that for any object  $A$ , there is a unique morphism  $0 \rightarrow A$ . A *terminal object*  $1$  of a category  $C$  is an object of  $C$  such that for any object  $B$ , there is a unique morphism  $B \rightarrow 1$ . We say an object is a *zero object* if it is both an initial object and a terminal object.

**Proposition.** Initial objects of a category  $C$  are unique up to isomorphism.

*Proof.* Let  $0$  and  $0'$  be initial objects in some category  $C$ . Let  $\varphi : 0 \rightarrow 0'$  and  $\psi : 0' \rightarrow 0$ . Since the diagram is commutative, we have  $\varphi \circ \psi \circ \varphi = \varphi \circ \text{id}_0$ , hence  $\varphi$  is an isomorphism.  $\square$

$$\begin{array}{ccc}
 0 & \xrightarrow{\varphi} & 0' \\
 & \searrow \text{id}_0 & \downarrow \psi \\
 & & 0 \xrightarrow{\varphi} 0'
 \end{array}$$

**Problem 6.9.** Prove that the initial object in a category  $C$  is a terminal object in  $C^{\text{op}}$ , conclude that terminal objects in  $C$  are unique up to isomorphism.

**Problem 6.10.** Prove that the initial object in the category of partially ordered set is the  $\leq$ -minimal element. Prove that the terminal object in this category is the  $\leq$ -maximal element.

**Proposition.** The initial object in **Set** is  $\emptyset$  and the terminal object in **Set** is a singleton.

*Proof.*  $\square$

We conclude that there is no zero object in **Set**.

**Universal property for quotient sets.** Let  $X$  and  $Y$  be sets. Let  $\sim$  be an equivalence relation on  $X$  and let  $f : X \rightarrow Y$  be invariant under the equivalence relation. Then there exists a unique function  $\bar{f} : X_{\sim} \rightarrow Y$  such that  $f = \bar{f} \circ \pi$ .

$$\begin{array}{ccc}
 X_{\sim} & \xleftarrow{\pi} & X \\
 & \searrow \bar{f} & \downarrow f \\
 & & Y
 \end{array}$$

The proof of the universal property has two parts. We first verify that a quotient set has the universal property. Then we verify that a quotient set is characterized by this universal property, that is, any set satisfying the universal property must be a quotient set. This proof will be separated into multiple propositions.

**Proposition.** There exists a map  $\bar{f} : X_{\sim} \rightarrow Y$  such that  $f = \bar{f} \circ \pi$ .

*Proof.* Define the relation  $\{([x], y)\} \in \bar{f}$  if and only if  $f(x) = y$ . Let  $[x] \in X$  and  $y, z \in Y$ . Suppose  $([x], y), ([x], z) \in \bar{f}$ , then there exists  $u, v \in X$  such that  $[u] = [v] = [x]$ ,  $f(u) = y$ , and  $f(v) = z$ , so  $u \sim v$ . Since  $f$  is invariant under  $\sim$ ,  $y = z$ . Hence  $\bar{f}$  is well-defined. For all  $x \in X$ ,  $(\bar{f} \circ \pi)(x) = \bar{f}([x]) = f(y)$ .  $\square$

**Problem 6.11.** Prove that such  $\bar{f}$  is unique.

**Problem 6.12.** Let  $f : X \rightarrow Y$  be a function that induces an equivalence relation  $\sim_f$  on  $X$ . Since  $X_{\sim_f}$  satisfies the universal property, prove that  $X_{\sim_f} \approx \text{im}(f)$ .

**Proposition.** Let  $S$  be any set satisfying the universal property for quotient sets. Let  $X$  be a set and let  $\sim_1$  be an equivalence relation on  $X$ , so  $\pi_1 : X \rightarrow X_{\sim_1}$  is invariant under the equivalence relation induced by  $\pi_2 : X \rightarrow S$ . Then  $S \approx X_{\sim_1}$ .

$$\begin{array}{ccc} S & \xleftarrow{\pi_2} & X \\ & \searrow \bar{\pi}_1 & \downarrow \pi_1 \\ & & X_{\sim_1} \end{array}$$

*Proof.* Since  $\pi_1$  is surjective,  $\bar{\pi}_1$  is surjective. Suppose  $\bar{\pi}_1(s_1) = \bar{\pi}_1(s_2)$ , where  $s_1, s_2 \in S$ . Since  $\pi_2$  is surjective, there exist  $x, y \in X$  with  $s_1 = \pi_2(x)$  and  $s_2 = \pi_2(y)$ . Then  $\bar{\pi}_1(s_1) = \bar{\pi}_1(\pi_2(x)) = \pi_1(x)$  and  $\bar{\pi}_1(s_2) = \bar{\pi}_1(\pi_2(y)) = \pi_1(y)$ , which implies  $\pi_1(x) = \pi_1(y)$ , so  $[x] = [y]$ . Since  $\pi_2$  is invariant under  $\sim_1$ , it follows that  $\pi_2(x) = \pi_2(y)$ , so  $s_1 = s_2$ . Hence  $S \approx X_{\sim_1}$ .  $\square$

This completes our proof of the universal property for quotient sets.

**Theorem 6.1** (canonical decomposition of functions). Let  $f : A \rightarrow B$  be a function, then there exists a surjective function  $\pi$ , a bijective function  $g$ , and an injective function  $i$ , such that  $f = i \circ g \circ \pi$ .

$$\begin{array}{ccccc} A & \xrightarrow{\pi} & A_{\sim} & \xrightarrow{g} & \text{im}(f) \\ & \searrow f & & & \downarrow i \\ & & & & B \end{array}$$

*Proof.* Let  $i : \text{im}(f) \rightarrow B$  be the identity map, then  $i$  is injective. The function  $f : A \rightarrow B$  induces an equivalence relation  $\sim$  on  $X$ , so  $g$  is bijective. The projection map  $\pi : A \rightarrow A_{\sim}$  is surjective. For all  $a \in A$ ,  $(i \circ g \circ \pi)(a) = (i \circ g)([a]) = i(f(a)) = f(a)$ .  $\square$


**Definition 6.6.** Let  $\mathcal{C}$  be a category. Let  $A$  and  $B$  be objects in  $\mathcal{C}$ . The *product* of  $A$  and  $B$ , denoted  $P$ , is an object in  $\mathcal{C}$  with morphisms  $p_1 : P \rightarrow A$  and  $p_2 : P \rightarrow B$  such that for all object  $X$  in  $\mathcal{C}$  with morphisms  $\varphi_1 : X \rightarrow A$  and  $\varphi_2 : X \rightarrow B$ , there exists a unique  $u : X \rightarrow P$  such that the following diagram commutes.

$$\begin{array}{ccccc} & & X & & \\ & \swarrow \varphi_1 & \downarrow u & \searrow \varphi_2 & \\ A & \xleftarrow{p_1} & P & \xrightarrow{p_2} & B \end{array}$$

**Problem 6.13.** Prove that in  $\mathbf{Set}$ , the Cartesian product satisfies the universal property for product.


**Proposition.** Let  $A$  and  $B$  be objects in a category  $\mathcal{C}$ . Then their product is unique up to isomorphism.

*Proof.* Let  $P$  with  $p_1 : P \rightarrow A$  and  $p_2 : P \rightarrow B$  be the product of  $A$  and  $B$ . Suppose  $Q$  with  $\varphi_1 : Q \rightarrow A$  and  $\varphi_2 : Q \rightarrow B$  is another product of  $A$  and  $B$ . By the universal property,  $u : Q \rightarrow P$  and  $v : P \rightarrow Q$  are unique such that  $p_1 \circ u = \varphi_1$ ,  $p_2 \circ u = \varphi_2$ ,  $\varphi_1 \circ v = p_1$ , and  $\varphi_2 \circ v = p_2$ . We have  $\varphi_1 \circ u \circ v = \varphi_1$  and  $\varphi_2 \circ u \circ v = \varphi_2$ , then  $u \circ v = \text{id}_Q$ . Similarly,  $v \circ u = \text{id}_P$ . Hence  $P \approx Q$ .  $\square$



**Problem 6.14.** Let  $A$ ,  $B$ , and  $C$  be sets. Use the universal property to prove that  $A \times (B \times C) = (A \times B) \times C$ .

**Definition 6.7.** Let  $\mathcal{C}$  be a category. Let  $A$  and  $B$  be objects in  $\mathcal{C}$ . The *coproduct* of  $A$  and  $B$ , denoted  $Q$ , is an object in  $\mathcal{C}$  with morphisms  $q_1 : A \rightarrow Q$  and  $q_2 : B \rightarrow Q$  such that for all object  $Y$  in  $\mathcal{C}$  with morphisms  $\varphi_1 : A \rightarrow Y$  and  $\varphi_2 : B \rightarrow Y$ , there exists a unique  $v : Q \rightarrow Y$  such that the diagram commutes.



**Problem 6.15.** Prove that in  $\mathbf{Set}$ , the disjoint union satisfies the universal property for coproduct. Prove that the universal property characterizes disjoint union.

**Problem 6.16.** Describe the product and coproduct in  $\mathbf{Set}^{op}$ .

Categorically, we can restate the axiom of choice as follows.

**Axiom of choice.** Any epimorphism in  $\mathbf{Set}$  has a right inverse.

We will show how axiom of choice implies our categorical reformulation.

## 7 Groups

**Definition 7.1.** A *monoid* is a triple  $(M, \circ, e)$ , where  $M$  is a set and  $\circ : M \times M \rightarrow M$ , called a *composition*, is a function such that:

1. for all  $a, b, c \in M$ ,  $a \circ (b \circ c) = (a \circ b) \circ c$ ; (associative)
2. there exists  $e \in M$  such that for all  $x \in M$ ,  $e \circ x = x = x \circ e$ . (identity)

Here  $e$  is called an *identity*.

**Remark.** We will use  $gf$  for  $g \circ f$ .

**Example.** The natural numbers  $(\mathbb{N}, +, 0)$  is a monoid.

**Example.** Given a set  $S$ , the set of all functions  $f : S \rightarrow S$  forms a monoid with the usual composition of function, denoted  $M(S)$ .

**Problem 7.1.** Prove that the identity element is unique in any monoid.

**Definition 7.2.** A triple  $(G, \circ, e)$  is a *group* if it is a monoid and for all  $g \in G$ , there exists  $h \in G$  such that  $hg = e = gh$ . Such  $f$  is called an *inverse* of  $g$ . If  $\circ$  is commutative, then the group is *abelian*. The *order* of a group  $G$ , denoted  $|G|$ , is the cardinality of the set  $G$ .

**Example.** The *trivial group* is the group of one element, that is,  $\{e\}$ .

**Example.** Consider the set  $D_n$  of all rotations and reflections that map a regular  $n$ -gon into itself, this is called the *dihedral group*. Let the reflection be  $S$ , which gives  $S^2 = e$ . Multiply the rotations by  $S$  on the left, then we have  $n$  distinct rotational symmetries. Hence there are  $n$  rotations and  $n$  reflections, that is,  $|D_n| = 2n$ .

**Example.** Let  $S$  be a set. All bijections  $S \rightarrow S$  forms a group called the *symmetric group*, denoted  $\text{Sym}(S)$ . If  $S$  is finite, then the symmetric group is denoted by  $\mathfrak{S}_n$ , where  $|S| = n$ . An element of the symmetric group is called a *permutation*. Let  $\pi$  be a permutation of  $\mathfrak{S}_n$ . A permutation can be expressed by two-line notation, that is,

$$\pi = \begin{pmatrix} 1 & 2 & \cdots & n \\ \pi(1) & \pi(2) & \cdots & \pi(n) \end{pmatrix}.$$

The first line can be dropped to form an one-line notation. Given  $1 \leq i \leq n$  and  $i \in \mathbb{Z}$ , we can write  $\pi$  in cycle notation, that is,  $(i, \pi(i), \pi^2(i), \dots, \pi^{p-1}(i))$ , where  $p$  is the first integer such that  $\pi^p(i) = i$ . Such a cycle means that  $\pi$  sends  $i$  to  $\pi(i)$ ,  $\pi(i)$  to  $\pi^2(i)$ , and eventually,  $\pi^{p-1}(i)$  back to  $i$ . The cycle type of a permutation is an expression of the form  $(1^{m_1}, 2^{m_2}, \dots, n^{m_n})$ , where  $m_k$  is the number of cycles of length  $k$  in  $\pi$ . An *involution* is a permutation  $\pi$  such that  $\pi^2 = e$ .

**Problem 7.2.** Prove that every permutation in  $\mathfrak{S}_n$  can be written as a finite composition of involutions. Is this expression unique?

**Definition 7.3.** A permutation  $\pi \in \mathfrak{S}_n$  is called an *odd permutation* if it can be written as the composition of an odd number of involutions. A permutation  $\pi \in \mathfrak{S}_n$  is called an *even permutation* if it can be written as the composition of an even number of involutions.

**Proposition.** Let  $G$  be a group. For all  $g \in G$ , the inverse of  $g$  is unique.

*Proof.* For all  $g \in G$ , let  $f$  and  $h$  be two inverses, then  $f = f(gh) = (fg)h = h$ , hence the inverse of  $g$  is unique.  $\square$

For all  $g \in G$ , where  $G$  is a group, the inverse of  $g$  is denoted by  $g^{-1}$ .

**Problem 7.3.** Let  $G$  be a group and let  $g \in G$ . A *left inverse* of  $g$  is an element  $h \in G$  such that  $hg = e$ . Similarly, we can define a right inverse of  $g$ . Prove that a left inverse is equivalent to a right inverse, conclude that any left or right inverse of an element is its inverse.

**Problem 7.4.** Let  $G$  be a group and let  $g, h \in G$ . Prove that  $(-e)g = -g$ ,  $-(-g) = g$ , and  $-g(-h) = gh$ .

**Problem 7.5.** Let  $G$  be a group and let  $g_i \in G$ , define the composition of finitely many elements by  $\prod_{i=1}^n x_1 \cdots x_n = (x_1 \cdots x_{n-1})x_n$ , prove that  $\prod_{i=1}^m x_i \prod_{j=1}^n x_{m+j} = \prod_{j=1}^{m+n} x_j$ . This is called the *generalized associativity*.

**Problem 7.6.** Prove that  $|\mathfrak{S}_n| = n!$

**Definition 7.4.** Let  $G$  and  $H$  be monoids. A function  $\varphi : G \rightarrow H$  is called a *monoid homomorphism* if  $\varphi(ab) = \varphi(a)\varphi(b)$  and  $\varphi(e_G) = e_H$  for all  $a, b \in G$ . Let  $G$  and  $H$  be groups, a function  $\varphi : G \rightarrow H$  is called a *group homomorphism* if  $\varphi$  is a monoid homomorphism. The set  $\{x \in G \mid \varphi(x) = e_H\}$  is called the *kernel* of  $\varphi$ , denoted  $\ker(\varphi)$ . If two groups  $G$  and  $H$  are isomorphic to each other, then we write  $G \approx H$ .

**Problem 7.7.** Prove that a group isomorphism is a bijective group homomorphism.

**Definition 7.5.** A *subgroup* of a group  $(G, \circ, e)$  is a subset  $H \subset G$  containing  $e$  such that  $(H, \circ, e)$  is a group. If  $H$  is a subgroup of  $G$ , we write  $H \leq G$ .

**Definition 7.6.** Let  $G$  be a group. Every homomorphism  $f : G \rightarrow G$  is called an *automorphism*. The set of all automorphisms onto a group  $G$  is denoted by  $\text{Aut}(G)$ .

**Problem 7.8.** Let  $G$  be a group and let  $\circ$  be the composition of functions. Prove that  $(\text{Aut}(G), \circ, \text{id}_G)$  is a group. This is called the *automorphism group*.

**Proposition.** Let  $G$  be a group, then  $H \leq G$  if and only if  $e \in H \subset G$ , for all  $g, f \in H$ ,  $gf^{-1} \in H$ .

*Proof.*  $(\Rightarrow)$  Trivial.  $(\Leftarrow)$  Since  $H \subset G$ , the composition is associative. For all  $g \in H$ , take  $f = g$ , then  $gg^{-1} = e$ ; take  $f = e$ , then  $g^{-1} \in H$  for all  $g$ , hence  $H \leq G$ .  $\square$

**Problem 7.9.** If  $A \leq B \leq G$ , prove that  $A \leq G$ .

**Problem 7.10.** Let  $G$  be a group. Prove that all subgroups of  $G$  form a set. Prove that the set is partially ordered under " $\subset$ ". This is called the *lattice of subgroups* of  $G$ .

**Definition 7.7.** Let  $H \leq G$  be a subgroup. A *left coset* for  $H$  is a set of the form  $xH = \{xh \mid h \in H\}$ . A right coset is of the form  $Hx = \{hx \mid h \in H\}$ . The set of all left cosets for  $H$  is denoted by  $G/H$  and the set of all right cosets for  $H$  is denoted by  $H \backslash G$ . The *index* of  $H$  in  $G$ , denoted  $[G : H]$ , is defined to be  $|G/H|$ .

**Problem 7.11.** Prove that there exists a bijection  $\psi : G/H \rightarrow H \backslash G$ , conclude that  $[G : H] = |H \backslash G|$ .

**Problem 7.12.** Let  $H \leq G$  be a subgroup. Prove that there exists a set of left cosets that partitions  $G$ .

**Proposition.** Let  $H \leq G$  be a subgroup and let  $1$  be the trivial group, then  $[G : 1] = [G : H][H : 1]$ .

*Proof.*  $\square$

**Problem 7.13.** Based on the previous proposition, prove a more general form:

**Theorem 7.1** (Cayley's theorem). Any monoid is isomorphic to a submonoid of  $M(S)$  for some set  $S$ . Any group is isomorphic to a subgroup of some symmetric group.

*Proof.* Let  $M$  be a monoid. For all  $\alpha \in M$ , let  $\alpha_l(\alpha) = \alpha x$  for all  $x \in M$ , then  $\alpha_l$  maps  $M$  to itself. Consider  $S = \{\alpha_l \mid \alpha \in M\}$ , which is a subset of  $M(S)$ . The identity map  $\alpha_e \in S$ . For all  $\alpha, \beta \in M$ ,  $\alpha_l(\beta_l(x)) = \alpha_l(\beta x) = \alpha \beta x = (\alpha \beta)_l(x) = (\alpha \beta)_l(x)$ , so  $S$  is a submonoid of  $M(S)$ . Consider the map  $\varphi(\alpha) = \alpha_l$ . For all  $\alpha, \beta \in M$ ,  $\varphi(\alpha)\varphi(\beta) = \alpha_l \beta_l = (\alpha \beta)_l = \varphi(\alpha \beta)$ . The map is trivially surjective. Let  $\varphi(\alpha) = \varphi(\beta)$ , then  $\alpha_l = \beta_l$ , that is,  $\alpha x = \beta x$ . Consider  $x = 1$ , then  $\alpha = \beta$ , hence  $\varphi$  is an isomorphism. Now consider a group  $G$  and construct the same set  $S$ . For all  $\alpha_l$ , the inverse is  $(\alpha^{-1})_l$ . We have  $\alpha_l(\alpha^{-1})_l(x) = \alpha_l(\alpha^{-1}x) = x$  and  $(\alpha^{-1})_l \alpha_l(x) = (\alpha^{-1})_l(\alpha x) = x$ , hence  $S$  is a subgroup of some symmetric group. Construct the same  $\varphi$ , then  $S$  is isomorphic to  $G$ .  $\square$



**Definition 7.8.** Let  $H \leq G$  be a subgroup. We say  $H$  is a *normal subgroup* of  $G$ , denoted  $H \trianglelefteq G$ , if  $xH = Hx$  for all  $x \in G$ . A non-trivial group  $G$  is said to be *simple* if the only normal subgroups of  $G$  are  $G$  and the trivial subgroup.

**Problem 7.14.** Prove that  $H \trianglelefteq G$  if and only if  $gHg^{-1} = H$  for all  $g \in G$ .

**Theorem 7.2** (first isomorphism theorem). Let  $\varphi : G \rightarrow H$  be a group homomorphism. Then  $\ker(\varphi) \trianglelefteq G$ ,  $\text{im}(\varphi) \leq H$ , and  $G/\ker(\varphi) \approx \text{im}(\varphi)$ .

$$\begin{array}{ccccc} G & \xrightarrow{\pi} & G/\ker(\varphi) & \xrightarrow{\quad} & \text{im}(\varphi) \\ & & & & \downarrow i \\ & & & \searrow \varphi & H \end{array}$$

*Proof.* Let  $a, b \in \ker(\varphi)$ , then  $\varphi(ab^{-1}) = \varphi(a)\varphi(b) = \varphi(b^{-1})$ . We have  $e_H = \varphi(bb^{-1}) = \varphi(b)\varphi(b^{-1})$ , then  $\varphi(b^{-1}) = e_H$ , which implies  $\ker(\varphi) \leq G$ . For all  $g \in G$ ,  $\varphi(g\ker(\varphi)g^{-1}) = \varphi(g)\varphi(\ker(\varphi))\varphi(g^{-1}) = e_H$ , then  $g\ker(\varphi)g^{-1} \subset \ker(\varphi)$ . For all  $g \in \ker(\varphi)$ , we have  $\square$

**Problem 7.15.** Let  $C_2$  be a group of order 2. Prove that  $C_2$  is unique up to isomorphism. Define the set  $C_2 = \{1, -1\}$ . Let 1 be the identity. For all  $\pi \in \mathfrak{S}_n$ , define  $\varphi : \mathfrak{S}_n \rightarrow C_2$  by  $\varphi(\pi) = -1$  if  $\pi$  is an odd permutation and  $\varphi(\pi) = 1$  if it is an even permutation. Prove that  $\text{sgn}$  is a well-defined group homomorphism. The kernel of  $\varphi$  is called the *alternating group*, denoted  $\mathfrak{A}_n$ . Prove that  $\mathfrak{A}_n$  is abelian if  $n \leq 3$  and prove that  $\mathfrak{A}_n$  is not abelian for  $n > 3$ , conclude that a normal subgroup is not necessarily abelian.

**Definition 7.9.** Let  $G$  be a group and  $N \trianglelefteq G$ . The *quotient group*  $G/N$  is defined to be  $\{aN \mid a \in G\}$ .

**Problem 7.16.** Take  $(aN)(bN) = (ab)N$  as the composition on  $G/N$ , prove that  $G/N$  is indeed a group.

**Definition 7.10.** Let  $X$  be a subset of a group  $F$ . Then  $F$  is a *free group* with *basis*  $X$  if for any function  $\varphi : X \rightarrow G$ , where  $G$  is a group, there exists a unique extension  $\bar{\varphi} : F \rightarrow G$  such that  $i \circ \bar{\varphi} = \varphi$ .

$$\begin{array}{ccc} F & \xleftarrow{i} & X \\ & \searrow \bar{\varphi} & \downarrow \varphi \\ & & G \end{array}$$

We will first show the existence of free groups. Consider an arbitrary set  $X$ . The elements of  $X$  are called *generators*. Let  $Y$  be a disjoint copy of  $X$  with a bijection  $f : X \rightarrow Y$ . For all  $x \in X$ , if  $f(x) = y$ , then we say  $y$  is the *inverse* of  $x$ , denoted  $x^{-1}$ . We can rewrite  $Y = X^{-1}$  and denote  $X^{\pm 1} = X \cup X^{-1}$ .

**Definition 7.11.** Let  $X$  be a set. The elements of  $X^{\pm 1}$  are called *letters*. A *word*  $w$  is a finite sequence  $(l_1, \dots, l_n)$  of letters. The *length* of  $w = (l_1, \dots, l_n)$ , denoted  $|w|$ , is the value  $n$ . If  $n = 0$ , we call it the *empty word*, denoted 1. A word is *reduced* if it contains no part in the form  $aa^{-1}$ .

Let the set  $F(X)$  be the set of all reduced words. Let  $x = (x_1, \dots, x_m), y = (y_1, \dots, y_n) \in F(X)$ . Pick the largest  $k$  such that  $0 \leq k \leq \min(m, n)$  and  $x_{m-i} = y_{i+1}^{-1}$  for all  $i \leq k$ . Define the composition  $\cdot : F(X) \times F(X) \rightarrow F(X)$  to be the following function:

$$x \cdot y = \begin{cases} (x_1, \dots, x_{m-k}, y_{k+1}, \dots, y_n) & \text{if } k < \min(m, n) \\ (x_1, \dots, x_{m-k}) & \text{if } k = n < m \\ (y_{k+1}, \dots, y_n) & \text{if } k = m < n \\ 1 & \text{if } k = m = n \end{cases}$$

**Problem 7.17.** Prove that  $\cdot$  is a well-defined composition and prove that 1 is the identity. Prove that the inverse of a reduced word is also reduced.

**Proposition.** The triple  $(F(S), \cdot, 1)$  is a group.

*Proof.* It is left to show that  $\cdot$  is associative. □

**Proposition.** The group  $F(S)$  satisfies the definition of a free group.

*Proof.* □

**Problem 7.18.** Prove that  $F(S)$  is the unique group, up to group isomorphism, that satisfies the definition.

**Definition 7.12.** A *group presentation* is a pair  $(S, R)$ , where  $S$  is a set and  $R \subset F(S)$ . The group presented by  $(S, R)$  is defined to be  $\langle S \mid R \rangle = F(S)/N$ , where  $N$  is the smallest normal subgroup of  $F(S)$  containing  $R$ .

**Definition 7.13.** Let  $G$  be a group with the presentation  $\langle S \mid R \rangle$ , then the elements are called the *generators* of  $G$ . A group is said to be *cyclic* if  $|S| = 1$ .

**Example.** The presentation of a free group  $F(S)$  is  $\langle S \mid \emptyset \rangle$ . If  $R = \emptyset$ , we denote  $\langle S \mid R \rangle$  by  $\langle S \rangle$ . By our construction,  $\langle S \mid \emptyset \rangle = \{\prod s_i \mid s_i \in S^{\pm 1}\}$ .

Let  $G$  be a group and let  $S \subset G$ . If  $\langle S \rangle = G$ , then we say  $G$  is *generated by*  $S$ .

**Example.** The presentation of the trivial group is  $\langle a, b \mid abaa^{-1}b^{-2}, bab^{-1}a^{-2} \rangle$ .

**Problem 7.19.** Prove that a finite cyclic group of order  $n$  admits the presentation  $\langle a \mid a^n \rangle$ . We denote it by  $C_n$ .

**Problem 7.20.** Prove that  $\langle X \mid \emptyset \rangle$  is the smallest group that contains  $X$ .

**Definition 7.14.** Let  $G$  be a group. A *commutator* of  $G$  is an element of the form  $xyx^{-1}y^{-1}$ , where  $x, y \in G$ . Let  $G^c$  be the subgroup of  $G$  generated by all commutators in  $G$ . Here  $G^c$  is called the *commutator subgroup* of  $G$ .

**Problem 7.21.** Prove  $G^c$  is normal in  $G$ . Prove that  $G/G^c$  is abelian.

**Problem 7.22.** A *Tarski monster group* is a finitely generated infinite group where every proper non-trivial subgroup is cyclic of order  $p$ , a fixed prime. Prove that every Tarski monster group is simple.

**Problem 7.23.** A *Baumslag-Solitar group*  $BS(m, n)$  admits the presentation  $\langle a, b \mid ba^mb^{-1} = a^n \rangle$ . Prove that  $BS(1, 1) \approx C_2$ . This shows that the group presentation of a group is not necessarily unique.

**Problem 7.24.** Prove that  $\varphi : BS(2, 3) \rightarrow BS(2, 3)$  defined by  $\varphi(a) = a^2$  and  $\varphi(b) = b$  is a surjective group homomorphism.

Every group can be written as a group presentation. We will later show an equivalent statement: every group is a quotient group of a free group.

## 8 Rings

**Definition 8.1.** A triple  $(G, \circ, e)$  is a *group* if it is a monoid and for all  $g \in G$ , there exists  $h \in G$  such that  $h \circ g = e = g \circ h$ . Such  $h$  is called an *inverse* of  $g$ . If  $\circ$  is commutative, then the group is *abelian*.

**Definition 8.2.** A *ring* is an abelian group  $(R, +, 0)$  with the operation  $\cdot$  satisfies the following properties.

1. For all  $a, b, c \in R$ ,  $a \cdot (b \cdot c) = (a \cdot b) \cdot c$ . (associativity)
2. For all  $a, b, c \in R$ ,  $a \cdot (b + c) = a \cdot b + a \cdot c$  and  $(a + b) \cdot c = a \cdot c + b \cdot c$ . (distributivity)
3. There exists  $1_R \in R$  such that  $1_R \cdot a = a = a \cdot 1_R$  for all  $a \in R$ . We call  $1_R$  the *multiplicative identity* of  $R$ .

**Definition 8.3** (unit). An element  $u$  of a ring  $R$  is a unit if there exists  $v \in R$  such that  $u \cdot v = 1_R = v \cdot u$ .

**Remark.** A ring is a data  $(R, +, \cdot)$  consisting of a set  $R$ , with two binary operations  $+, \cdot$ .  $R$  is an abelian group under  $+$ . We use  $R^\times$  to represent units of  $R$ .

**Definition 8.4** (commutative ring). A ring  $R$  is commutative if and only if for all  $a, b \in R$ ,  $a \cdot b = b \cdot a$ .

**Example** (ring of functions). Let  $X$  be a set,  $R$  be a ring,  $R^X$  is the set of all functions from  $X$  to  $R$ .  $R^X$  is a ring. For any  $f, g \in R^X$ , addition and multiplication are defined as

$$\begin{aligned} + : \quad (f + g)(x) &= f(x) + g(x) \\ \cdot : \quad (f \cdot g)(x) &= f(x) \cdot g(x) \end{aligned}$$

for all  $x \in X$ .

**Definition 8.5** (polynomial ring). Let  $R$  be a commutative ring. Define  $R[x] = \{a_0 + a_1x + \dots + a_nx^n \mid n \geq 0, a_i \in R\}$ . Assuming  $n > m$ , we define addition and multiplication as follows:

$$\begin{aligned} + : \quad & \sum_{i=0}^n a_i x^i + \sum_{i=0}^m b_i x^i = \sum_{i=0}^n (a_i + b_i) x^i \\ \cdot : \quad & \left( \sum_{i=0}^n a_i x^i \right) \left( \sum_{j=0}^m b_j x^j \right) = \sum_{i=0}^{n+m} \left( \sum_{i+j=k} a_i b_j \right) x^k \end{aligned}$$

**Problem 8.1.** Prove that a polynomial ring is indeed a ring. What are 0 and 1 in  $R[x]$ ?

**Definition 8.6** (degree). The degree of  $0 \neq p(x) = a_0 + a_1x + \dots + a_nx^n \in R[x]$  is  $\deg(p) = \max\{n \mid a_n \neq 0\}$ . If  $p(x) = 0$ ,  $\deg(p) = -\infty$ .

**Definition 8.7** (field). A commutative (non-zero) ring  $F$  in which any non-zero element is a unit (i.e. for all  $a \in F$ , there exists  $b \in F$  such that  $a \cdot b = 1 = b \cdot a$ ).

**Lemma.** For any ring  $R$ , any  $a \in R$  satisfies  $a \cdot 0 = 0 = 0 \cdot a$ .

*Proof.*  $0 = 0 + 0$  implies  $a \cdot 0 = a \cdot (0 + 0) = a \cdot 0 + a \cdot 0$ . Since  $R$  is also a group, there exists an additive inverse  $-a \cdot 0 \in R$ . We can add  $-a \cdot 0$  to both sides:

$$a \cdot 0 - a \cdot 0 = (a \cdot 0 + a \cdot 0) - a \cdot 0 = a \cdot 0 + (a \cdot 0 - a \cdot 0) \iff 0 = a \cdot 0$$

Doing a symmetric argument to  $0 \cdot a = (0 + 0) \cdot a = 0 \cdot a + 0 \cdot a$ , we conclude that  $a \cdot 0 = 0 = 0 \cdot a$ . □

**Corollary.** If  $R$  is a ring and  $0 = 1$ , then  $R = \{0\}$ , we call  $\{0\}$  the *zero ring*.

*Proof.* For all  $a \in R$ ,  $a = a \cdot 1 = a \cdot 0 = 0$ . □

**Proposition.** Let  $R$  be a commutative ring. If there exist nonzero  $x, y \in R$  such that  $x \cdot y = 0$ , then  $x, y \notin R^\times$ .

*Proof.* Assume  $x$  is a unit. Then there exists some  $x^{-1} \in R$  and  $0 = x^{-1} \cdot 0 = x^{-1} \cdot (x \cdot y) = (x^{-1} \cdot x) \cdot y = 1 \cdot y = y$ . This contradicts the assumption that  $y \neq 0$ . Therefore,  $x$  is not a unit. By a similar argument, we can show  $y$  is also not a unit.  $\square$

**Lemma.** Let  $R$  be a commutative ring. For any  $f, g \in R[x]$ ,  $\deg(f \cdot g) \leq \deg(f) + \deg(g)$ .

*Proof.* If  $f$  or  $g$  is 0, then  $f \cdot g = 0$ . And  $\deg(f \cdot g)$  is  $-\infty$ .  $\deg(f) + \deg(g)$  is either  $-\infty$  or  $-\infty$  plus something. If  $f, g \neq 0$ , then let  $f = \sum_{i=0}^n a_i x^i$  and  $g = \sum_{j=0}^m b_j x^j$ . Their product would be  $\sum_{i=0}^{n+m} \left( \sum_{i+j=k} a_i b_j x^k \right)$ . If  $a_n, b_m \neq 0$ , then  $\deg(f \cdot g) = n + m$  which is equal to  $\deg(f) + \deg(g)$ . Otherwise  $\deg(f \cdot g) < n + m = \deg(f) + \deg(g)$ .  $\square$

**Remark.**  $0(x) = 0_R, \forall x \in X$  is the zero function. Constant function  $1(x) = 1_R, \forall x \in X$  is the unity.

**Definition 8.8** (subring). A subset  $S$  of a ring  $R$  is a subring if and only if

1.  $S$  is a subgroup of  $(R, +, 0)$ .
2. for all  $a, b \in S$ ,  $ab \in S$ . This means  $S$  is closed under multiplication.

**Definition 8.9** (ring homomorphism). Let  $R, R'$  be two rings, a map  $f : R \rightarrow R'$  is a ring homomorphism if and only if  $f$  preserves both  $+$  and  $\cdot$ :

$$f(a + b) = f(a) + f(b)$$

$$f(ab) = f(a)f(b)$$

for all  $a, b \in R$ .

**Definition 8.10** (unital homomorphism). A ring homomorphism  $f : R \rightarrow R'$  is unital if  $f(1_R) = 1_{R'}$ .

**Definition 8.11** (ideal). An ideal  $I$  in a ring is a subgroup of  $(R, +, 0)$  so that ①  $I \neq \emptyset$  and ② for all  $a, b \in I$ ,  $ab^{-1} = a - b \in I$ . Or equivalently, for any  $r \in R$  and  $i \in I$ ,  $ri \in I$ .

## 9 Real Numbers

Consider the function  $|\cdot| : \mathbb{Q} \rightarrow \mathbb{Q}$  defined by  $|a| = a$  if  $a \geq 0$  and  $|a| = -a$  if  $a < 0$ .

**Problem 9.1.** Use the definition of  $|\cdot|$ , prove that for all  $a, b \in \mathbb{Q}$ , the following conditions hold.

1.  $|a| \geq 0$ .
2. If  $|a| = 0$ , then  $a = 0$ .
3.  $|ab| = |a||b|$ .
4.  $|a + b| \leq |a| + |b|$ .

**Definition 9.1.** A *sequence* of rationals is a function  $a : \{n \in \mathbb{Z} \mid n \geq 1\} \rightarrow \mathbb{Q}$ . We denote a sequence by  $\{a_n\}_{n=1}^{\infty}$ .

**Remark.** We will use the notation  $\{a_n\}$  for a sequence  $\{a_n\}_{n=1}^{\infty}$ .

**Definition 9.2.** A sequence  $\{a_n\}$  *converges* to  $a \in \mathbb{Q}$  if for all  $r \in \mathbb{Q}^+$ , there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$ ,  $|a_n - a| < r$ . If a sequence  $\{a_n\}$  converges to  $a$ , then we write  $\{a_n\} \rightarrow a$ . Here  $a$  is called the *limit* of  $\{a_n\}$ .

**Proposition.** If a sequence converges to a limit, then this limit is unique.

*Proof.* Suppose  $\{a_n\} \rightarrow a$  and  $\{a_n\} \rightarrow b$ . Fix  $r = |a - b|/3$ , then for sufficiently large  $n$ , we have  $|a_n - a| < r$  and  $|a_n - b| < r$ . Then  $|a - b| \leq |a - a_n| + |a_n - b| < 2|a - b|/3$ , a contradiction.  $\square$

**Example.** Consider the sequence  $a_n = 1 + 1/n$ . We claim it converges to 1. Choose some  $r$ , and we want  $|a_n - 1| < r$  always hold. Notice that  $|a_n - 1| = |1/n| = 1/n$ . Let the integer part of  $1/r$  be  $c$ , so  $c \leq 1/r < c + 1$ , pick  $N = c + 1$ .

**Example.** Not every sequence of rationals has a limit. Consider  $\{3.1, 3.14, 3.141, 3.1415, \dots\} \rightarrow \pi \notin \mathbb{Q}$ .

**Problem 9.2.** Find the limits of the following sequences. If some limit does not exist, prove it.

1.  $a_n = 1/n$ .
2.  $a_n = \sum_{i=1}^n 2^{-i}$ .
3.  $a_n = (-1)^n$ .

**Definition 9.3.** A sequence  $\{a_n\}$  of rationals is *Cauchy* if for all  $r \in \mathbb{Q}^+$ , there exists  $N \in \mathbb{N}$  such that for all  $n, m \geq N$  and  $|a_n - a_m| < r$ .

**Proposition.** If  $\{a_n\}$  converges in  $\mathbb{Q}$ , then  $\{a_n\}$  is Cauchy.

*Proof.* Take some arbitrary  $r \in \mathbb{Q}^+$ . Since  $\{a_n\} \rightarrow a$ , there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$ ,  $|a_n - a| < r/2$  and for all  $m \geq N$ ,  $|a_m - a| < r/2$ . Then  $|a_n - a_m| = |(a_n - a) + (a - a_m)| \leq |a_n - a| + |a_m - a| < r$ . Therefore,  $\{a_n\}$  is Cauchy.  $\square$

**Problem 9.3.** Not every Cauchy sequence in  $\mathbb{Q}$  is bounded. Consider our previous example  $\{3.1, 3.14, 3.141, \dots\}$ , prove it is Cauchy but does not converge in  $\mathbb{Q}$ .

**Definition 9.4.** Let  $S \subset \mathbb{Q}$ . We say  $S$  is *bounded* if there exists  $L > 0$  such that  $|x| \leq L$  for all  $x \in S$ .

**Definition 9.5.** A rational sequence  $\{a_n\}$  is *bounded* if  $S = \{a_n \mid n \in \mathbb{N}\}$  is bounded in  $\mathbb{Q}$ .

**Proposition.** If  $\{a_n\}$  is a Cauchy sequence in  $\mathbb{Q}$ , then it is bounded in  $\mathbb{Q}$ .

*Proof.* Let  $r = 347$ . If the sequence  $\{a_n\}$  is Cauchy, then there exists  $N \in \mathbb{N}$  such that for all  $m, n \geq N$ ,  $|a_n - a_m| < 347$ . If  $m = N$ , then for all  $n \geq N$ ,  $|a_n - a_N| < 347$ . Since  $|a_n - a_N| < 347$ , replace  $|a_n - a_N|$  with 347. When  $n \geq N$ ,  $|a_n| < 347 + |a_N|$ . Let  $U = \max\{|a_1|, \dots, |a_{N-1}|, 347 + |a_N|\}$ . We have that  $|a_n| \leq U$  for all  $n \in \mathbb{N}$ . If  $n \geq N$ , then use  $347 + |a_N|$  as the bound. If  $n < N$ , then there finitely many values, and they are bounded by  $\max\{|a_1|, \dots, |a_{N-1}|\}$ .  $\square$

Let  $\{a_n\}$  and  $\{b_n\}$  be sequences. Define the addition as  $\{a_n\} + \{b_n\} = \{a_n + b_n\}$ . Define the multiplication as  $\{a_n\} \cdot \{b_n\} = \{a_n b_n\}$ .

**Proposition.** The set of all rational Cauchy sequence is closed under addition and multiplication.

*Proof.* Let  $\{a_n\}$  and  $\{b_n\}$  be Cauchy. Fix  $r$ . Since  $\{a_n\}$  is Cauchy, there exists  $N_1$  such that for all  $n, m \geq N_1$ ,  $|a_n - a_m| < r/2$ . Similarly, we have  $N_2$  for  $\{b_n\}$ . Let  $N = \max(N_1, N_2)$ . For all  $n, m \geq N$ ,  $|(a_m + b_m) - (a_n + b_n)| \leq |a_m - a_n| + |b_m - b_n| < r$ . Let  $\{a_k\}$  and  $\{b_k\}$  be Cauchy sequences in  $\mathbb{Q}$ . Then  $|a_n b_n - a_m b_m| \leq |a_n| |b_n - b_m| + |b_m| |a_n - a_m| \leq A |b_n - b_m| + B |a_n - a_m|$ , where  $A$  and  $B$  are upper bounds for sequences  $\{|a_k|\}$  and  $\{|b_k|\}$ . For all  $\varepsilon(2A)^{-1}, \varepsilon(2B)^{-1} > 0$ , there exists  $N \in \mathbb{N}$  such that for all  $m, n \geq N$ ,  $|a_m - a_n| < \varepsilon(2B)^{-1}$  and  $|b_m - b_n| < \varepsilon(2A)^{-1}$ . Now  $|a_m b_m - a_n b_n| < A\varepsilon(2A)^{-1} + B\varepsilon(2B)^{-1} = \varepsilon$ .  $\square$

**Problem 9.4.** Prove that the addition and multiplication satisfies commutativity, associativity, and distributivity.

Define the sequence 1 as  $a_n = 1$  and define the sequence 0 as  $a_n = 0$ .

**Problem 9.5.** Prove 1 and 0 are the identities of multiplication and addition, respectively. Moreover, prove that additive inverse of a Cauchy sequence is Cauchy.

We conclude that the set of all rational Cauchy sequences is a commutative ring with 1, denoted  $\mathcal{C}$ .

**Problem 9.6.** Prove that  $\mathcal{C}$  is not an integral domain.

**Proposition.** Let  $I := \{\text{Cauchy sequences that converge to } 0\}$ .  $I \subset \mathcal{C}$  is an ideal.

*proof sketch.* Let  $I$  be the set of sequences  $\{a_k\}$  in  $\mathcal{C}$  with the property that, given any rational  $r > 0$ , there exists an integer  $N$  such that if  $n \geq N$ , then  $|a_n| < r$ . For any  $c \in \mathcal{C}$  and  $i \in I$ , the sequence  $ci$  can be eventually scaled down to 0 because  $c$  is bounded. For any  $i' \in I$ , obviously  $i + i' \in I$ .  $\square$

**Proposition.** If  $\{a_k\} \in \mathcal{C} \setminus I$ , then there exist a positive rational number  $r$  and an integer  $N$  so that  $|a_k| \geq r$  for all  $n \geq N$ . In other words, if a Cauchy sequence  $\{a_k\}$  does not converge to 0, then after certain point  $n \geq N$  the sequence stays at least  $r$  distance away from 0.

*Proof.* If  $\{a_k\} \not\rightarrow 0$ , then there exists  $r \in \mathbb{Q}^+$  such that for all  $N \in \mathbb{N}$  there exists  $k \geq N$  such that  $|a_k| \geq 2r$  (the negation of converging to 0: there are some  $r$  that  $a_k$  is  $2r$  away from 0). Since  $\{a_k\}$  is Cauchy, for all  $n, m \geq N$  there exists  $N \in \mathbb{N}$  such that  $|a_n - a_m| < r$ .

$$\begin{aligned} \Rightarrow |a_m - a_n| < r & \quad (\text{property of absolute value}) \\ \Rightarrow |a_m| - |a_n| \leq |a_m - a_n| < r & \quad (\text{reverse triangle inequality}) \\ \Rightarrow -|a_n| < -|a_m| + r \Rightarrow |a_n| > |a_m| - r \end{aligned}$$

Now fix  $m \geq N$  and let  $|a_m| \geq 2r$ :  $|a_n| > |a_m| - r \geq 2r - r$ . This implies  $|a_n| > r$  and holds for all  $n \geq N$ .  $\square$

**Corollary.** if a Cauchy sequence does not converge to 0, all terms of the sequence eventually have the same sign.

*Proof.* If  $\{a_k\}$  is a Cauchy sequence that doesn't converge to 0, then  $\{a_k\} \notin I$ . By the above proposition, there exists some  $r > 0$  and  $N$  so that  $|a_k| \geq r$  for all  $n \geq N$ . Since  $\{a_k\}$  is Cauchy and  $r > 0$ , there exists  $M$  such that if  $m, n \geq M$ , then  $|a_n - a_m| < r$ . Assume without loss of generality that there is some  $k > \max(M, N)$  such that  $a_k$  is positive. Fix  $a_k$ , pick any  $n \geq k$  from the sequence:

- ① If  $a_n > a_k$ , then we are done ( $a_n$  is for sure positive).
- ② If  $a_n < a_k$ , then  $a_k - a_n < r$  which implies  $a_n$  is positive for all  $n$ .

$\square$

**Definition 9.6** (equivalence classes of Cauchy sequences). Let  $\{a_k\}$  and  $\{b_k\}$  be Cauchy sequences in  $\mathbb{Q}$ . We say that  $\{a_k\}$  is equivalent to  $\{b_k\}$ , denoted by  $\{a_k\} \sim \{b_k\}$ , if  $\{c_k\} = \{a_k - b_k\}$  is in  $I$ .

**Proposition.** The " $\sim$ " defined above gives an equivalence relation.

*Proof.* We need to check " $\sim$ " is reflexive, symmetric, and transitive.

Reflexive: For all  $\{a_k\}$ ,  $\{a_k\} - \{a_k\} = \{0\} \in I$ .

Symmetric: If  $\{a_k - b_k\} \in I$ , then for all  $\epsilon > 0$ , there exists some  $N \in \mathbb{N}$  such that for all  $n \geq N$ ,  $|a_n - b_n - 0| < \epsilon$ .

This means that  $|b_n - a_n - 0| < \epsilon$ , so  $\{b_k - a_k\} \in I$ .

Transitive: If  $\{a_k - b_k\} \in I$ , then for all  $\epsilon/2 > 0$ , there exists  $N_1 \in \mathbb{N}$  such that for all  $n \geq N_1$ ,  $|a_n - b_n - 0| < \epsilon/2$ .

If also  $\{b_k - c_k\} \in I$ , then for all  $\epsilon > 0$ , there exists  $N_2 \in \mathbb{N}$  such that for all  $n \geq N_2$ ,  $|b_n - c_n - 0| < \epsilon/2$ .

Let  $N = \max(N_1, N_2)$ . Then for all  $n > N$ ,  $|a_n - b_n - 0| < \epsilon/2$  and  $|b_n - c_n - 0| < \epsilon/2$ . It follows that  $|a_n - b_n - 0 + b_n - c_n - 0| = |a_n - c_n - 0| < \epsilon$ . So  $\{a_k - c_k\} \in I$ .

$\square$

**Definition 9.7** (set of equivalence classes). Define  $\mathbf{R} := \mathcal{C}/I$  as the set of all equivalence classes in  $\mathcal{C}$ .

**Proposition.** Term-wise addition and multiplication are well-defined on  $\mathbf{R}$ : If  $\{a_k\}$  is a Cauchy sequence, denote its equivalence class by  $[a_k]$ , then the sum and product of equivalence classes are defined as  $[a_k] + [b_k] = [a_k + b_k]$  and  $[a_k][b_k] = [a_k b_k]$ .

**Proposition.** Assume  $\mathbf{R} = \mathcal{C}/I$  is a commutative ring<sup>1</sup>. It turned out  $\mathbf{R}$  is a field.

*Proof.* We only need to show that multiplicative inverses exist for nonzero elements of  $\mathbf{R}$ . Consider some arbitrary  $[a_k] \neq I$ . By Proposition ???, since  $[a_k] \nrightarrow 0$ , the absolute value of  $[a_k]$  is eventually bounded below. In other words, there exists  $N \in \mathbb{N}$  and  $r > 0$ , such that  $|a_n| > r$  for all  $n \geq N$ . Therefore, we define  $[b_k]$  as follows:

$$\{b_k\} = \begin{cases} 1 & (k < N) \\ \frac{1}{a_k} & (k \geq N) \end{cases}$$

We include the  $k < N$  case since  $a_k$  may equal to 0 when  $k < N$ . Now show  $\{b_k\}$  is Cauchy. Take some arbitrary  $\epsilon > 0$ . Since  $\{a_k\}$  is Cauchy, there exists  $M$  such that for all  $n, m \geq M$ ,  $|a_n - a_m| < \epsilon r^2$ . Moreover,

$$|b_n - b_m| = \left| \frac{1}{a_n} - \frac{1}{a_m} \right| = \frac{|a_n - a_m|}{|a_n a_m|} \leq \frac{1}{r^2} |a_n - a_m| \leq \frac{1}{r^2} \epsilon r^2 = \epsilon$$

This implies that  $\{b_k\}$  is Cauchy. Furthermore,  $\{a_k\}\{b_k\} \rightarrow 1$ , which means that  $\{b_k\}$  is the multiplicative inverse of  $\{a_k\}$ . Therefore,  $\mathbf{R}$  is a field.  $\square$

**Definition 9.8** (order of  $\mathbf{R}$ ). Let  $a = [a_k], b = [b_k]$  be distinct elements of  $\mathbf{R}$ . We define  $a < b$  if  $a_k < b_k$  eventually and  $b < a$  if  $b_k < a_k$  eventually.

**Definition 9.9** (order axioms). Here are the order axioms

- O1** Trichotomy: Either  $a = b$ ,  $a < b$  or  $b < a$ ;
- O2** Transitivity: If  $a < b$  and  $b < c$ , then  $a < c$ ;
- O3** Addition Law:  $a < b$  if and only if  $a + c < b + c$ ;
- O4** Multiplication Law: If  $c > 0$ , then  $ac < bc$  if and only if  $a < b$ . If  $c < 0$ , then  $ac < bc$  if and only if  $b < a$ .

**Proposition.** The order relation on  $\mathbf{R}$  is well-defined and makes  $\mathbf{R}$  an ordered field.

*Proof.* Suppose  $\{a_k\} \sim \{a'_k\}$  and  $\{b_k\} \sim \{b'_k\}$ . We want to show  $a_k < b_k$  implies  $a'_k < b'_k$ . Since  $\{a_k\} \sim \{a'_k\}$  and  $\{b_k\} \sim \{b'_k\}$ , for all  $r > 0$ , there exists  $N_1$  such that for all  $n > N$ ,  $|a_n - a'_n| < r$ . This gives ①  $-r < a_n - a'_n < r$ . Similarly, for all  $r > 0$ , there exists  $N_2$  such that for all  $n > N$ ,  $|b_n - b'_n| < r$ . This gives ②  $r < b_n - b'_n < r$ . Notice that if we subtract inequality ① from ②, we get  $a'_k - b'_k < a_k - b_k$ . Since  $a_k < b_k$ ,  $a_k - b_k < 0$ . Therefore,  $a'_k < b'_k$ . Now we check all the order axioms:

- O1** Only one of  $a_k < b_k$ ,  $a_k = b_k$ , and  $a_k > b_k$  holds. So only one of  $[a_k] < [b_k]$ ,  $[a_k] = [b_k]$ , and  $[a_k] > [b_k]$  holds.
- O2** If  $[a_k] < [b_k]$  and  $[b_k] < [c_k]$ , then eventually  $a_k < b_k$  and  $b_k < c_k$ , so  $a_k < c_k$ . Finally,  $[a_k] < [c_k]$ .
- O3** If  $[a_k] < [b_k]$ , then eventually  $a_k < b_k$ , so  $a_k + c_k < b_k + c_k$ . Hence  $[a_k] + [c_k] < [b_k] + [c_k]$ .
- O4** If  $[a_k] < [b_k]$ , then eventually  $a_k < b_k$ , so  $a_k c_k < b_k c_k$  if  $c_k$ . Hence  $[a_k][c_k] < [b_k][c_k]$ .

$\square$

**Definition 9.10** (Archimedean). For all  $x \in F$  in an ordered field, there exists  $N \in \mathbb{Z}$  such that  $x < N$ .

**Proposition.**  $\mathbf{R}$  is an Archimedean ordered field.

*Proof.* Consider an arbitrary  $[a_k] \in \mathbf{R}$ . Since  $\{a_k\}$  is Cauchy, it is bounded in  $\mathbb{Q}$ . Furthermore,  $\mathbb{Q}$  is Archimedean. Therefore, there exists  $U \in \mathbb{Q}$  and  $N \in \mathbb{N}$ , such that for all  $n \in \mathbb{N}$ ,  $a_n \leq U < N$ . Therefore  $[a_k] < [N, N, N, \dots]$ , which means that  $\mathbf{R}$  is Archimedean.  $\square$

**Definition 9.11** (least upper bound). Let  $F$  be an ordered field, and let  $A$  be a nonempty subset of  $F$  that is bounded above. We say that  $L \in F$  is a least upper bound for  $A$  if the following two conditions hold:

1.  $L$  is an upper bound for  $A$ ;

<sup>1</sup>This is not hard to show. But for right now, we do not have the construction needed to prove this fact.



2. If  $M$  is any upper bound for  $A$ , then  $L \leq M$ .

**Definition 9.12** (least upper bound property). Let  $S \subseteq F$  be a nonempty set that is bounded above. Then  $S$  has a least upper bound in  $F$ .

**Proposition.** The least upper bound property holds in  $\mathbf{R}$ .

*proof sketch.* Let  $\emptyset \neq A \subseteq \mathbf{R} = \mathcal{C}/I$ , that is bounded above by  $m$ . Then there exists  $M \in \mathbb{Z}$  such that  $m \leq M$ . Since  $A \neq \emptyset$ , we can choose some  $a \in A$ . Hence there exists  $n \in \mathbb{Z}$  such that  $n < a$ . Consider

$$S_p = \{k2^{-p} \mid k \in \mathbb{Z}, n \leq k2^{-p} \leq M\}$$

Note that  $S_p$  is nonempty and finite. Let  $a_p = \min\{x \mid x \in S_p \text{ and } x \text{ is an upper bound of } A\}$ . If  $q > p$  implies  $a_q \leq a_p$ , then  $a_p - 2^{-p} < a_q$ . (Notice that  $a_p$  is not an upper bound of  $A$ , and  $a_q$  is an upper bound of  $A$ ). Therefore,  $|a_p - a_q| \leq 2^{-p}$ , for all  $p < q$ . In other words,  $\{a_p\}$  is Cauchy. So  $[a_p] = \text{lub}(A) \in \mathcal{C}/I = \mathbf{R}$ .  $\square$

**Definition 9.13** (real number). Real number is  $\mathbf{R}$ . Now denoted as  $\mathbb{R}$ .

**Proposition.** Real numbers are not countable.

*Proof.* Use Cantor's diagonal argument on real numbers.  $\square$

**Proposition.** Real numbers are Archimedean. That is for any  $a, b \in \mathbb{R}^+$ , there exists some  $n \in \mathbb{N}$  such that  $an > b$ .

*Proof.* If  $a > b$ , this is trivial. If  $a < b$ , assume for contradiction that we do not have an  $n \in \mathbb{N}$  such that  $an > b$ . This is the same as saying for all  $n \in \mathbb{N}$ ,  $an < b$ . Consider the set

$$S = \{an \mid n \in \mathbb{N}\}$$

Since all  $an < b$ ,  $S$  is bounded above by  $b$ . By the least upper bound property,  $S$  has a least upper bound denoted as  $L$ . Now consider  $L - a$ . Because  $a > 0$ ,  $L - a < L$ . Since  $L$  is the **least** upper bound,  $L - a$  is not an upper bound of  $S$ . In other words,  $L - a < an_0$ , for some  $n_0 \in \mathbb{N}$ . Therefore

$$L < an_0 + a \quad \text{and} \quad L < a(n_0 + 1)$$

Since  $n_0 \in \mathbb{N}$ ,  $n_0 + 1 \in \mathbb{N}$ . Therefore,  $a(n_0 + 1) \in S$ . But  $a(n_0 + 1) > L$  is a contradiction. We conclude that there must exist some  $n \in \mathbb{N}$  such that  $an > b$ .  $\square$

**Remark.** Ordered fields without the least upper bound property may not be Archimedean.

**Proposition.** The least upper bound is unique.

*Proof.* Suppose  $L_1$  and  $L_2$  are least upper bounds for  $\emptyset \neq A \subseteq F$ . By definition 6.1,  $L_1 \leq L_2$  and  $L_1 \geq L_2$ . By order axiom **O1**,  $L_1 = L_2$ .  $\square$

**Proposition.** Every real is between two consecutive integers. Formally, if  $a \in \mathbb{R}$ , there exists  $N \in \mathbb{Z}$  such that  $N - 1 \leq a \leq N$ .

*Proof.* Let  $S = \{n \in \mathbb{Z} \mid n > a\}$ . Then by the Archimedean property,  $S \neq \emptyset$  and  $S$  is bounded below. By the well-ordering principle,  $S$  has a least element. Then  $N - 1 \notin S$ , so  $N - 1 \leq a \leq N$ .  $\square$

**Theorem 9.1** (rationals are dense). Let  $a, b \in \mathbb{R}$ . If  $a < b$ , there exists some  $r \in \mathbb{Q}$  such that  $a < r < b$ .

*Proof.* Intuitively, if you multiply  $a$  and  $b$  by a large enough integer  $q$ , then the gap between them will be big enough to contain another integer,  $p$ . Let  $a, b$  be arbitrary real numbers such that  $a < b$ .



*Step 1:* We claim that there exists a  $q \in \mathbb{N}$  such that  $qb - qa > 1$ . Notice that  $qb - qa = q(b - a)$ . Since  $b - a > 0$  and they are both reals, by the Archimedean property, there exists  $q \in \mathbb{N}$  such that  $q(b - a) > 1$ .

*Step 2:* We claim that there exists a  $p \in \mathbb{Z}$ , such that  $qa < p < qb$ . Since every real is between two consecutive integers, we know that there exists  $p \in \mathbb{Z}$  such that  $p - 1 \leq qa < p$ . Since  $qb - qa > 1$ , we know that  $qb > qa + 1$ . And because  $p - 1 \leq qa$ , we know that  $p \leq qa + 1$ . This means that  $qa < p < qb$ , which is what we needed to show.

*Step 3:* Since  $qa < p < qb$ , if we divide the inequality by  $q$ , we get  $a < p/q < b$ . Moreover, since  $p, q$  are both integers,  $p/q \in \mathbb{Q}$ . This is all we need to show.  $\square$

**Corollary.** The irrationals are dense over the reals. In other words, if  $a < b \in \mathbb{R}$ , there exists  $r \in \mathbb{Q}^c$  such that  $a < r < b$ .

*Proof.* Let  $a, b$  be arbitrary real numbers such that  $a < b$ . Since  $\sqrt{2} > 0$ , we know that  $a/\sqrt{2} < b/\sqrt{2}$ . Because the rational numbers are dense, we know that there exists some  $p/q \in \mathbb{Q}$  such that  $a/\sqrt{2} < p/q < b/\sqrt{2}$ . Therefore,  $a < \sqrt{2}(p/q) < b$ . Because  $p/q$  is rational and  $\sqrt{2}$  is irrational, we know that their product  $\sqrt{2}(p/q)$  is irrational. Let  $r = \sqrt{2}(p/q)$ . We have  $a < r < b, r \in \mathbb{Q}^c$ , which is what we needed to show.  $\square$

**Proposition.** Let  $([a_n, b_n])_{n \in \mathbb{N}}$  be a nested sequence of closed bounded intervals in  $\mathbb{R}$ . That is, for any  $n$ , we have  $[a_{n+1}, b_{n+1}] \subseteq [a_n, b_n]$ , or equivalently,  $a_n \leq a_{n+1} \leq b_{n+1} \leq b_n$  for all  $n$ . Then  $\bigcap_{n \in \mathbb{N}} [a_n, b_n] \neq \emptyset$ .

*Proof.* Let  $A = \{a_n \mid n \in \mathbb{N}\}$ .  $A$  is bounded above by some  $b_1$ , which implies for every  $a_n \in A$ ,  $a_n \leq b_1$ . If  $a_m = \text{lub}(A)$ ,  $a_n \leq a_m \leq b_1$ .  $a_m \in [a_n, b_1]$  and  $a_m \in \bigcap_{n \in \mathbb{N}} [a_n, b_n]$ .  $\square$

**Remark.** What happens if we replace  $[a_n, b_n]$  with  $(a_n, b_n]$  and assume  $a_n < b_n$ ? Does the proposition still hold? No. An example would be  $a_n = 0$  and  $b_n = 1/n$  for all  $n \in \mathbb{N}$ . The intersection is empty (this is "clear" but can be proved using the Archimedean property).

## 10 Topological Spaces

**Definition 10.1.** A *metric* on a set  $X$  is a function  $d : X \times X \rightarrow [0, \infty)$  satisfying:

1.  $d(x, y) = 0$  if and only if  $x = y$  for all  $x, y \in X$ ;
2.  $d(x, y) = d(y, x)$  for all  $x, y \in X$ ;
3.  $d(x, z) \leq d(x, y) + d(y, z)$  for all  $x, y, z \in X$ .

If  $d$  is a metric on  $X$ , we say  $(X, d)$  is a *metric space*.

**Definition 10.2.** Let  $(X, d)$  be a metric space and let  $x_0 \in X$ . An *open ball* centered at  $x_0$  with radius  $r$ , where  $r \geq 0$ , is the set  $B(x_0, r) = \{x \mid x \in X, d(x, x_0) < r\}$ . An *open set* is a subset of  $X$  that can be written as a union of open balls.

**Definition 10.3.** A *topology* on a set  $X$  is a set  $\mathcal{T}$  of subsets of  $X$  such that:

1.  $\emptyset$  and  $X$  are in  $\mathcal{T}$ ;
2. the union of the elements of any subcollection of  $\mathcal{T}$  is in  $\mathcal{T}$ ;
3. the intersection of the elements of any finite subcollection of  $\mathcal{T}$  is in  $\mathcal{T}$ .

If  $\mathcal{T}$  is a topology on  $X$ , we say the pair  $(X, \mathcal{T})$  is a *topological space*.

It is trivial that  $\mathcal{T}$  must be a subcollection of  $\mathcal{P}(X)$ , so  $\mathcal{T}$  is a set.

**Example.** Let  $X$  be a set. The set  $\mathcal{P}(X)$  is a topology on  $X$ , called the *discrete topology*. The set  $\{X, \emptyset\}$  is also a topology on  $X$ , called the *trivial topology*.

**Definition 10.4.** Let  $\mathcal{T}$  be a topology on  $X$ . A subset  $U \subset X$  is said to be *open* if  $U \in \mathcal{T}$ . A subset  $V$  is said to be *closed* if  $X \setminus V$  is open.

**Definition 10.5.** Let  $\mathcal{T}$  and  $\mathcal{T}'$  be topologies on  $X$ . If  $\mathcal{T} \subset \mathcal{T}'$ ,  $\mathcal{T}$  is said to be *coarser* than  $\mathcal{T}'$ ; if  $\mathcal{T}' \subset \mathcal{T}$ ,  $\mathcal{T}$  is said to be *finer* than  $\mathcal{T}'$ .

**Proposition.** Every metric space is a topological space.

*Proof.* Denote the family of all open sets in  $X$  by  $\mathcal{T}$ . Fix  $x_0 \in X$ , then the open ball  $B(x_0, 0) = \emptyset$ . There exists an open set consisting of all open balls in  $X$ , which is  $X$ . Let  $U_1$  and  $U_2$  be open sets, we can express them as unions of open balls, and the union of open balls is again an open set, similarly, the finite intersection is an open set, so  $(X, d)$  induces a topology.  $\square$

**Problem 10.1.** Let  $(X, \mathcal{T}_1)$  and  $(X, \mathcal{T}_2)$  be two topological spaces. Are  $(X, \mathcal{T}_1 \cap \mathcal{T}_2)$  and  $(X, \mathcal{T}_1 \cup \mathcal{T}_2)$  topological spaces? Prove or state counterexamples.

**Remark.** Given a topological space  $(X, \mathcal{T})$ , we will denote it by  $X$  for convenience.

**Definition 10.6.** A function  $f : X \rightarrow Y$  between two topological spaces is *continuous* if  $f^{-1}(U)$  is open in  $X$  whenever  $U$  is open in  $Y$ . A continuous function with a left and right inverse is called a *homeomorphism*.

**Problem 10.2.** Let  $(X, \mathcal{T}_X)$  and  $(Y, \mathcal{T}_Y)$  be topological spaces. Let  $f : X \rightarrow Y$  be a function, prove that  $f$  is continuous if and only if for any closed set  $C \in Y$ ,  $f^{-1}(C)$  is closed in  $X$ .

**Problem 10.3.** Not every continuous bijection between topological spaces is a homeomorphism. Consider the identity function  $\text{id} : (\mathbb{R}, \mathcal{T}_1) \rightarrow (\mathbb{R}, \mathcal{T}_2)$ , where  $\mathcal{T}_1$  is the discrete topology and  $\mathcal{T}_2$  is the standard topology. Prove that  $\text{id}$  is not a continuous bijection but not a homeomorphism.

**Definition 10.7.** Let  $(X, \mathcal{T}_X)$  be a topological space and  $Y \subset X$ . The *subspace topology* on  $Y$  is  $\mathcal{T}_Y = \{U \cap Y \mid U \in \mathcal{T}_X\}$ .

**Problem 10.4.** Prove that a subspace topology is indeed a topology.

**Problem 10.5.** Let  $Y$  be a subspace of  $X$ . If  $U$  is open in  $Y$  and  $Y$  is open in  $X$ , then  $U$  is open in  $X$ .

**Proposition.** Let  $(X, \mathcal{T}_X)$  be a topological space and  $Y \subset X$ . The subspace topology on  $Y$  is the coarsest topology on  $Y$  for which the canonical inclusion  $i : Y \hookrightarrow X$  is continuous.

*Proof.* Given a topological space  $(X, \mathcal{T})$  and an arbitrary set  $Y$ , consider the topology on  $Y$  that makes  $f : Y \rightarrow X$  continuous. There exists such topology and the discrete topology is an example. Therefore, the intersection of topologies such that  $f : Y \rightarrow X$  is continuous is trivially the coarsest such topology, denoted  $\mathcal{T}_f$ . Now let  $Y \subset X$  and  $f = i$ , the inclusion map, then  $\mathcal{T}_i = \{i^{-1}(U) \mid U \in \mathcal{T}_X\}$ , where  $\mathcal{T}_X$  is the topology on  $X$  and  $i^{-1}(U) = U \cap Y$ , hence  $\mathcal{T}_i$  is the subspace topology.  $\square$

**Universal property for subspace topology.** Let  $(X, \mathcal{T}_X)$  be a topological space, let  $Y$  be a subset of  $X$ , and let  $i : Y \hookrightarrow X$  be the natural inclusion. For every topological space  $(Z, \mathcal{T}_Z)$  and every function  $f : Z \rightarrow Y$ ,  $f$  is continuous if and only if  $i \circ f : Z \rightarrow X$  is continuous.

$$\begin{array}{ccc} Z & \xrightarrow{f} & Y \\ & \searrow i \circ f & \downarrow i \\ & & X \end{array}$$

**Problem 10.6.** Prove that the composition of two continuous functions is continuous.

**Proposition.** Any topology satisfies this property is the subspace topology.

$$\begin{array}{ccc} \mathcal{T} & \xrightarrow{\text{id}_Y} & \mathcal{T} \\ & \searrow i & \downarrow i \\ & & \mathcal{T}_X \end{array} \qquad \begin{array}{ccc} \mathcal{T}_Y & \xrightarrow{\text{id}_Y} & \mathcal{T} \\ & \searrow i & \downarrow i \\ & & \mathcal{T}_X \end{array}$$

*Proof.* Let  $i \circ f$  be a continuous function. Since  $i$  is continuous, for all  $U \in \mathcal{T}_X$ , there exists  $V \in \mathcal{T}_Y$  and  $W \in \mathcal{T}_Z$  such that  $(i \circ f)^{-1}(U) = W$  and  $i^{-1}(U) = V$ , so  $f^{-1}(V) = W$ , so  $f$  is continuous. Let  $\mathcal{T}$  be any topology on  $Y$  with the property and let  $\mathcal{T}_Y$  be the subspace topology. Consider the left diagram, since  $\text{id}_Y$  is continuous,  $i \circ \text{id}_Y = i$  is continuous, then  $\mathcal{T}_Y \subset \mathcal{T}$ . Now consider the right diagram, since  $i$  is continuous,  $\text{id}_Y$  is continuous, then  $\mathcal{T} \subset \mathcal{T}_Y$ . Hence  $\mathcal{T} = \mathcal{T}_Y$ .  $\square$

**Definition 10.8.** Let  $(X, \mathcal{T}_X)$  be a topological space, let  $S$  be a set, and let  $\pi : X \twoheadrightarrow S$  be surjective. The *quotient topology* on  $S$  is the finest topology for which  $\pi$  is continuous, and  $\pi$  is called a *quotient map*.

**Example.** A *torus* is

We denote a torus by  $T^2$ .

## 11 Categorical Reformulations

Recall that *Set* does not have a zero object. We will take *Grp* as an example to show more categorical definitions and reformulate our notions of *Grp* and *Top*.

**Problem 11.1.** Prove that in *Grp*, the product of two groups  $G$  and  $H$  is the Cartesian product  $G \times H$ .

**Proposition.** In *Grp*, the trivial group  $1$  is the zero object.

*Proof.*

$\square$

**Definition 11.1.** Let  $\mathcal{C}$  be a category with zero object  $0$ . The *zero morphism*  $0_{A,B}$  between objects  $A$  and  $B$  is the unique morphism that factors through  $0$ .

Let  $G$  and  $H$  be groups. Since  $0$  is the zero object, there exists unique group homomorphisms  $\varphi : G \rightarrow 0$  and  $\psi : 0 \rightarrow H$ ,  $\psi \circ \varphi$  factors through the zero morphism  $0_{G,H}$  and it is the desired zero morphism.

**Definition 11.2.** Let  $\mathcal{C}$  be a category and let  $f : X \rightarrow Y$  be a morphism in  $\mathcal{C}$ . An object  $\ker(f)$  is said to be the *kernel* of  $f$  if for every object  $Z$  and  $h : Z \rightarrow X$  such that  $f \circ h = 0$ , where  $0$  is the zero morphism, there is a unique morphism  $\varphi : Z \rightarrow \ker(f)$  such that  $h = \varphi \circ \varphi$ .

**Problem 11.2.** Prove the set-theoretic definition of kernels coincide with the categorical definition of kernels.

**Proposition.** A group homomorphism is an epimorphism if and only if it is surjective.

*Proof.* ( $\Rightarrow$ ) Let  $f : H \rightarrow K$  be an epimorphism. Let  $X = K/f(H)$  be the set of right cosets of  $f(H)$  in  $K$ . Let  $\alpha$  not in  $X$ . Consider the set  $Y = X \cup \{\alpha\}$

( $\Leftarrow$ ) Let  $f : G \rightarrow H$  be surjective, so  $f$  is an epimorphism in  $\mathbf{Set}$ , since  $\mathbf{Grp}$  is a subcategory of  $\mathbf{Set}$  and  $f$  is a group homomorphism,  $f$  is a group epimorphism.  $\square$

**Universal property for quotient groups.** Let  $G$  be a group and let  $N \trianglelefteq G$ . Let  $\pi : G \rightarrow G/N$  be the quotient epimorphism. For all group homomorphism  $\varphi : G \rightarrow H$  with  $N \subset \ker(\varphi)$ , there exists a unique group homomorphism  $\bar{\varphi} : G/N \rightarrow H$  such that  $\bar{\varphi} \circ \pi = \varphi$ .

$$\begin{array}{ccc} G/N & \xleftarrow{\pi} & G \\ & \searrow \bar{\varphi} & \downarrow \varphi \\ & & H \end{array}$$

**Problem 11.3.** Prove that the quotient groups satisfy the universal property and the universal property characterizes quotient groups.

**Proposition.** Every group can be written as a quotient group of a free group.

*Proof.*  $\square$

**Universal property for quotient topology.** Let  $(X, \mathcal{T}_X)$  be a topological space, let  $S$  be a set, and let  $\pi : X \rightarrow S$  be surjective. For every topological space  $(Z, \mathcal{T}_Z)$  and every function  $f : S \rightarrow Z$ ,  $f$  is continuous if and only if  $f \circ \pi : X \rightarrow Z$  is continuous.

$$\begin{array}{ccc} X & \xrightarrow{\pi} & S \\ & \searrow f \circ \pi & \downarrow f \\ & & Z \end{array}$$

**Problem 11.4.** Prove that the quotient topology satisfies the universal property. Prove that any topology satisfies this property is the quotient topology.

## References

- [1] Awodey, S. *Introduction to Category Theory*. 2nd ed. Oxford, Oxford University Press. 2008.
- [2] Bradley, Tai-Danae, et al. *Topology*. 1st ed. MIT Press. 2020.
- [3] Jech, T. *Set Theory*. 3rd ed. Berlin ; London, Springer. 2011.
- [4] Kaplansky, I. *Set Theory and Metric Spaces*. 1st ed. Chelsea Publishing Co., New York. 1977.
- [5] Lang, S. *Algebra*. 3rd ed. New York, Springer. 2002.
- [6] Tao, T. *Analysis I*. 3rd ed. Singapore Springer Singapore, Imprint: Springer. 2016.

## Alphabetical Index

- abelian, 23, 27
- addition, 14, 15
- alternating group, 25
- antecedent, 2
- automorphism, 24
- automorphism group, 24
- axiom, 1
- axiom of choice, 13
- axiom of empty set, 4
- axiom of extensionality, 4
- axiom of infinity, 6
- axiom of pairing, 5
- axiom of power set, 7
- axiom of regularity, 6
- axiom of union, 6
- axiom schema of replacement, 12
- axiom schema of separation, 5
  
- basis, 25
- Baumslag-Solitar group, 26
- biconditional proposition, 3
- bijective, 10
- binary operation, 7
- bounded, 29
- Burali-Forti paradox, 13
  
- canonical decomposition of
  - functions, 21
- Cantor's theorem, 16
- Cantor-Schröder-Bernstein theorem, 16
- cardinality, 10
- Cartesian product, 7
- category, 19
- Cauchy, 29
- Cayley's theorem, 24
- choice function, 13
- class, 12
- closed, 35
- coarser, 35
- codomain, 9, 19
- commutative, 9
- commutative ring, 27
- commutator, 26
- commutator subgroup, 26
- complement, 6
- complete finite induction, 9
- composition, 9, 23
- composition operation, 19
- conditional proposition, 2
- conjunction, 2
- consequent, 2
- continuous, 35
- contrapositive, 3
- converges, 29
- converse, 3
- coproduct, 22
- countable, 16
- cyclic, 26
  
- De Morgan's law, 6
- degree, 27
- dihedral group, 23
- discrete topology, 35
- disjoint, 6
- disjoint union, 7
- disjunction, 2
- division, 15
- domain, 3, 7, 19
- dual category, 19
  
- elements, 4
- empty set, 4
- empty word, 25
- epimorphism, 20
- equivalence class, 7
- equivalence relation, 7
- Euclid's theorem, 15
- even function, 10
- even permutation, 23
- existential quantifier, 3
  
- field, 27
- finer, 35
- finite, 10
- finite induction, 8
- free group, 25
- full, 19
- function, 9
  
- fundamental theorem of
  - arithmetic, 15
  
- generalized associativity, 24
- generated by, 26
- generators, 25, 26
- group, 23, 27
- group homomorphism, 24
- group presentation, 26
  
- homeomorphism, 35
  
- ideal, 28
- identity, 23
- identity function, 11
- identity morphism, 19
- image, 9
- index, 24
- inductive set, 6
- infimum, 8
- infinite, 10
- initial object, 20
- initial segment, 8
- injective, 10
- integers, 14
- intersection, 6
- invariant, 11
- inverse, 11, 14, 23, 25, 27
- involution, 23
- isomorphic, 20
- isomorphism, 20
  
- kernel, 24, 37
  
- lattice of subgroups, 24
- left coset, 24
- left inverse, 11, 23
- length, 25
- letters, 25
- limit, 29
- limit ordinal, 13
- linearly ordered, 8
- locally small category, 19
- logically equivalent, 2
- lower bound, 8

- metric, 35
- metric space, 35
- monoid, 23
- monoid homomorphism, 24
- monomorphism, 20
- morphisms, 19
- multiplication, 14, 15
- multiplicative identity, 27
  
- n-tuple, 5
- natural numbers, 7
- negation, 2
- normal subgroup, 25
  
- objects, 19
- odd function, 10
- odd permutation, 23
- open, 35
- open ball, 35
- open set, 35
- order, 23
- order-preserving, 11
- order-type, 11
- ordered pair, 5
- ordinal, 12
  
- partial ordering, 8
- partially ordered set, 8
- partition, 6
- permutation, 23
- polynomial ring, 27
- power set, 7
- preimage, 9
- prime, 14
  
- product, 21
- proper subset, 5
- proposition, 1
- propositional function, 3
  
- quotient group, 25, 37
- quotient map, 36
- quotient set, 11, 20
- quotient topology, 36
  
- range, 7
- rational numbers, 15
- recursion, 14
- reduced, 25
- restriction, 11
- right inverse, 11
- ring, 27
- ring homomorphism, 28
  
- sequence, 29
- set, 4
- simple, 25
- singleton, 5
- subcategory, 19
- subgroup, 24
- subring, 28
- subset, 5
- subspace topology, 35
- subtraction, 15
- successor, 6
- successor ordinal, 12
- supremum, 8
- surjective, 10
- symmetric difference, 7
  
- symmetric group, 23
  
- Tarski monster group, 26
- terminal object, 20
- theorem, 1
- topological space, 35
- topology, 35
- torus, 36
- transfinite induction, 13
- transitive, 12
- trivial group, 23
- trivial topology, 35
- truth table, 2
- truth value, 1
  
- uncountable, 16
- union, 6
- unit, 27
- unital homomorphism, 28
- universal quantifier, 3
- upper bound, 8
  
- vacuous truth, 2
- variable, 3
  
- well-ordered, 8
- well-ordering principle, 13
- well-ordering theorem, 8
- word, 25
  
- zero morphism, 37
- zero object, 20
- zero ring, 27