

Observable time traces (100-step resolution)
HB: bond count from hb_obs.dat | Dist: min(6 pair distances) from dist_obs.dat
RT thresholds — HB: hb ≤ 0.0 / hb ≥ 6.0 | Dist: mindist ≤ 2.0 / mindist ≥ 12.0
Colour = seed identity (tab10). Alpha = σ_naive. Legend shows round-trip count per seed.
pulling — HB
bond count
steps (×10⁶)
wB_s4 (1RT)
wC_s3 (1RT)
wC_s5 (1RT)
wC_s7 (1RT)
wC_s9 (1RT)
noclamps — HB
bond count
steps (×10⁶)
wC_s7 (1RT)
wC_s9 (2RT)
pulling — dist
min pair dist (oxDNA units)
steps (×10⁶)
wA_s2 (2RT)
wA_s3 (1RT)
wA_s4 (1RT)
wA_s6 (1RT)
wA_s7 (1RT)
wA_s8 (1RT)
wA_s9 (1RT)
wB_s1 (2RT)
wB_s3 (1RT)
wB_s4 (1RT)
wB_s7 (1RT)
wB_s8 (1RT)
wC_s1 (3RT)
wC_s10 (2RT)
wC_s2 (2RT)
wC_s3 (2RT)
wC_s4 (3RT)
wC_s5 (1RT)
wC_s6 (1RT)
wC_s7 (3RT)
wC_s8 (3RT)
wC_s9 (5RT)
noclamps — dist
min pair dist (oxDNA units)
steps (×10⁶)
wB_s5 (1RT)
wC_s6 (1RT)
pushing — dist
min pair dist (oxDNA units)
steps (×10⁶)
wB_s1 (3RT)
wB_s2 (1RT)
wB_s9 (1RT)