

Reaction-diffusion spatial modeling of COVID-19 in Chicago

Trent Gerew*

Department of Applied Mathematics, Illinois Institute of Technology, Chicago, Illinois

November 2, 2021




Figure 1: Timeline of the progression of COVID-19 in Chicago with key public policy events marked. The COVID-19 data was obtained from the City of Chicago Data Portal [3]. The dates of the policy events were gathered from the Illinois.gov press releases [8], [6], [7], [9], [5], the Chicago Tribune [2], and NBC Chicago [1].

1 Model Setup

We estimate the non-COVID death rate μ using the average United States death rate in urban regions from 2019 [4]. That is, $\mu = 1.8997 \times 10^{-5}$ [per day].

*tgerew@hawk.iit.edu

Table 1: Time sequence of events and simulation times.

Initial simulation time	Imposed lockdown	Effective lockdown	Last fitting day
March 18, 2020	March 21, 2020	April 1, 2020	June 24, 2020
$t_i = 1$	$t_q = 4$	$t_q = 15$	$t_f = 99$




Figure 2: Schematic diagram of the model. The dashed lines indicate the interaction of the infected populations with the susceptible populations that leads to infection.

$$S_t = \mathfrak{D}_S \Delta S - \beta_{SA} S A - \beta_{SI} S I - \mu S, \quad (1)$$

$$E_t = \mathfrak{D}_E \Delta E + \beta_{SA} S A + \beta_{SI} S I - (\sigma_A + \sigma_I) E, \quad (2)$$

$$AR_t = M_{AR} A, \quad (3)$$

$$A_t = \mathfrak{D}_A \Delta A + \sigma_A E - M_{AR} A, \quad (4)$$

$$I_t = \sigma_I E - M I, \quad (5)$$

$$H_t = \gamma M I - (1 - \omega) \chi H - \omega \psi H, \quad (6)$$

$$R_t = (1 - \gamma) M I + (1 - \omega) \chi H, \quad (7)$$

$$D_t = \omega \psi H. \quad (8)$$

Table 2: Population values for Chicago. Initial populations are determined from March 13, 2020.

	Population	
Total population	N	2,695,598
Initial infected	I_0	162
Initial hospitalized	H_0	38
Initial deceased	D_0	3

2 ODE Dynamics

We want to understand the trajectories of the dynamics of the ODE system under different initial conditions. To do this we first find the equilibrium points by solving

$$S_t = E_t = A_t = I_t = H_t = R_t = D_t = 0$$

Table 3: Fitting parameters for Chicago: optimal (best-fitting), median and interquartile range, and variation range used in the optimization algorithm. Initial parameter guesses were uniformly sampled within these ranges.

	Median (interquartile range)	Initial value
Transmission rate, $S \rightarrow I$ [per day]	β_{SI}	$c \in \mathcal{U}[0, 1]$
Transmission rate, $S \rightarrow A$ [per day]	β_{SA}	$c \in \mathcal{U}[0, 1]$
Lockdown effect, $S \rightarrow I$	η_{SI}	$c \in \mathcal{U}[0, 1]$
Lockdown effect, $S \rightarrow A$	η_{SA}	$c \in \mathcal{U}[0, 1]$
Incubation period, $E \rightarrow I$ [days]	$1/\sigma_I$	$1/k, k \in \mathcal{U}[2, 7]$
Latent period, $E \rightarrow A$ [days]	$1/\sigma_A$	$1/k, k \in \mathcal{U}[2, 7]$
Infectivity period [days]	$1/M$	$1/k, k \in \mathcal{U}[5, 12]$
Recovery period, $A \rightarrow AR$ [days]	$1/M_{AR}$	$1/k, k \in \mathcal{U}[5, 12]$
Recovery period, $H \rightarrow R$ [days]	$1/\chi$	$1/k, k \in \mathcal{U}[5, 20]$
Period to deceased, $H \rightarrow D$ [days]	$1/\psi$	$1/k, k \in \mathcal{U}[5, 20]$
Conversion fraction ($I \xrightarrow{\gamma} H, I \xrightarrow{1-\gamma} R$)	γ	$c \in \mathcal{U}[0.25, 0.75]$
Conversion fraction ($H \xrightarrow{\omega} D, H \xrightarrow{1-\omega} R$)	ω	$c \in \mathcal{U}[0.1, 0.5]$
Initial population fraction, exposed	E_0/I_0	$c \in \mathcal{U}[1, 5]$
Initial population fraction, asymptomatic	A_0/I_0	$c \in \mathcal{U}[1, 5]$

simultaneously for $\mathbf{x} = (S, E, A, AR, I, H, R, D)$. The solutions of this system are of the form $\mathbf{x}^* = (0, 0, 0, AR, 0, 0, R, D)$. This implies there are infinitely many non-isolated equilibrium points. We determine the stability of these equilibrium points by analyzing the linearized system near the points. The Jacobian of the system is

$$J = \begin{pmatrix} -A\beta_{SA} - I\beta_{SI} - \mu & 0 & -S\beta_{SA} & 0 & -S\beta_{SI} & 0 & 0 & 0 \\ A\beta_{SA} + I\beta_{SI} & -\sigma_A - \sigma_I & S\beta_{SA} & 0 & S\beta_{SI} & 0 & 0 & 0 \\ 0 & \sigma_A & -M_{AR} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & M_{AR} & 0 & 0 & 0 & 0 & 0 \\ 0 & \sigma_I & 0 & 0 & -M & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & M\gamma & -\chi(1-\omega) - \psi\omega & 0 & 0 \\ 0 & 0 & 0 & 0 & M(1-\gamma) & \chi(1-\omega) & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \psi\omega & 0 & 0 \end{pmatrix} \quad (9)$$

Now evaluating J at the equilibrium point \mathbf{x}^* and calculating the eigenvalues, we have

$$\lambda = \{0, 0, 0, -M, -M_{AR}, -\mu, -\sigma_A - \sigma_I, -\chi + \chi\omega - \psi\omega\}. \quad (10)$$

Note that the first three eigenvalues are 0, which implies the equilibrium points are non-isolated. This agrees with our earlier observation.

The equilibrium points are stable when $\lambda_i < 0$ for $4 \leq i \leq 8$. Since all the system parameters are positive, this implies $\lambda_i < 0$ for $4 \leq i \leq 7$. Thus the stability depends on the sign of λ_8 . There are two cases when $\lambda_8 = -\chi + \chi\omega - \psi\omega < 0$ is true:

1. $0 < \omega \leq 1$ implies $\lambda_8 < 0$, and
2. $\omega > 1$ and $\chi < \frac{\psi\omega}{\omega-1}$ implies $\lambda_8 < 0$.

That is, whenever we have either of these conditions the equilibrium points are stable. We call this situation *endemic*. If $\lambda_8 > 0$, the equilibrium points are unstable and the situation is an *epidemic*.

References

- [1] Read the full ‘restore Illinois’ plan aimed at reopening the state during coronavirus. *NBC Chicago*, May 2020. <https://www.nbcchicago.com/news/coronavirus/read-the-full-restore-illinois-plan-aimed-at-reopening-the-state-during-coronavirus/2267039/>.
- [2] J. Byrne, D. Petrella, and A. Lukach. Mayor Lori Lightfoot says Chicago will move to phase 3 of her reopening plan on June 3 but warns: ‘COVID-19 is still very much part of our present’. *Chicago Tribune*, May 2020. <https://www.chicagotribune.com/coronavirus/ct-coronavirus-chicago-lightfoot-reopening-20200528-cefwuidwnfd7a57m25uavq6me-story.html>.
- [3] City of Chicago. Daily chicago covid-19 cases, deaths, and hospitalizations, 2021. Data retrieved from Chicago Data Portal, <https://data.cityofchicago.org/Health-Human-Services/Daily-Chicago-COVID-19-Cases-Deaths-and-Hospitaliz/kxzd-kd6a>.
- [4] S. C. Curtin and M. R. Spencer. Trends in death rates in urban and rural areas: United States, 1999-2019. *National Center for Health Statistics*, NCHS Data Brief, no 417, September 2021. <https://www.cdc.gov/nchs/products/databriefs/db417.htm>.
- [5] Illinois.gov. Gov. Pritzker releases guidelines to safely reopen additional businesses and industries as state advances to next phase of restore Illinois. Press Release, June 2020. <https://www.illinois.gov/news/press-release.21714.html>.
- [6] Illinois.gov. Gov. Pritzker aligns Illinois mask guidance with CDC for fully vaccinated people. Press Release, May 2021. <https://www.illinois.gov/news/press-release.23322.html>.
- [7] Illinois.gov. Gov. Pritzker announces metrics-based pathway for Illinois to fully reopen; expands vaccine eligibility to all residents 16+ on April 12. Press Release, March 2021. <https://www.illinois.gov/news/press-release.22961.html>.
- [8] Illinois.gov. Gov. Pritzker issues guidelines for Illinois reopening on June 11. Press Release, June 2021. <https://www.illinois.gov/news/press-release.23399.html>18.
- [9] Illinois.gov. United Center vaccination appointments open Thursday for Illinois seniors. Press Release, March 2021. <https://www.illinois.gov/news/press-release.22868.html>.