

ROUGEOLE

1. Case definition

- + Fever > 38°C
- + Generalized maculopapular rash lasting 3 days or more
- + At least **one of the following symptoms**:
 - cough
 - cold or respiratory infection
 - conjunctivitis

2. Case management

*** Systematically look for danger signs:**

- weight loss > 10% of initial weight or weight < 3th percentile
- polypnoea or draught
- severe dehydration
- consciousness disorders

If any of these signs are present, the child should be kept under observation (or referred in accordance with the SPT discharge); if refused, treatment should be started and the child asked to return the following day.

*** Processing :**

- Amoxicillin: 80 mg/kg/day in 2 doses (see FT dosage for children) for 7 days. If the child cannot be monitored, give Benzathine benzyl penicillin instead.
60,000U/kg in 1 single injection.
- Vitamin A: 100,000 IU if < 12 months, 200,000 IU if > 12 months
- Tetracycline ophthalmic ointment 1% 4x/day
- Paracetamol 30 mg/kg/d in 3 doses
- If T° > 38.5, give malaria treatment according to FT Palu

If **signs of dehydration**: see SPT DRHe and FT Rehydration/Cholera

If **convulsions** : Diazepam injection 0.5 mg/kg IR or IM, consult SPT Convulsion

3. Action to be taken

Notification:

In a register (specify age, place of residence, vaccination status) and in the monthly SNIS report with information on age and vaccination status. Record the number of new cases and deaths each week, and report by telephone to the District Health Office (BDS).

Take a sample and fill in the sampling form for the national laboratory of the Institut National de Santé Publique -INSP. The sampling form must be approved by the BDS before the sample is sent to the laboratory.

Epidemic detection

Analyze data before sending:

- count the cases
- check whether they are more numerous than in the same week of the previous non-epidemic year
- verify the existence or otherwise of local outbreaks.

Any anomalies must be reported to the hierarchy without delay, together with the data.

***Epidemic : vaccination
(see next page)***

an epidemic: vaccination

Where	Who
- Around the outbreak - In high-risk areas (peri-urban concentrations, camps for displaced persons or refugees, nomadic populations) - In the health center to nosocomial transmission.	- Children aged 6 months to 5 years. - Children vaccinated before age of 11 months, if possible (second dose).

Strengthen routine immunization in areas not reached by EPI