

INFORMATION - EPILEPSY

Does the person convulsive epileptic seizures?

1. Symptoms :

- Loss of consciousness
- Stiffness more than 1-2 minutes
- Convulsions lasting 1-2 minutes
- Tongue bite or wound
- Loss urine and/or stools
- After the attack: fatigue, drowsiness, sleepiness, confusion, abnormal behaviour, headaches or muscle pain

2. Disease education: explaining

What is a seizure/epilepsy:

- A seizure is caused by a problem in the brain.
- Epilepsy is a disease characterized by repeated seizures.
- It is not contagious and is not transmitted by spirits or witchcraft.

How a crisis unfolds and possible causes:

- It is a chronic condition
- Seizures can be controlled in 2/3 of cases by medication
- The patient can then lead a normal life
- He not need medication for the rest of his life.

3. Case management

• Measures to be taken during seizures

Do not immobilize the subject

Keep away anything that could obstruct the person or cause injury Do not insert anything into the person's mouth

Loosen your tie or collar, as appropriate

Once the seizure is over, place a cushion or folded piece of clothing under his head and gently turn his head to one side.

Make they can breathe easily and let them rest.

- **At CDS and then at home**, administration of a single anticonvulsant drug (1 ampoule of Diazepam 10 mg IM at CDS, then phenobarbital cp 100 mg: 1cp/dr at home) is enough to stop seizures or reduce their number and intensity.
- **Refer to district hospital** if seizure lasts longer than 5 minutes, new seizure occurs immediately after first seizure and loss of consciousness persists for several minutes after seizure ends