



I. Practical help for screening children under 5 for malnutrition

1. Active community screening: based on BP measurement and oedema testing



2. Passive screening in health facilities :



Use the table for boys and girls (highest weight regardless of sex)													
Size lying	Weight Kg - Z-score						Size lying	Weight Kg - Z-score					
cm	-4.0	-3	-2	-1.5	-1	0	cm	-4.0	-3	-2	-1.5	-1	0
For children with a lying height of less than 87 cm													
45	1.75	1.90	2.07	2.16	2.25	2.46	66	5.5	5.92	6.4	6.65	6.92	7.50
45.5	1.81	1.97	2.14	2.23	2.33	2.55	66.5	5.6	6.02	6.5	6.75	7.03	7.62
46	1.88	2.03	2.21	2.30	2.41	2.63	67	5.7	6.11	6.6	6.86	7.14	7.74
46.5	1.94	2.10	2.28	2.38	2.48	2.72	67.5	5.8	6.2	6.69	6.96	7.24	7.85
47	2.00	2.16	2.35	2.45	2.56	2.80	68	5.8	6.29	6.79	7.06	7.35	7.97
47.5	2.06	2.23	2.42	2.53	2.64	2.89	68.5	5.9	6.38	6.89	7.16	7.45	8.1
48	2.12	2.30	2.50	2.61	2.72	2.97	69	6.0	6.47	6.99	7.26	7.56	8.2
48.5	2.18	2.37	2.57	2.68	2.80	3.06	69.5	6.1	6.56	7.08	7.36	7.66	8.3
49	2.25	2.44	2.65	2.76	2.89	3.16	70	6.2	6.65	7.18	7.46	7.77	8.4
49.5	2.32	2.51	2.73	2.85	2.97	3.25	70.5	6.3	6.74	7.27	7.56	7.87	8.5
50	2.39	2.59	2.81	2.94	3.07	3.35	71	6.3	6.82	7.37	7.66	7.97	8.6
50.5	2.46	2.67	2.90	3.03	3.16	3.46	71.5	6.4	6.91	7.46	7.76	8.1	8.8
51	2.54	2.75	2.99	3.12	3.26	3.56	72	6.5	7.00	7.55	7.86	8.2	8.9
51.5	2.62	2.84	3.08	3.22	3.36	3.68	72.5	6.6	7.08	7.65	7.95	8.3	9.0
52	2.70	2.93	3.18	3.32	3.47	3.79	73	6.7	7.16	7.74	8.0	8.4	9.1
52.5	2.79	3.02	3.28	3.42	3.58	3.91	73.5	6.7	7.25	7.83	8.1	8.5	9.2
53	2.88	3.12	3.38	3.53	3.69	4.03	74	6.8	7.33	7.91	8.2	8.6	9.3
53.5	2.98	3.22	3.49	3.64	3.80	4.16	74.5	6.9	7.41	8.0	8.3	8.7	9.4
54	3.08	3.33	3.61	3.76	3.92	4.29	75	6.9	7.49	8.1	8.4	8.8	9.5
54.5	3.18	3.44	3.73	3.88	4.05	4.42	75.5	7.0	7.56	8.2	8.5	8.8	9.6
55	3.29	3.55	3.85	4.01	4.18	4.55	76	7.1	7.64	8.3	8.6	8.9	9.7
55.5	3.39	3.67	3.97	4.14	4.31	4.69	76.5	7.2	7.72	8.3	8.7	9.0	9.8
56	3.50	3.78	4.10	4.26	4.44	4.83	77	7.2	7.79	8.4	8.8	9.1	9.9
56.5	3.61	3.90	4.22	4.40	4.58	4.98	77.5	7.3	7.87	8.5	8.8	9.2	10.0
57	3.7	4.02	4.35	4.53	4.71	5.13	78	7.4	7.94	8.6	8.9	9.3	10.1
57.5	3.8	4.13	4.47	4.66	4.85	5.27	78.5	7.4	8	8.7	9.0	9.4	10.2
58	3.9	4.25	4.6	4.79	4.99	5.42	79	7.5	8.1	8.7	9.1	9.5	10.3
58.5	4.1	4.37	4.72	4.92	5.12	5.56	79.5	7.6	8.2	8.8	9.2	9.5	10.4
59	4.2	4.49	4.85	5.05	5.25	5.71	80	7.6	8.2	8.9	9.2	9.6	10.4
59.5	4.3	4.6	4.97	5.17	5.39	5.85	80.5	7.7	8.3	9.0	9.3	9.7	10.5
60	4.4	4.71	5.09	5.3	5.52	5.99	81	7.8	8.4	9.1	9.4	9.8	10.6
60.5	4.5	4.82	5.21	5.42	5.65	6.13	81.5	7.8	8.5	9.1	9.5	9.9	10.7
61	4.6	4.93	5.33	5.54	5.77	6.26	82	7.9	8.5	9.2	9.6	10.0	10.8
61.5	4.7	5.04	5.44	5.66	5.89	6.40	82.5	8.0	8.6	9.3	9.7	10.1	10.9

62	4.8	5.14	5.56	5.78	6.01	6.53	83	8.1	8.7	9.4	9.8	10.2	11.0
62.5	4.9	5.25	5.67	5.89	6.13	6.65	83.5	8.2	8.8	9.5	9.9	10.3	11.2
63	5.0	5.35	5.77	6.00	6.25	6.78	84	8.3	8.9	9.6	10.0	10.4	11.3
63.5	5.1	5.45	5.88	6.12	6.36	6.9	84.5	8.3	9	9.7	10.1	10.5	11.4
64	5.1	5.54	5.99	6.23	6.48	7.03	85	8.4	9.1	9.8	10.2	10.6	11.5
64.5	5.2	5.64	6.09	6.33	6.59	7.15	85.5	8.5	9.2	9.9	10.3	10.7	11.6
65	5.3	5.74	6.19	6.44	6.7	7.27	86	8.6	9.3	10.0	10.4	10.8	11.7
65.5	5.4	5.83	6.29	6.55	6.81	7.39	86.5	8.7	9.4	10.1	10.5	11.0	11.9


Use the table for boys and girls (highest weight regardless of sex)													
Size stand	Weight Kg - Z-score						Size stand	Weight Kg - Z-score					
cm	-4.0	-3	-2	-1.5	-1	0	cm	-4.0	-3	-2	-1.5	-1	0
For children with a standing height of 87cm and over													
87	9.0	9.6	10.4	10.8	11.2	12.2	104	12.0	13.0	14.0	14.6	15.2	16.5
87.5	9.0	9.7	10.5	10.9	11.3	12.3	104.5	12.1	13.1	14.2	14.7	15.4	16.7
88	9.1	9.8	10.6	11.0	11.5	12.4	105	12.2	13.2	14.3	14.9	15.5	16.8
88.5	9.2	9.9	10.7	11.1	11.6	12.5	105.5	12.3	13.3	14.4	15.0	15.6	17.0
89	9.3	10.0	10.8	11.2	11.7	12.6	106	12.4	13.4	14.5	15.1	15.8	17.2
89.5	9.4	10.1	10.9	11.3	11.8	12.8	106.5	12.5	13.5	14.7	15.3	15.9	17.3
90	9.5	10.2	11.0	11.5	11.9	12.9	107	12.6	13.7	14.8	15.4	16.1	17.5
90.5	9.6	10.3	11.1	11.6	12.0	13.0	107.5	12.7	13.8	14.9	15.6	16.2	17.7
91	9.7	10.4	11.2	11.7	12.1	13.1	108	12.8	13.9	15.1	15.7	16.4	17.8
91.5	9.8	10.5	11.3	11.8	12.2	13.2	108.5	13.0	14.0	15.2	15.8	16.5	18.0
92	9.9	10.6	11.4	11.9	12.3	13.4	109	13.1	14.1	15.3	16.0	16.7	18.2
92.5	9.9	10.7	11.5	12.0	12.4	13.5	109.5	13.2	14.3	15.5	16.1	16.8	18.4
93	10.0	10.8	11.6	12.1	12.6	13.6	110	13.3	14.4	15.6	16.3	17.0	18.6
93.5	10.1	10.9	11.7	12.2	12.7	13.7	110.5	13.4	14.5	15.8	16.4	17.1	18.8
94	10.2	11.0	11.8	12.3	12.8	13.8	111	13.5	14.6	15.9	16.6	17.3	19.0
94.5	10.3	11.1	11.9	12.4	12.9	13.9	111.5	13.6	14.8	16.0	16.7	17.5	19.2
95	10.4	11.1	12.0	12.5	13.0	14.1	112	13.7	14.9	16.2	16.9	17.7	19.4
95.5	10.4	11.2	12.1	12.6	13.1	14.2	112.5	13.9	15.0	16.3	17.1	17.9	19.6
96	10.5	11.3	12.2	12.7	13.2	14.3	113	14.0	15.2	16.5	17.2	18.0	19.8
96.5	10.6	11.4	12.3	12.8	13.3	14.4	113.5	14.1	15.3	16.7	17.4	18.2	20.0
97	10.7	11.5	12.4	12.9	13.4	14.6	114	14.2	15.4	16.8	17.6	18.4	20.2
97.5	10.8	11.6	12.5	13.0	13.6	14.7	114.5	14.3	15.6	17.0	17.8	18.6	20.5
98	10.9	11.7	12.6	13.1	13.7	14.8	115	14.5	15.7	17.2	18.0	18.8	20.7
98.5	11.0	11.8	12.8	13.3	13.8	14.9	115.5	14.6	15.9	17.3	18.1	19.0	20.9
99	11.1	11.9	12.9	13.4	13.9	15.1	116	14.7	16.0	17.5	18.3	19.2	21.1
99.5	11.2	12.0	13.0	13.5	14.0	15.2	116.5	14.8	16.2	17.7	18.5	19.4	21.3
100	11.2	12.1	13.1	13.6	14.2	15.4	117	15.0	16.3	17.8	18.7	19.6	21.5

100.5	11.3	12.2	13.2	13.7	14.3	15.5		117.5	15.1	16.5	18.0	18.9	19.8	21.7
101	11.4	12.3	13.3	13.9	14.4	15.6		118	15.3	16.6	18.2	19.0	19.9	22.0
101.5	11.5	12.4	13.4	14.0	14.5	15.8		118.5	15.4	16.8	18.4	19.2	20.1	22.2
102	11.6	12.5	13.6	14.1	14.7	15.9		119	15.5	16.9	18.5	19.4	20.3	22.4
102.5	11.7	12.6	13.7	14.2	14.8	16.1		119.5	15.7	17.1	18.7	19.6	20.5	22.6
103	11.8	12.8	13.8	14.4	14.9	16.2		120	15.8	17.3	18.9	19.8	20.7	22.8
103.5	11.9	12.9	13.9	14.5	15.1	16.4								

3. Look for bilateral oedema: apply pressure for 3 seconds, release and see if the thumbprint forms a depression (cup sign).

Severity of edema	Coding
Mild oedema: both feet	+
Moderate oedema: of the 2 feet and the lower part of the 2 forearms Intermediate between mild and severe edema severe	++
Severe oedema: generalized or including both feet, legs, arms and face	+++

II. Summary of acute malnutrition management services and admission criteria (children under 5)



III. Indicators and definition of categories of acute malnutrition

Classification of acute malnutrition

Indicators	Age	Malnutrition moderate acute	Malnutrition malnourished
Nutritional edema	All	No	Yes
P/T2008	Children	≥ -3 to < -2 Z-score	< -3 Z-score
P/T WHO	Teenagers	≥ 70 à $< 80\%$	$< 80\%$
BMI	Adults	≥ 16 à < 17	< 16 with recent weight loss
PB	Children over 6 months	≥ 115 to < 125 mm	< 115 mm
	Adults	≥ 160 to < 180 mm	< 160 mm with recent weight loss

IV. Management of severe acute malnutrition in ATS

MANAGEMENT OF SEVERE ACUTE MALNUTRITION WITHOUT MEDICAL COMPLICATIONS STA

IDENTIFICATION



Weight gain

Take PB



Look for edema



PATIENT CIRCUIT

1. Home
2. Anthropometry
3. Sorting
4. Admission to the program



Measure standing height (≥87cm) or lying height (<87cm)



SURVEILLANCE

For each visit :

- appetite test
- PB measurement
- weight gain
- search for edema
- complete clinical examination
- promoting essential family practices

For patients admitted on P/T:

- Measure size on admission
- Use target weight

TREATMENT FAILURE

Criteria for non-response to treatment	Time elapsed after admission
Stable weight (non-children) edematous)	21 days
Weight loss since admission in the program (non-edematous children)	14 days
No signs of melting edema	14 days
Edema still present	21 days
Failed appetite test	Any visit
Weight loss of 5% of body weight (non-edematous children)	Any visit
Weight loss for two consecutive visits	Any visit
Unsatisfactory weight gain after total melting of oedema (kwashiorkor) or from day 14 'marasmus' (<2.5g/kg/d)	Any visit

ADMISSION CRITERIA

Children from 6 months to 120cm or 2 to 3 kg
 Nutritional edema+ or++ OR
 -P/T<-3 z-scores OR
 -PB<115 mm
 Adolescents: P/T<70% or nutritional edema
 Adults: BMI<16 or BP<180 with recent weight loss or bilateral edema

AND

-Good to moderate appetite
 -No medical complications

NUTRITIONAL THERAPY

Weight class	ATPE sachets (92g)	
	Per day	Per week
3,0 - 3,4	1 ¼	8
3,5 - 4,9	1 ½	10
5,0 - 6,9	2	15
7,0 - 9,9	3	20
10,0 - 14,9	4	30
15,0 - 19,9	5	35
20,0 and more	6	40

APPETIT TEST

Weight class	Proportion a 30gATPE			Quantity ATPE in grams		
	Low	Medium	Good	Low	Medium	Good
Less than 4 kg	<1/8s	1/8 - 1/4s	>1/4s	<15gr	15 - 25gr	>25gr
4 - 6,9	<1/4s	1/4 - 1/3s	>1/3s	<25gr	25 - 30gr	>35gr
7 - 9,9	<1/3s	1/3 - 1/2s	>1/2s	<35gr	35 - 50gr	>50gr
10 - 14,9	<1/2s	1/2 - 3/4s	>3/4s	<50gr	50 - 75gr	>75gr
15 - 19,9	<3/4s	3/4 - 1s	>1s	<100gr	100 - 150gr	>150gr
Over 20 kg	<1s	>1s		<150gr	>150gr	

SYSTEMATIC PROCESSING

-Amoxicillin on admission and for 7 days
 <5kg 125mg * 2 20-35kg 750mg * 2
 5-10kg 250mg * 2 >35kg 1000mg * 2
 10-20kg 500mg * 2

-Vitamin A: at 4th visit EXCEPT if received within the last 6 months

*6-8 Kg: 1 cap of 100,000 IU;
 *8 Kg: 1 cap of 200,000 IU

-Deworming at 4th visit IF child aged 12 months and over
 *Either Mebendazole ≥ 12 months 1 tablet 500mg
 . Either Albendazole 400mg
 *12 to 23 months: ½ tablet
 *≥ 24 months: 1 tablet

-Measles at 4th visit for children ≥ 9 months without vaccination
 Anti-malarial treatment according to national protocol on admission

OUTPUT

"Healed

- If admitted on P/T: P/T ≥ -1.5 z-scores after 2 consecutive visits
- If admitted on PB: PB ≥ 125mm after 2 consecutive visits
- Patients with nutritional oedema: discharge at least 14 days after oedema has completely melted away

"deceased

"Abandonment confirmed / not confirmed

"Transferred to ESS if appetite test fails - complications

"Non-responsive" / Refusal to transfer to SST/Medical referral

SOME KEY MESSAGES ABOUT ATPE

1. RUTF is a food and medicine given to children to treat severe acute malnutrition. It must not be shared with family members or consumed by non-malnourished people.
2. Breastfeeding should be continued and encouraged until the child is two years of age or older. Always give the child breast milk before offering him/her RUTF.
3. Always give the child drinking water when taking the RUTF.

For more details, consult the national protocol, December 2018

If blater oedemas +++ OR marasmo-kwashi (P/T<-3Z-score and presence of oedemas) AND/OR medical complications
 AND/OR Appetite low, organize the transfer internal to the SST

