

REHYDRATION / CHOLERA

1. Definition of cholera cases

- Patients over 5 years of age who develop severe dehydration or die as a result acute watery diarrhea.
- Sudden increase in the number of patients with acute watery diarrhea, especially those with the "rice water" stools typical of cholera.

2. General management diarrhoeal dehydration

- * **Step 1:** Assess the level of dehydration
- * **Step 2:** rehydrate the patient according to the level of dehydration and monitor frequently, then reassess the patient's condition.

PLAN C.

SEVERE DEHYDRATION: INTRAVENOUS REHYDRATION (see FT PCIME)

Intravenous administration lactated Ringer's solution (or saline).

* Patients one year or more :

100 ml/kg for 3 hours, i.e. 30 ml/kg very quickly in 30 minutes, then the rest in 2 hours 30 minutes

PLAN CAN BE REPEATED - If there is no improvement - REFER

* Patients under one year of age :

100 ml/kg for 6 hours, i.e. 30 ml/kg for the first hour and the rest for 5 hours

If no improvement - REFER

Reminder: 1 ml of solute = 20 drops

PLAN B.

MODERATE DEHYDRATION: ORAL REHYDRATION QUANTITY OF SRO TO BE ADMINISTERED DURING THE FIRST FOUR HOURS :

Age	< 4 months	4-14 months	12-13 months	2-4 years	5-14 years	> 15 years
Weight	< 5 kg	5-9 kg	9-10 kg	11-15 kg	16-29 kg	> 30 kg
ORS in ml	200-400	400-600	600-800	800-1200	1200-2200	2200-4000

PLAN A.

- * **MAINTAIN REHYDRATION**
- * **COMPENSATE FOR FLUID LOSS THROUGH STOOL WITH SRO**

- * **Step 3:** feed the patient, continue breastfeeding the child.

3. Specific measures to be taken for cholera

Administer an oral antibiotic to the patient:

- * Doxycycline cp100mg :300mg **in adults** as a single dose, or
Tetracycline cp 250mg 3x 2cp/d
- * Cotrimoxazole 400/80mg: 50mg/kg/day in 2 doses, for 3 days **in children**.

Withdrawals

As soon as the first case of cholera is suspected, a fresh stool sample or rectal swab should be collected and sent to the laboratory for confirmation of the diagnosis:

- label the sample and fill out a laboratory
- transport to the laboratory :
- place samples in a clean container if they can reach the laboratory in less than two hours, or soak the stool liquid in blotting paper and send it to the laboratory in a sealed plastic bag, or place the sample in alkaline peptone water if it can reach the laboratory in less than 24 hours.

The fight against cholera

- * Disinfection/excreta disposal
- * Water bleaching
- * Careful hand washing treating patients
- * Selective prophylaxis in contact subjects: Doxycycline 100mg 1cp in one dose.