

CHOKING - HEIMLICH MANEUVER

Measures for CLEARING UPPER RESPIRATORY TRACKS

5 Slaps on the back

if no success

HEIMLICH MANEUVER



Classification of choking according to severity:

1) Mild airway obstruction

The patient can still answer questions about the cause of the blockage, he breathes and coughs.

Treatment: Encourage coughing, no other intervention

2) Severe airway obstruction


Patient cannot speak, cannot breathe; wheezes, tries to cough or may be unconscious

Treatment: lean forward - guide the patient to stand upright and lean forward.

A maximum of five slaps on the back with the inside of the wrist between the shoulder blades, with the aim of releasing the obstruction with the slap. After each slap, check that the airway is clear.

(*Treatment cont.*) Remove/clean foreign bodies, such as vomit, from the oral cavity. **Perform 5 compressions on stomach area (Heimlich maneuver) in the reclined position (patient sitting or standing).**


If the procedure is unsuccessful, repeat the 5 back slaps.



HEIMLICH MANEUVER

1. Patient standing or sitting (Heimlich maneuver)

- Stand behind the victim, place your arms under his and wrap around his waist;
- Then place your fist, palm down, in the epigastric fossa, with your other free hand grasping your wrist. Make a sudden upward and backward movement. Repeat immediately if the foreign body has not been vomited or spat out by the patient(s) or child.



2. Unconscious patient :

Sit astride the victim's thighs. Place the heel of one hand on the epigastric cavity (NOT the ribs), covering it with the other free hand. Apply sudden upward and backward pressure, with arms outstretched. Repeat several times until the foreign body or object is vomited or spat out.

