

RECEPTION A VICTIM OF SEXUAL VIOLENCE.

1. Definition.

Sexual violence refers to sexual acts imposed by physical violence, coercion or any form intimidation: threats, emotional blackmail, abuse of power or trust, use of weapons or in situations where the person was not fully capable of consenting (minors, infirm, etc.).

2. Treatment in the event of rape

The first interview must be carried out in good conditions, with great tact, care and kindness, explaining in advance all the examinations that will be necessary (for children, demonstration on a doll). Minors must be accompanied by a parent or guardian.

questioning:

❖ note down the facts, grievances (advise the victim to write them down +++) if possible, otherwise use the victim's exact words and expressions.

Physical examination :

- ❖ Examination of the entire body for wounds, bruises, erosions, burns, bruises, hematomas, fractures;
- ❖ detailed examination of genitalia, perineum, anus, mouth

Paraclinical examinations

- ❖ vaginal, vulval, anal and buccal swabs for spermatozoa testing
- ❖ pregnancy test for women of childbearing age
- ❖ HIV, VDRL, TPHA, hepatitis B: Ag Hbs, Ac anti-HBc, Chlamydia serological tests, with an invitation to repeat them after 3 months

Treatments to be administered

If woman of childbearing age :

1. Taking the pregnancy test:

- If positive, allow pregnancy to progress
- If negative, give emergency contraception:

☐ Levonorgestrel (Postinor) 0.75mg, 2cp: 1 tablet every 12 hours or Levonorgestrel (Postinor) 1, 5 mg single dose (first choice)

☐ Microgynon 30 (COC) 4 tablets , then 4 tablets 12 hours later (second choice)

☐ Microlut (low-dose COP) 25 tablets immediately and 25 tablets 12 hours later (third choice).

☐ IUD (fourth choice); if the woman presents 3 to 7 days after the rape incident, the IUD is the best choice.

only method to use.

Taking a pregnancy test after 4 weeks

STI prophylaxis :

☑ Azithromycin 1 gram+ Cefixime 400mg +Metronidazole 2g in one dose

☑ Or Doxycycline 100 mg x 2/d or 200 mg once/d for 7 days+ Benzathine penicillin 2.4 M IU single dose, half a dose in each buttock + Ceftriaxone 250 mg IM single dose + Metronidazole 2 g (second choice)

Tetanus prophylaxis

Tetanus prophylaxis for unvaccinated subjects, Td1 (first dose on first contact), Td2 after 1 month, Td3 after 6 months, Td4 after one year and Td5 after 1 to 3 years. If the subject has been vaccinated and the last vaccination was more than 10 years ago, give a booster.

Hepatitis B prevention :

Hepatitis B vaccination (Engerix B): M0 (within 2 4 h), M1, M2 and M12. Give passive immunization with immunoglobulin if available.

HIV prophylaxis :

As PEP is an emergency, it should be started as soon as possible, at best within the first few minutes, and in any case before 48 hours. The duration of prophylaxis is four weeks; the recommended regimen is :

- ✓ TDF/3TC /DTG from age 10, 1 tablet per day
- ✓ ABC/ 3TC +DTG between 6 and 10 years of age,
- ✓ ABC/3TC+LPV/r for children under 6.

3. Care levels

➤ **At community level.**

Rape victims need :

- ✓ to be believed, to be listened to sympathetically
- ✓ to be protected, put out of danger
- ✓ to be understood, not to be judged,
- ✓ to be recognized as victims, and for the perpetrators to be denounced and punished
- ✓ to be supported, helped and informed about all the steps to take, about the law, about all their rights,
- ✓ to be supported and enhanced.

➤ **CS level:**

- Take care of the victim (local care if injured, psychological care),
- prevent STIs,
- give emergency contraception if there is a risk of pregnancy)
- encourage victims to lodge complaints and transfer hospital so that a doctor can draw up a certificate attesting to the violence.

➤ **HD, HR, HN:**

In addition to what is done at CDS,

- draw up a medical certificate of violence

- detect STIs

- encourage victims to lodge complaints

Screening for HIV and hepatitis is offered at the first consultation: depending on the results, treatment may need to be adapted or repeated 3 months later, due to the incubation period for STIs.