

# Iterative methods for linear systems

Seidel iteration, Successive over-relaxation

## Seidel iteration

$$\mathbf{Ax} = \mathbf{b} \quad \longrightarrow \quad \mathbf{A} = \mathbf{L} + \mathbf{D} + \mathbf{U}$$

lower    diagonal    upper

Rewrite

$$\mathbf{D}\mathbf{x} + (\mathbf{L} + \mathbf{U})\mathbf{x} = \mathbf{b}$$

# Seidel's and Jacobi iteration

- ▶ Jacobi iteration:  $\mathbf{D}\mathbf{x}^{(n+1)} + (\mathbf{L} + \mathbf{U})\mathbf{x}^{(n)} = \mathbf{b}$

- ▶ Seidel iteration:  $\mathbf{D}\mathbf{x}^{(n+1)} + \mathbf{L}\mathbf{x}^{(n+1)} + \mathbf{U}\mathbf{x}^{(n)} = \mathbf{b}$

## Seidel's iteration in components

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1m} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2m} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3m} \\ \vdots & \vdots & \vdots & \ddots & \vdots \end{pmatrix}$$

$$a_{11}x_1^{(n+1)} + a_{12}x_2^{(n)} + a_{13}x_3^{(n)} + \cdots + a_{1m}x_m^{(n)} = b_1$$

$$a_{22}x_2^{(n+1)} + a_{21}x_1^{(n+1)} + a_{23}x_3^{(n)} + \cdots + a_{2m}x_m^{(n)} = b_2$$

...

At each iteration, sweep down the system of equations; at each step use  $\mathbf{x}^{(n+1)}$ -s from previous steps: for  $x_i^{(n+1)}$  use  $x_1^{(n+1)} \cdots x_{i-1}^{(n+1)}$  (Jacobi iteration uses  $x_1^{(n)} \cdots x_{i-1}^{(n)}$ )

# Seidel iterations: Sufficient conditions for convergence

Write the original system as

$$\mathbf{x} = \mathbf{B}\mathbf{x} + \mathbf{c}$$

with  $\mathbf{B} = -\mathbf{D}^{-1}(\mathbf{L} + \mathbf{U})$

## Theorem $S_1$

Let  $\|\mathbf{B}\|_1 < 1$  or  $\|\mathbf{B}\|_\infty < 1$ .

Then, Seidel's iterations converge for any initial  $\mathbf{x}^{(0)}$  with the rate of a geometric series,

$$\|\mathbf{x}^{(n)} - \hat{\mathbf{x}}\| \leq q^n \|\mathbf{x}^{(0)} - \hat{\mathbf{x}}\|,$$

with  $q < \|\mathbf{B}\|$

## A weaker convergence condition

Define  $\mathbf{B}_L = -\mathbf{D}^{-1}\mathbf{L}$  and  $\mathbf{B}_U = -\mathbf{D}^{-1}\mathbf{U}$ .

### Theorem $S_2$

Let  $\|\mathbf{B}_U\| + \|\mathbf{B}_L\| < 1$ . Then, Seidel's iterations converge for any initial  $\mathbf{x}^{(0)}$ , and

$$\|\mathbf{x}^{(n)} - \hat{\mathbf{x}}\| \leq q^n \|\mathbf{x}^{(0)} - \hat{\mathbf{x}}\|,$$

with

$$q = \frac{\|\mathbf{B}_U\|}{1 - \|\mathbf{B}_L\|}$$

# Symmetric positive definite matrices

In practice, we often have symmetric positive definite  $\mathbf{A}$ -s

## Theorem $S_3$

Let  $\mathbf{A}$  is symmetric positive definite. Then for any initial  $\mathbf{x}^{(0)}$ , Seidel's iteration converges with the rate of a geometric series.

**Note:** No conditions on the norm of  $\mathbf{A}$  or  $\mathbf{B}$

# Necessary and sufficient conditions for convergence

Rewrite Seidel's algorithm as a simple iteration:

$$\mathbf{x}^{(n+1)} = \tilde{\mathbf{B}}\mathbf{x}^{(n)} + \tilde{\mathbf{c}}$$

with

$$\tilde{\mathbf{B}} = (\mathbf{D} + \mathbf{L})^{-1}\mathbf{U}$$

## Theorem $S_4$

Seidel's iterations converge for any initial  $\mathbf{x}^{(0)}$  if and only if

$$\rho(\tilde{\mathbf{B}}) < 1$$



# Jacobi vs Seidel

Loosely speaking,

- ▶ Jacobi method is suitable for  $\mathbf{A} \approx \text{diagonal}$ .
- ▶ Seidel's method is suitable for  $\mathbf{A} \approx \text{triangular (or symmetric)}$ .

# Successive over-relaxation method (SOR)

At step  $j$  of iteration  $n + 1$ ,

1. Do a Seidel's iteration step: compute  $\tilde{x}_j^{(n+1)}$
2. Shift  $x_i^{(n+1)} = \omega \tilde{x}_i^{(n+1)} + (1 - \omega)x_i^{(n)}$

Here  $\omega$  is an arbitrary parameter  $\omega \in [0, 2]$ . Tune  $\omega$  to speed up convergence.

In the matrix form

$$\text{Seidel iteration:} \quad \mathbf{x}^{(n+1)} = -\mathbf{D}^{-1}\mathbf{L}\mathbf{x}^{(n+1)} - \mathbf{D}^{-1}\mathbf{U}\mathbf{x}^{(n)} + \mathbf{c}$$

$$\begin{aligned} \text{SOR iteration:} \quad \mathbf{x}^{(n+1)} &= (1 - \omega)\mathbf{x}^{(n)} \\ &+ \omega \left( -\mathbf{D}^{-1}\mathbf{L}\mathbf{x}^{(n+1)} - \mathbf{D}^{-1}\mathbf{U}\mathbf{x}^{(n)} + \mathbf{c} \right) \end{aligned}$$

## Further variations of SOR

- ▶ Use  $j$ -dependent  $\omega_j$
- ▶ Symmetric successive over-relaxation
- ▶ ...