

2D Physical World Report

F06 Team 06

Dicson Wijaya (1002289), Wenkie Lau (1002219),
Mok Jun Neng (1002219), Charlotte Phang (1002277),
Martin Tan (1002173)

April 13, 2017




Figure 1

A #how the experiment is conducted, to find λ_{algae}

#how to find the theoretical λ_{algae} value is a range

#experimental λ_{algae} should be within theoretical λ_{algae}

B (a) The equation of control volume in the heat exchanger

$$\frac{dE_{cv}}{dt} = \dot{Q}_{cv} - \dot{W}_{cv} + \dot{m}_i \left(h_i + \frac{V_i^2}{2} + gz_i \right) + \dot{m}_e \left(h_e + \frac{V_e^2}{2} + gz_e \right)$$

*Assume steady state, so $\dot{m}_i = \dot{m}_e = \dot{m}$ No change in potential and kinetic energy
Air is ideal gas.*

$$\frac{dE_{cv}}{dt} = \dot{Q}_{cv} - \dot{W}_{cv} + \dot{m} (h_1 - h_2)$$

(b) The rate of heat entering control volume

$$\dot{Q}_{cv} = \dot{Q}_{exchanged}$$

$$\dot{Q}_{cv} = m_a c_a \left(\frac{dT_a}{dt} \right) - \dot{Q}_{ambient}$$

$$\dot{Q}_{cv} = m_a c_a \left(\frac{dT_a}{dt} \right) - \lambda_{algaebottle} (T_a - T_{amb})$$

By substituting value of $\lambda_{algaebottle} = 1.76$

$$\dot{Q}_{cv} = m_a c_a \left(\frac{dT_a}{dt} \right) - 1.76 (T_a - T_{amb})$$

(c) The equation of control volume in the heat exchanger

From simplified $\frac{dE_{cv}}{dt}$ equation in the previous part

$$\frac{dE_{cv}}{dt} = \dot{Q}_{cv} - \dot{W}_{cv} + \dot{m} (h_1 - h_2)$$

(d) Express $\dot{W}_{cv} = f(T_a)$ assuming \dot{W}_{cv} is a constant Assume that $\frac{dE_{cv}}{dt} = 0$

$$0 = m_a c_a \left(\frac{dT_a}{dt} \right) - \lambda_{algaebottle} (T_a - T_{amb}) - \dot{W}_{cv} + \dot{m} [c_w (T_i - T_e)]$$

$$\lambda_{algaebottle} (T_a - T_{amb}) + \dot{W}_{cv} - \dot{m} [c_w (T_i - T_e)] = m_a c_a \left(\frac{dT_a}{dt} \right)$$

$$\int 1 dt = \int \frac{m_a c_a}{\lambda_{algaebottle} (T_a - T_{amb}) + \dot{W}_{cv} - \dot{m} [c_w (T_i - T_e)]} dT_a$$

$$t + c = \frac{m_a c_a}{\lambda_{algaebottle}} \ln(\lambda_{algaebottle} (T_a - T_{amb}) + \dot{W}_{cv} - \dot{m} [c_w (T_i - T_e)])$$

$$\frac{\lambda_{algaebottle} (t + c)}{m_a c_a} = \ln(\lambda_{algaebottle} (T_a - T_{amb}) + \dot{W}_{cv} - \dot{m} [c_w (T_i - T_e)])$$

$$e^{\frac{\lambda_{algaebottle} (t+c)}{m_a c_a}} = \lambda_{algaebottle} (T_a - T_{amb}) + \dot{W}_{cv} - \dot{m} [c_w (T_i - T_e)]$$

$$\dot{W}_{cv} = e^{\frac{\lambda_{algaebottle} (t+c)}{m_a c_a}} - \lambda_{algaebottle} (T_a - T_{amb}) + \dot{m} [c_w (T_i - T_e)]$$

C Plot \dot{W}_{cv} vs t

To find the value of c , substitute $t = 0$, $\dot{W}_{cv} = -(6V * 0.22A) = -1.32$

$$T_a = 308.65K \text{ and } T_i - T_e = 0$$

$$\dot{W}_{cv} = K e^{\frac{\lambda_{algaebottle}(t+c)}{m_a c_a}} - \lambda_{algaebottle} (T_a - T_{amb}) + \dot{m}[c_w (T_i - T_e)] \text{ where } K = e^{\frac{\lambda_{algaebottle}(t+c)}{m_a c_a}}$$

$$-1.32 = K - 1.76 (T_a - T_{amb}) + \dot{m}[c_w (T_i - T_e)]$$

D None Yet