

Physical World 2D Report

Dicson Wijaya (1002289), Wenkie Lau (1002219),
Mok Jun Neng (1002286), Charlotte Phang (1002277),
Martin Tan (1002173)

2D Project

Systems World

F06 Team 06

Dicson Candra Wijaya 1002289

Lau Weng Kei Wenkie 1002219

Phang Li Wen Charlotte 1002277

Mok Jun Neng 1002286

Tan Guan Yuan Martin 1002173

Question 2.

Our group used Excel's matrix multiplication (=MMULT) of the entries of error and the k coefficients to get the linear combination of calculated $v(n)$ for every order controller.

We decided to use minimum chi-square in assessing the goodness of fit between the set of calculated values of $v(n)$ and actual values of $v(n)$ given in the question.

Pearson's Chi-Square Test:

$$X^2 = \sum_{i=1}^n \frac{(O_i - E_i)^2}{E_i}$$

where

X^2 = Pearson's cumulative test statistic

O_i = calculated $v(n)$

E_i = actual $v(n)$ given

n = number of cells in the table

A lower X^2 value indicates that the calculated $v(n)$ is closer to the actual $v(n)$ given. Therefore, our objective is to minimize the chi-square value (X^2) for each order controller.

Using Excel Solver, we set the chi-square value (X^2) as the objective function to a minimum and set the unknown k coefficients (k_0, k_1, \dots, k_9) as the changing variable.

Result:

[CHART]

| Order | X^2 |
|-------|-------------|
| 0 | 20.26071659 |
| 1 | 3.089836451 |
| 2 | 0.006754156 |
| 3 | 0.006716346 |
| 4 | 0.006493217 |
| 5 | 0.005102069 |
| 6 | 0.003122692 |
| 7 | 0.002727727 |
| 8 | 0.000374947 |
| 9 | 3.57577E-10 |

The next step is to analyze the result obtained. Since 10 data set is given, the degree of freedom ($N-1$) is 9. A chi-square distribution with 9 degrees of freedom for a lower one-sided test at significance level $\alpha = 0.001$ (extreme) has a critical value of 1.152.

From the table of values, we can see that the minimized chi-square values (X^2) of 2nd order onwards do not exceed this critical value and hence satisfy the condition to be considered “correct”. However, it is also apparent, from the table of values and the graph, that X^2 already approaches 0 at 2nd order and rate of change of X^2 from 2nd order onwards to the 9th order is negligible.

Therefore, we conclude that the team from last year’s 2D project used controller of 2nd order.

This approach is still reasonable even if we have more data points. In fact, minimized chi-square test becomes more relevant with increasing data points. The plot of X^2 vs order would give data points forming the full similarly converging curve. With more data points, there is increased precision in assessing the data point which chi-square value (X^2) first converges to approximately 0. Therefore, the chi-square test is a reliable tool for assessing the goodness of fit between observed and expected values.

The k coefficients in the 2nd order controller is:

| | |
|-------|-------------|
| k_0 | 0.107782668 |
| k_1 | 0.546854887 |
| k_2 | 0.769143476 |

| | |
|-------|-------------|
| k_0 | 0.107782668 |
|-------|-------------|

We have observed that the coefficients of k_0 and k_1 are the same for 2nd and 3rd order controller, with a different coefficient of k_2 because the 3rd order controller has an additional k_3 coefficient. The unchanged values of k_0 and k_1 suggest that the 2nd order controller is indeed sufficient and these k coefficients are “best” or true.

Therefore, we conclude that the sufficient 2nd order controller was used and the corresponding k coefficients are correct.

Bibliography

Appendix C Critical Values for the Chi-Squared Distribution. (2015). *Beyond Basic Statistics*,167-167.

Moore, D. S. (1976). *Chi-Square Tests* (No. Mimeograph). PURDUE UNIV LAFAYETTE IND DEPT OF STATISTICS.