

chapter 5 Performance Analysis

2021/6/27

Regret Analysis

- A player chooses an action $\theta^{(t)} \in K$ every t period, where K is a feasible set of actions.
- The cost function $f^{(t)}$ determines the cost $f^{(t)}(\theta^{(t)})$ for action $\theta^{(t)}$.
- The player decides his action based on the strategy.

Regret Analysis

- How does the player choose an action which minimizes a total cost $\sum f^{(t)}(\theta^{(t)})$?
- Can the cost function be minimized even if it is not unknown?
- We introduce a regret about the strategy.

definition (Regret)

The difference between the total cost of an action based on a strategy A and the total cost of the optimal strategy θ^ is defined as the regret $\text{Regret}(A)$ of strategy A .*

$$\text{Regret}(A) = \sum_{t=1}^T f^{(t)}(\theta^{(t)}) - \sum_{t=1}^T f^{(t)}(\theta^*)$$

Regret Analysis

Regret analysis in online learning

- Let action be the parameter of the online learner $\theta^{(t)} \in \mathbb{R}^m$ given the training data $(\mathbf{x}^{(t)}, y^{(t)})$.
- Let the cost function be a loss function $f^{(t)} = (\mathbf{x}^{(t)}, y^{(t)}, \theta)$.
- In this case, the optimal strategy is the strategy that chooses the action that minimizes the cost function for all training data.

Follow the Leader

- At first, We consider strategy for choosing action that minimizes the total cost to date.

$$\boldsymbol{\theta}^{(t)} = \arg \min_{\boldsymbol{\theta} \in K} \sum_{i=1}^{t-1} f^{(i)}(\boldsymbol{\theta})$$

- This strategy is called Follow the Leader (FTL).

Follow the Leader

- However, there are cases where FTL doesn't work.
- Consider action $\theta \in [-1, 1]$ and cost function $f^{(t)}(\theta) = (1/2)(-1)^t\theta$.
- In this case, the action goes back and forth between -1 and 1 except for the first, as $\theta^{(1)} = 0, \theta^{(2)} = -1, \theta^{(3)} = 1, \dots$
- The cost function is $1/2$ except for the first, as $f^{(1)}(\theta^{(1)}) = 0, f^{(2)}(\theta^{(2)}) = 1/2, f^{(3)}(\theta^{(3)}) = 1/2, \dots$

Follow the Leader

- On the other hand, The optimal strategy is $\theta = 0$ and the total cost of it is 0.
- Therefore, FTL regret in this case doesn't approach 0.
- We have to expand FTL.

Regularized Follow the Leader

- Regularized Follow the Leader (RFTL)

$$\boldsymbol{\theta}^{(t)} = \arg \min_{\boldsymbol{\theta} \in K} \eta \sum_{i=1}^{t-1} f^{(i)}(\boldsymbol{\theta}) + R(\boldsymbol{\theta})$$

- Let $R(\boldsymbol{\theta})$ be convex regularization function. Let $\eta \geq 0$ be parameter that determines the degree of regularization.
- When choosing the first action, the cost function is not presented, so action is determined only by the regularization term.

$$\boldsymbol{\theta}^{(1)} = \arg \min_{\boldsymbol{\theta} \in K} R(\boldsymbol{\theta})$$

Regularized Follow the Leader

- We introduce lemma and definitions to derive RFTL regret.

lemma

For any vector $\mathbf{u} \in K$, the following holds.

$$\sum_{t=1}^T \mathbf{f}^{(t)T} (\boldsymbol{\theta}^{(t)} - \mathbf{u}) \leq \sum_{t=1}^T \mathbf{f}^{(t)T} (\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{t+1}) + \frac{1}{\eta} \left\{ R(\mathbf{u}) - R(\boldsymbol{\theta}^{(1)}) \right\} \quad (1)$$

Regularized Follow the Leader

Proof.

For simplicity, let us assume that $\mathbf{f}^{(0)} = \frac{1}{\eta}R(\boldsymbol{\theta})$ and the algorithm starts at $t = 0$.

$$\sum_{t=0}^T \mathbf{f}^{(t)}(\boldsymbol{\theta}) = \sum_{t=1}^T \mathbf{f}^{(t)}(\boldsymbol{\theta}) + \frac{1}{\eta}R(\boldsymbol{\theta})$$

In this time, the lemma can be expressed as following.

$$\sum_{t=0}^T \mathbf{f}^{(t)T}(\boldsymbol{\theta}^{(t)} - \mathbf{u}) \leq \sum_{t=0}^T \mathbf{f}^{(t)T}(\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)})$$

Regularized Follow the Leader

Proof.

At $t = 0$,
by definition, $\boldsymbol{\theta}^{(1)} = \arg \min_{\boldsymbol{\theta}} R(\boldsymbol{\theta})$ and $\mathbf{f}^{(0)}(\boldsymbol{\theta}^{(1)}) \leq \mathbf{f}^{(0)}(\mathbf{u})$ holds.
therefore,

$$\mathbf{f}^{(0)}(\boldsymbol{\theta}^{(0)}) - \mathbf{f}^{(0)}(\mathbf{u}) \leq \mathbf{f}^{(0)}(\boldsymbol{\theta}^{(0)}) - \mathbf{f}^{(0)}(\boldsymbol{\theta}^{(1)})$$



Regularized Follow the Leader

Proof.

At $t > 0$,
assume that lemma holds for $t = T$.
In this time,

$$\boldsymbol{\theta}^{(T+2)} = \arg \min_{\boldsymbol{\theta}} \sum_{t=0}^{T+1} \mathbf{f}^{(t)}(\boldsymbol{\theta}) \quad (2)$$

$$\boldsymbol{\theta}^{(T+1)} = \arg \min_{\boldsymbol{\theta}} \sum_{t=0}^T \mathbf{f}^{(t)}(\boldsymbol{\theta}) \quad (3)$$



Regularized Follow the Leader

Proof.

Using equation (2) and (3),

$$\begin{aligned} & \sum_{t=0}^{T+1} \mathbf{f}^{(t)T} (\boldsymbol{\theta}^{(t)} - \mathbf{u}) \\ & \leq \sum_{t=0}^{T+1} \mathbf{f}^{(t)} (\boldsymbol{\theta}^{(t)}) - \sum_{t=0}^{T+1} \mathbf{f}^{(t)} (\boldsymbol{\theta}^{(T+2)}) \\ & = \sum_{t=0}^T (\mathbf{f}^{(t)} (\boldsymbol{\theta}^{(t)}) - \mathbf{f}^{(t)} (\boldsymbol{\theta}^{(T+2)})) + \mathbf{f}^{(T+1)} (\boldsymbol{\theta}^{(T+1)}) - \mathbf{f}^{(T+1)} (\boldsymbol{\theta}^{(T+2)}) \\ & \leq \sum_{t=0}^T (\mathbf{f}^{(t)} (\boldsymbol{\theta}^{(t)}) - \mathbf{f}^{(t)} (\boldsymbol{\theta}^{(T+1)})) + \mathbf{f}^{(T+1)} (\boldsymbol{\theta}^{(T+1)}) - \mathbf{f}^{(T+1)} (\boldsymbol{\theta}^{(T+2)}) \\ & = \sum_{t=0}^{T+1} \mathbf{f}^{(t)} (\boldsymbol{\theta}^{(t)}) - \mathbf{f}^{(t)} (\boldsymbol{\theta}^{(t+1)}) \\ & = \sum_{t=0}^{T+1} \mathbf{f}^{(t)} (\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)}) \end{aligned}$$



Regularized Follow the Leader

definition (norm based on positive semi-definite matrix)

We define $\|\mathbf{x}\|_A = \sqrt{\mathbf{x}^T \mathbf{A} \mathbf{x}}$ as the norm of vector \mathbf{x} based on positive semi-definite matrix \mathbf{A} .

We also define $\|\mathbf{x}\|_{A^{-1}} = \|\mathbf{x}\|_A^*$ as a dual norm.

- In this time, from generalized Cauchy-Schwarz inequality, the following holds.

$$\mathbf{x}^T \mathbf{y} \leq \|\mathbf{x}\|_A \|\mathbf{y}\|_A^*$$

Regularized Follow the Leader

definition (norm of cost function)

A norm of cost function measured by the regularization function is defined as

$$\lambda = \max_{t, \boldsymbol{\theta} \in K} \mathbf{f}^{(t)T} \{\nabla^2 R(\boldsymbol{\theta})\}^{-1} \mathbf{f}^{(t)}$$

Regularized Follow the Leader

definition (diameter of feasible area)

A diameter measured by the regularization function is defined as

$$D = \max_{\boldsymbol{\theta} \in K} R(\boldsymbol{\theta}) - R(\boldsymbol{\theta}^{(1)})$$

Regularized Follow the Leader

theorem (regret of RFTL)

RFTL achieves the following regret for any vector $\mathbf{u} \in K$.

$$\text{Regret}(A) = \sum_{t=1}^T \mathbf{f}^{(t)T} (\boldsymbol{\theta}^{(t)} - \mathbf{u}) \leq 2\sqrt{2\lambda DT}$$

Regularized Follow the Leader

Proof.

At first, we define Φ as following.

$$\Phi^{(t)}(\boldsymbol{\theta}) = \eta \sum_{i=1}^t f^{(i)}(\boldsymbol{\theta}) + R(\boldsymbol{\theta})$$

By Taylor expansion of $\Phi^{(t)}$ around $\boldsymbol{\theta}^{(t+1)}$ and using intermediate value theorem, we can show following.

$$\begin{aligned}\Phi^{(t)}(\boldsymbol{\theta}^{(t)}) &= \Phi^{(t)}(\boldsymbol{\theta}^{(t+1)}) + (\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)})^T \nabla \Phi^{(t)}(\boldsymbol{\theta}^{(t+1)}) \\ &\quad + \frac{1}{2} \|\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)}\|_{\mathbf{z}^{(t)}}^2 \\ &\geq \Phi^{(t)}(\boldsymbol{\theta}^{(t+1)}) + \frac{1}{2} \|\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)}\|_{\mathbf{z}^{(t)}}^2\end{aligned}$$



Regularized Follow the Leader

Proof.

Here, from intermediate value theorem, $\mathbf{z}^{(t)} \in [\boldsymbol{\theta}^{(t+1)}, \boldsymbol{\theta}^{(t)}]$

This inequality holds because $\boldsymbol{\theta}^{(t)}$ achieves the minimum value of $\Phi^{(t)}$.

By transforming the equation, we can get following.

$$\begin{aligned}\|\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)}\|_{\mathbf{z}^{(t)}}^2 &\leq 2(\Phi^{(t)}(\boldsymbol{\theta}^{(t)}) - \Phi^{(t)}(\boldsymbol{\theta}^{(t+1)})) \\ &= 2(\Phi^{(t-1)}(\boldsymbol{\theta}^{(t)}) - \Phi^{(t-1)}(\boldsymbol{\theta}^{(t+1)})) + 2\eta \mathbf{f}^{(t)T}(\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)}) \\ &\leq 2\eta \mathbf{f}^{(t)T}(\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)})\end{aligned}$$



Regularized Follow the Leader

Proof.

Also, using generalized Cauchy-Schwarz inequality, the following holds.

$$\begin{aligned} \mathbf{f}^{(t)T}(\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)}) &\leq \|\mathbf{f}^{(t)}\|_z^* \|\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)}\|_z \\ &\leq \|\mathbf{f}^{(t)}\|_z^* \sqrt{2\eta \mathbf{f}^{(t)T}(\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)})} \end{aligned}$$

furthermore,

$$\mathbf{f}^{(t)T}(\boldsymbol{\theta}^{(t)} - \boldsymbol{\theta}^{(t+1)}) \leq 2\eta \|\mathbf{f}^{(t)}\|_z^{*2} \leq 2\eta\lambda \quad (4)$$



Regularized Follow the Leader

Proof.

Using (1) and (4), adding up to T and letting η be $\eta = \sqrt{\frac{D}{2\lambda T}}$ which minimizes this equation, we can get

$$\text{Regret}(A) \leq \min_{\eta} \left[2\eta\lambda T + \frac{1}{\eta} \{R(\mathbf{u}) - R(\boldsymbol{\theta}^{(t)})\} \right] \leq 2\sqrt{2D\lambda T}$$

