

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Unit 0: ...](#) > [Contin...](#) > 15. Su...

Audit Access Expires Mar 10, 2021

You lose all access to this course, including your progress, on Mar 10, 2021.

15. Summary

Definition of continuity at a point

We say that a function f is **continuous at a point** $x = a$ if

$$\lim_{x \rightarrow a} f(x) = f(a).$$

In particular, if either $f(a)$ or $\lim_{x \rightarrow a} f(x)$ fails to exist, then f is discontinuous at a .

We say that a function f is **right-continuous at a point** $x = a$ if

$$\lim_{x \rightarrow a^+} f(x) = f(a).$$


We say that a function f is **left-continuous at a point** $x = a$ if $\lim_{x \rightarrow a^-} f(x) = f(a)$.

Types of Discontinuities


It is sometimes useful to classify certain types of discontinuities.

If the left-hand limit $\lim_{x \rightarrow a^-} f(x)$ and the right-hand limit $\lim_{x \rightarrow a^+} f(x)$ both exist at a point $x = a$, but they are not equal, then we say that f has a **jump discontinuity** at $x = a$.





If the overall limit $\lim_{x \rightarrow a} f(x)$ exists (i.e., the left- and right-hand limits agree), but the overall limit does not equal $f(a)$, then we say that f has a **removable discontinuity** at $x = a$.



Definition of continuous functions

A function $f(x)$ is **continuous** if for every point c in the domain of $f(x)$, the function f is continuous at the point $x = c$.

Basic Continuous Functions

Note: we have not proven all of the following facts, but you should feel free to use them.

The following functions are continuous at *all real numbers*:

- all polynomials

- $\sqrt[3]{x}$

- $|x|$



- $\cos x$ and $\sin x$
- exponential functions a^x with base $a > 0$

The following functions are continuous *at the specified values of x* :

- \sqrt{x} , for $x > 0$
- $\tan x$, at all x where it is defined
- logarithmic functions $\log_a x$ with base $a > 0$, for $x > 0$

Limit laws and continuity

If the functions f and g are continuous everywhere, then:

- $f + g$ is continuous everywhere.
- $f - g$ is continuous everywhere.
- $f \cdot g$ is continuous everywhere.
- $\frac{f}{g}$ is continuous where it is defined.

Intermediate Value Theorem

If f is a function which is continuous on the interval $[a, b]$, and M lies between the values of $f(a)$ and $f(b)$, then there is at least one point c between a and b such that $f(c) = M$.

(A function f is **continuous on a closed interval** $[a, b]$ if it is right-continuous at a , left-continuous at b , and continuous at all points between a and b .)

15. Summary

[Hide Discussion](#)

Topic: Unit 0: Limits / 15. Summary

Add a Post

Show all posts ▼

by recent ▼



?	<u>Continuity in the case of imaginary numbers.</u>	1
	Are functions of the form $\sqrt{x + y}$, where y is a positive real number, continuous at every point?...	
?	<u>CUBE ROOT VS SQUARE ROOT</u>	3
	Why is $\sqrt[3]{x}$ continuous at all points even when $x=0$, and \sqrt{x} is continuous for $x>0$?	
💬	<u>Format has changed</u>	3
	It may just be my computer, but in this most recent section the format is quite different. For instan...	

© All Rights Reserved

