NURS 1017 Unit 7 Case Study 2

19-year-old male client who fell from a longboard without a helmet: recognizing a head injury, early and late signs of increased ICP during neurological checks (epidural hematoma with a lucid interval), the nurse's response to deterioration, nursing care to prevent secondary brain injury after craniotomy (positioning, suctioning, fever, sedation and hypertonic saline monitoring), and discharge teaching on medication effects, graded return to activity and TBI prevention.

Readiness problems: none

NURS 1017 Unit 7 Case Study 2, screen 1

Practice NCLEX RN Test

Question 1 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 1 of 6

The nurse reviews the Nurses' Notes and the Vital Signs. **Highlight** the findings that require follow-up.

Nurses' Notes | **Vital Signs**

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 2, screen 1 (answer key, score 6/6)

Practice NCLEX RN Test

Question 1 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 1 of 6

The nurse reviews the Nurses' Notes and the Vital Signs. **Highlight** the findings that require follow-up.

Nurses' Notes | **Vital Signs**

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 2, screen 2

Practice NCLEX RN Test

Question 2 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 2 of 6

The nurse in the emergency department is caring for a 19-year-old male client who fell from a longboard and struck his head.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

1615: Neurosurgery consulted; plan is observation in the ED with neurological checks every 15 minutes and a repeat CT in 4 hours. Client states his headache is "a bit better" and asks to go home.

The nurse has reviewed the Diagnostic Results, the Laboratory Results and the Nurses' Notes from 1615.

➤ The nurse is performing neurological checks every 15 minutes. For each finding, click to specify whether it would be an **early sign** of increased intracranial pressure (ICP), a **late sign** of increased ICP, or **not a sign** of increased ICP.

Finding	Early sign	Late sign	Not a sign
New restlessness, irritability, or confusion	○	○	○
One pupil dilated and not reacting to light	○	○	○
Headache that is getting worse, with vomiting	○	○	○
Abrasion over the right temple	○	○	○
Rising systolic BP with a widening pulse pressure, a slowing pulse, and irregular breathing	○	○	○
Takes longer to answer, or needs a louder voice or a touch to stay awake	○	○	○
Abnormal flexion or extension of the arms in response to a painful stimulus	○	○	○
Pupils equal, round, and briskly reactive	○	○	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 2, screen 2 (answer key, score 8/8)

Practice NCLEX RN Test

Question 2 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 2 of 6

The nurse in the emergency department is caring for a 19-year-old male client who fell from a longboard and struck his head.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

1615: Neurosurgery consulted; plan is observation in the ED with neurological checks every 15 minutes and a repeat CT in 4 hours. Client states his headache is "a bit better" and asks to go home.

The nurse has reviewed the Diagnostic Results, the Laboratory Results and the Nurses' Notes from 1615.

➤ The nurse is performing neurological checks every 15 minutes. For each finding, click to specify whether it would be an **early sign** of increased intracranial pressure (ICP), a **late sign** of increased ICP, or **not a sign** of increased ICP.

Finding	Early sign	Late sign	Not a sign
New restlessness, irritability, or confusion	●	○	○
One pupil dilated and not reacting to light	○	●	○
Headache that is getting worse, with vomiting	●	○	○
Abrasion over the right temple	○	○	●
Rising systolic BP with a widening pulse pressure, a slowing pulse, and irregular breathing	○	●	○
Takes longer to answer, or needs a louder voice or a touch to stay awake	●	○	○
Abnormal flexion or extension of the arms in response to a painful stimulus	○	●	○
Pupils equal, round, and briskly reactive	○	○	●

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 2, screen 3

Practice NCLEX RN Test

Question 3 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 3 of 6

The nurse in the emergency department is caring for a 19-year-old male client who fell from a longboard and struck his head.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

1615: Neurosurgery consulted; plan is observation in the ED with neurological checks every 15 minutes and a repeat CT in 4 hours. Client states his headache is "a bit better" and asks to go home.

1720: Client difficult to rouse: opens eyes only to a painful stimulus, speaks in confused, inappropriate words, and localizes pain (GCS 10, E2 V3 M5). Vomited twice since 1700. Right pupil 6 mm and sluggish; left pupil 3 mm and brisk. Left arm and leg weaker than the right (3/5). Respirations irregular.

The nurse has reviewed the Nurses' Notes from 1720 and the Vital Signs from 1720.

Complete the following sentences by choosing from the lists of options.

The client's findings indicate Select... , as shown by Select... . The nurse's first action is to Select... .

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 2, screen 3 (answer key, score 3/3)

Practice NCLEX RN Test

Question 3 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 3 of 6

The nurse in the emergency department is caring for a 19-year-old male client who fell from a longboard and struck his head.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

1615: Neurosurgery consulted; plan is observation in the ED with neurological checks every 15 minutes and a repeat CT in 4 hours. Client states his headache is "a bit better" and asks to go home.

1720: Client difficult to rouse: opens eyes only to a painful stimulus, speaks in confused, inappropriate words, and localizes pain (GCS 10, E2 V3 M5). Vomited twice since 1700. Right pupil 6 mm and sluggish; left pupil 3 mm and brisk. Left arm and leg weaker than the right (3/5). Respirations irregular.

The nurse has reviewed the Nurses' Notes from 1720 and the Vital Signs from 1720.

➤ Complete the following sentences by choosing from the lists of options.

The client's findings indicate [rapidly rising ICP from the expanding epidural hematoma] , as shown by [a falling GCS, a dilated right pupil, left-sided weakness, and Cushing's triad] . The nurse's first action is to

[notify the provider immediately and prepare him for emergency surgery] .

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 2, screen 4

Practice NCLEX RN Test

Question 4 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 4 of 6

The nurse in the emergency department is caring for a 19-year-old male client who fell from a longboard and struck his head.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

1615: Neurosurgery consulted; plan is observation in the ED with neurological checks every 15 minutes and a repeat CT in 4 hours. Client states his headache is "a bit better" and asks to go home.

1720: Client difficult to rouse: opens eyes only to a painful stimulus, speaks in confused, inappropriate words, and localizes pain (GCS 10, E2 V3 M5). Vomited twice since 1700. Right pupil 6 mm and sluggish; left pupil 3 mm and brisk. Left arm and leg weaker than the right (3/5). Respirations irregular.

2130: Returned from the operating room after emergency right temporal craniotomy and evacuation of the epidural hematoma. Admitted to the neurological ICU. Intubated and mechanically ventilated; sedated with propofol and fentanyl infusions. ICP monitor in place: ICP 12 mm Hg. Pupils 3 mm, equal and reactive. Moves all limbs to painful stimuli; left side weaker.

The nurse has reviewed the Nurses' Notes from 2130 and the Provider Orders.

➤ The nurse plans care to prevent secondary brain injury. For each intervention, click to specify whether it is **indicated** or **contraindicated**.

Intervention	Indicated	Contraindicated
Keep the head and neck in midline alignment and avoid tight endotracheal tube ties around the neck	○	○
Position the client with the hips flexed more than 90 degrees	○	○
Compare each neurological check with the previous one and report any decline	○	○
Change the maintenance IV fluid to 5% dextrose in water	○	○
Monitor the BP, respirations and sedation level closely while the propofol and fentanyl infusions run	○	○
Hyperoxygenate before suctioning and limit each suction pass to 10 seconds or less	○	○
Group the bath, linen change, suctioning, and repositioning into one long session	○	○
Turn the client every 2 hours, keeping the head in line with the body	○	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 2, screen 4 (answer key, score 8/8)

Practice NCLEX RN Test

Question 4 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 4 of 6

The nurse in the emergency department is caring for a 19-year-old male client who fell from a longboard and struck his head.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

1615: Neurosurgery consulted; plan is observation in the ED with neurological checks every 15 minutes and a repeat CT in 4 hours. Client states his headache is "a bit better" and asks to go home.

1720: Client difficult to rouse: opens eyes only to a painful stimulus, speaks in confused, inappropriate words, and localizes pain (GCS 10, E2 V3 M5). Vomited twice since 1700. Right pupil 6 mm and sluggish; left pupil 3 mm and brisk. Left arm and leg weaker than the right (3/5). Respirations irregular.

2130: Returned from the operating room after emergency right temporal craniotomy and evacuation of the epidural hematoma. Admitted to the neurological ICU. Intubated and mechanically ventilated; sedated with propofol and fentanyl infusions. ICP monitor in place: ICP 12 mm Hg. Pupils 3 mm, equal and reactive. Moves all limbs to painful stimuli; left side weaker.

The nurse has reviewed the Nurses' Notes from 2130 and the Provider Orders.

➤ The nurse plans care to prevent secondary brain injury. For each intervention, click to specify whether it is **indicated** or **contraindicated**.

Intervention	Indicated	Contraindicated
Keep the head and neck in midline alignment and avoid tight endotracheal tube ties around the neck	●	○
Position the client with the hips flexed more than 90 degrees	○	●
Compare each neurological check with the previous one and report any decline	●	○
Change the maintenance IV fluid to 5% dextrose in water	○	●
Monitor the BP, respirations and sedation level closely while the propofol and fentanyl infusions run	●	○
Hyperoxygenate before suctioning and limit each suction pass to 10 seconds or less	●	○
Group the bath, linen change, suctioning, and repositioning into one long session	○	●
Turn the client every 2 hours, keeping the head in line with the body	●	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 2, screen 5

Practice NCLEX RN Test

Question 5 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 5 of 6

The nurse in the emergency department is caring for a 19-year-old male client who fell from a longboard and struck his head.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

1615: Neurosurgery consulted; plan is observation in the ED with neurological checks every 15 minutes and a repeat CT in 4 hours. Client states his headache is "a bit better" and asks to go home.

1720: Client difficult to rouse: opens eyes only to a painful stimulus, speaks in confused, inappropriate words, and localizes pain (GCS 10, E2 V3 M5). Vomited twice since 1700. Right pupil 6 mm and sluggish; left pupil 3 mm and brisk. Left arm and leg weaker than the right (3/5). Respirations irregular.

2130: Returned from the operating room after emergency right temporal craniotomy and evacuation of the epidural hematoma. Admitted to the neurological ICU. Intubated and mechanically ventilated; sedated with propofol and fentanyl infusions. ICP monitor in place: ICP 12 mm Hg. Pupils 3 mm, equal and reactive. Moves all limbs to painful stimuli; left side weaker.

Day 2 0400: Client coughing and pushing against the ventilator during repositioning; sedation scale shows light sedation. ICP 26 mm Hg for the last 10 minutes; MAP 78 mm Hg. Head turned to the right on the pillow. Skin warm and flushed. Has received two doses of 3% sodium chloride overnight.

The nurse has reviewed the Nurses' Notes from Day 2 0400, the Vital Signs from Day 2 0400 and the Laboratory Results from Day 2 0200.

Which actions should the nurse take? **Select all that apply.**

- [ ] 1. Lower the head of the bed flat to increase blood flow to the brain
- [ ] 2. Check the IV site of the hypertonic saline for redness, swelling or pain
- [ ] 3. Give the prescribed acetaminophen and apply cooling measures
- [ ] 4. Complete the morning bed bath now while the client is awake
- [ ] 5. Reposition the head to midline and confirm the head of the bed is at 30 degrees
- [ ] 6. Administer the prescribed PRN fentanyl
- [ ] 7. Suction the endotracheal tube every 15 minutes to prevent coughing
- [ ] 8. Notify the provider of the ICP and CPP and the latest sodium result

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 2, screen 5 (answer key, score 5/5)

Practice NCLEX RN Test

Question 5 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 5 of 6

The nurse in the emergency department is caring for a 19-year-old male client who fell from a longboard and struck his head.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

1615: Neurosurgery consulted; plan is observation in the ED with neurological checks every 15 minutes and a repeat CT in 4 hours. Client states his headache is "a bit better" and asks to go home.

1720: Client difficult to rouse: opens eyes only to a painful stimulus, speaks in confused, inappropriate words, and localizes pain (GCS 10, E2 V3 M5). Vomited twice since 1700. Right pupil 6 mm and sluggish; left pupil 3 mm and brisk. Left arm and leg weaker than the right (3/5). Respirations irregular.

2130: Returned from the operating room after emergency right temporal craniotomy and evacuation of the epidural hematoma. Admitted to the neurological ICU. Intubated and mechanically ventilated; sedated with propofol and fentanyl infusions. ICP monitor in place: ICP 12 mm Hg. Pupils 3 mm, equal and reactive. Moves all limbs to painful stimuli; left side weaker.

Day 2 0400: Client coughing and pushing against the ventilator during repositioning; sedation scale shows light sedation. ICP 26 mm Hg for the last 10 minutes; MAP 78 mm Hg. Head turned to the right on the pillow. Skin warm and flushed. Has received two doses of 3% sodium chloride overnight.

The nurse has reviewed the Nurses' Notes from Day 2 0400, the Vital Signs from Day 2 0400 and the Laboratory Results from Day 2 0200.

Which actions should the nurse take? **Select all that apply.**

- [ ] 1. Lower the head of the bed flat to increase blood flow to the brain
- [x] 2. Check the IV site of the hypertonic saline for redness, swelling or pain
- [x] 3. Give the prescribed acetaminophen and apply cooling measures
- [ ] 4. Complete the morning bed bath now while the client is awake
- [x] 5. Reposition the head to midline and confirm the head of the bed is at 30 degrees
- [x] 6. Administer the prescribed PRN fentanyl
- [ ] 7. Suction the endotracheal tube every 15 minutes to prevent coughing
- [x] 8. Notify the provider of the ICP and CPP and the latest sodium result

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 2, screen 6

Practice NCLEX RN Test

Question 6 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 6 of 6

The nurse in the emergency department is caring for a 19-year-old male client who fell from a longboard and struck his head.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

1615: Neurosurgery consulted; plan is observation in the ED with neurological checks every 15 minutes and a repeat CT in 4 hours. Client states his headache is "a bit better" and asks to go home.

1720: Client difficult to rouse: opens eyes only to a painful stimulus, speaks in confused, inappropriate words, and localizes pain (GCS 10, E2 V3 M5). Vomited twice since 1700. Right pupil 6 mm and sluggish; left pupil 3 mm and brisk. Left arm and leg weaker than the right (3/5). Respirations irregular.

2130: Returned from the operating room after emergency right temporal craniotomy and evacuation of the epidural hematoma. Admitted to the neurological ICU. Intubated and mechanically ventilated; sedated with propofol and fentanyl infusions. ICP monitor in place: ICP 12 mm Hg. Pupils 3 mm, equal and reactive. Moves all limbs to painful stimuli; left side weaker.

Day 2 0400: Client coughing and pushing against the ventilator during repositioning; sedation scale shows light sedation. ICP 26 mm Hg for the last 10 minutes; MAP 78 mm Hg. Head turned to the right on the pillow. Skin warm and flushed. Has received two doses of 3% sodium chloride overnight.

Day 9 1000: Extubated on Day 4; transferred to the neurosurgical unit on Day 5. GCS 15. Mild

The nurse has reviewed the Nurses' Notes from Day 9 1000 and the Vital Signs from Day 9 1000.

For each statement the client makes, click to specify whether it indicates an **understanding** of the discharge teaching or a **need for further teaching**.

Client Statement	Understanding	Needs further teaching
"I'll wear a certified helmet every time I ride my longboard or bike."	○	○
"I can go back to playing hockey as soon as my headache is gone."	○	○
"If I feel unusually irritable, down, or very sleepy on the levetiracetam, I'll tell my doctor."	○	○
"A few beers at the party this weekend won't affect how my brain heals."	○	○
"I'll drive myself to my follow-up appointment next week."	○	○
"If my headache gets much worse, I keep vomiting, or I get very sleepy or confused, I'll go to emergency."	○	○
"I'll start with short periods of light activity and limit screen time, then increase slowly if my symptoms don't get worse."	○	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Finish attempt →

NURS 1017 Unit 7 Case Study 2, screen 6 (answer key, score 7/7)

Practice NCLEX RN Test

Question 6 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 6 of 6

The nurse in the emergency department is caring for a 19-year-old male client who fell from a longboard and struck his head.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results** | **Provider Orders**

1540: Arrived by ambulance after falling from his longboard at about 1500 and striking the right side of his head on a curb. He was not wearing a helmet. His friend states he was "knocked out for about a minute", then woke up and was talking. Client is alert. Oriented to person and place but not to time; repeatedly asks "What happened?" and cannot remember the fall. Reports a headache rated 6/10 with nausea; vomited once in the ambulance. 4-cm boggy swelling with an abrasion over the right temporal area. No drainage from the nose or ears. Pupils equal, 3 mm, briskly reactive. Moves all extremities with equal strength. Glasgow Coma Scale (GCS) 14 (E4 V4 M6). Neck nontender with full range of motion; cervical collar removed by the ED physician. Plays recreational hockey; had a concussion last winter.

1615: Neurosurgery consulted; plan is observation in the ED with neurological checks every 15 minutes and a repeat CT in 4 hours. Client states his headache is "a bit better" and asks to go home.

1720: Client difficult to rouse: opens eyes only to a painful stimulus, speaks in confused, inappropriate words, and localizes pain (GCS 10, E2 V3 M5). Vomited twice since 1700. Right pupil 6 mm and sluggish; left pupil 3 mm and brisk. Left arm and leg weaker than the right (3/5). Respirations irregular.

2130: Returned from the operating room after emergency right temporal craniotomy and evacuation of the epidural hematoma. Admitted to the neurological ICU. Intubated and mechanically ventilated; sedated with propofol and fentanyl infusions. ICP monitor in place: ICP 12 mm Hg. Pupils 3 mm, equal and reactive. Moves all limbs to painful stimuli; left side weaker.

Day 2 0400: Client coughing and pushing against the ventilator during repositioning; sedation scale shows light sedation. ICP 26 mm Hg for the last 10 minutes; MAP 78 mm Hg. Head turned to the right on the pillow. Skin warm and flushed. Has received two doses of 3% sodium chloride overnight.

Day 9 1000: Extubated on Day 4; transferred to the neurosurgical unit on Day 5. GCS 15. Mild

The nurse has reviewed the Nurses' Notes from Day 9 1000 and the Vital Signs from Day 9 1000.

➤ For each statement the client makes, click to specify whether it indicates an **understanding** of the discharge teaching or a **need for further teaching**.

Client Statement	Understanding	Needs further teaching
"I'll wear a certified helmet every time I ride my longboard or bike."	●	○
"I can go back to playing hockey as soon as my headache is gone."	○	●
"If I feel unusually irritable, down, or very sleepy on the levetiracetam, I'll tell my doctor."	●	○
"A few beers at the party this weekend won't affect how my brain heals."	○	●
"I'll drive myself to my follow-up appointment next week."	○	●
"If my headache gets much worse, I keep vomiting, or I get very sleepy or confused, I'll go to emergency."	●	○
"I'll start with short periods of light activity and limit screen time, then increase slowly if my symptoms don't get worse."	●	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Finish attempt →

NURS 1017 Unit 7 Case Study 3

19-year-old university student with meningococcal meningitis and early sepsis: recognizing the hallmark signs of meningeal irritation, increased ICP and sepsis (including the nonblanching rash), nursing priorities, droplet precautions, a quiet environment, seizure precautions and public health follow-up, monitoring for vancomycin and dexamethasone adverse effects, care during and after a seizure, communicating hydrocephalus, and evaluation of complications such as hearing loss.

Readiness problems: none

NURS 1017 Unit 7 Case Study 3, screen 1

Practice NCLEX RN Test

Question 1 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 1 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | **Vital Signs**

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

The nurse reviews the client's history and assessment. Which of the following findings would be consistent with **bacterial meningitis**? **Select all that apply.**

- [ ] 1. Photophobia
- [ ] 2. Abdomen soft and nontender
- [ ] 3. Confusion and drowsiness
- [ ] 4. Neck stiffness with pain on flexion
- [ ] 5. Positive Kernig and Brudzinski signs
- [ ] 6. Temperature 39.6° C
- [ ] 7. Last menstrual period 2 weeks ago
- [ ] 8. SpO_2 96% on room air
- [ ] 9. Severe headache with vomiting
- [ ] 10. Pupils equal and reactive

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 3, screen 1 (answer key, score 6/6)

Practice NCLEX RN Test

Question 1 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 1 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | **Vital Signs**

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

The nurse reviews the client's history and assessment. Which of the following findings would be consistent with **bacterial meningitis**? **Select all that apply.**

- [x] 1. Photophobia
- [ ] 2. Abdomen soft and nontender
- [x] 3. Confusion and drowsiness
- [x] 4. Neck stiffness with pain on flexion
- [x] 5. Positive Kernig and Brudzinski signs
- [x] 6. Temperature 39.6° C
- [ ] 7. Last menstrual period 2 weeks ago
- [ ] 8. SpO_2 96% on room air
- [x] 9. Severe headache with vomiting
- [ ] 10. Pupils equal and reactive

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 3, screen 2

Practice NCLEX RN Test

Question 2 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 2 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

Day 1 2215: Two sets of blood cultures drawn. Dexamethasone IV given at 2218, then ceftriaxone IV and vancomycin IV started at 2220. Droplet precautions in place.

Day 1 2310: Lumbar puncture done after the head CT; CSF sent to the laboratory.

The nurse has reviewed the Nurses' Notes from Day 1 2215 and 2310, the Diagnostic Results and the Laboratory Results.

➤ For each finding, click to specify whether it is a sign of **meningeal irritation**, **increased intracranial pressure (ICP)**, or **sepsis**. Each finding may be a sign of more than one problem.

Finding	Meningeal irritation	Increased ICP	Sepsis
Neck stiffness with pain on flexion	☐	☐	☐
Drowsy and oriented to person only	☐	☐	☐
Nonblanching petechial rash on the legs and trunk	☐	☐	☐
Positive Kernig and Brudzinski signs	☐	☐	☐
BP 96/58 mm Hg, cool hands and feet, and a capillary refill of 4 seconds	☐	☐	☐
Photophobia	☐	☐	☐
CSF opening pressure of 32 cm H_2O	☐	☐	☐
Lactate 4.1 mmol/L	☐	☐	☐

Note: Each column must have at least 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 3, screen 2 (answer key, score 9/9)

Practice NCLEX RN Test

Question 2 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 2 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

Day 1 2215: Two sets of blood cultures drawn. Dexamethasone IV given at 2218, then ceftriaxone IV and vancomycin IV started at 2220. Droplet precautions in place.

Day 1 2310: Lumbar puncture done after the head CT; CSF sent to the laboratory.

The nurse has reviewed the Nurses' Notes from Day 1 2215 and 2310, the Diagnostic Results and the Laboratory Results.

➢ For each finding, click to specify whether it is a sign of **meningeal irritation**, **increased intracranial pressure (ICP)**, or **sepsis**. Each finding may be a sign of more than one problem.

Finding	Meningeal irritation	Increased ICP	Sepsis
Neck stiffness with pain on flexion	☑	☐	☐
Drowsy and oriented to person only	☐	☑	☑
Nonblanching petechial rash on the legs and trunk	☐	☐	☑
Positive Kernig and Brudzinski signs	☑	☐	☐
BP 96/58 mm Hg, cool hands and feet, and a capillary refill of 4 seconds	☐	☐	☑
Photophobia	☑	☐	☐
CSF opening pressure of 32 cm H_2O	☐	☑	☐
Lactate 4.1 mmol/L	☐	☐	☑

Note: Each column must have at least 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 3, screen 3

Practice NCLEX RN Test

Question 3 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 3 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

Day 1 2215: Two sets of blood cultures drawn. Dexamethasone IV given at 2218, then ceftriaxone IV and vancomycin IV started at 2220. Droplet precautions in place.

Day 1 2310: Lumbar puncture done after the head CT; CSF sent to the laboratory.

Complete the following sentences by choosing from the lists of options.

The client is at highest risk for Select... as a result of Select... . She is also at risk for Select... , as shown by Select... .

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 3, screen 3 (answer key, score 4/4)

Practice NCLEX RN Test

Question 3 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 3 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

Day 1 2215: Two sets of blood cultures drawn. Dexamethasone IV given at 2218, then ceftriaxone IV and vancomycin IV started at 2220. Droplet precautions in place.

Day 1 2310: Lumbar puncture done after the head CT; CSF sent to the laboratory.

Complete the following sentences by choosing from the lists of options.

The client is at highest risk for [increased intracranial pressure] as a result of [cerebral edema from the inflammatory response]. She is also at risk for [septic shock and disseminated intravascular coagulation], as shown by [hypotension, a nonblanching rash, a high lactate, and a low platelet count].

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 3, screen 4

Practice NCLEX RN Test

Question 4 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 4 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

Day 1 2215: Two sets of blood cultures drawn. Dexamethasone IV given at 2218, then ceftriaxone IV and vancomycin IV started at 2220. Droplet precautions in place.

Day 1 2310: Lumbar puncture done after the head CT; CSF sent to the laboratory.

The nurse has reviewed the Provider Orders.

➤ Which interventions should the nurse include in the client's plan of care? **Select all that apply.**

- [ ] 1. Maintain droplet precautions until she has received 24 hours of effective antibiotic therapy
- [ ] 2. Watch for flushing, itching and low BP during the vancomycin infusion
- [ ] 3. Keep the room quiet and dimly lit
- [ ] 4. Restrict fluids to 500 mL per day to prevent cerebral edema
- [ ] 5. Support the neck in flexion on two pillows to ease the neck pain
- [ ] 6. Elevate the head of the bed to 30 degrees
- [ ] 7. Place her in a negative-pressure room and wear an N95 respirator for all care
- [ ] 8. Implement seizure precautions
- [ ] 9. Confirm that public health has been notified so close contacts can receive preventive antibiotics
- [ ] 10. Check her capillary glucose while she receives dexamethasone

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 3, screen 4 (answer key, score 7/7)

Practice NCLEX RN Test

Question 4 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 4 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

Day 1 2215: Two sets of blood cultures drawn. Dexamethasone IV given at 2218, then ceftriaxone IV and vancomycin IV started at 2220. Droplet precautions in place.

Day 1 2310: Lumbar puncture done after the head CT; CSF sent to the laboratory.

The nurse has reviewed the Provider Orders.

➤ Which interventions should the nurse include in the client's plan of care? **Select all that apply.**

- [x] 1. Maintain droplet precautions until she has received 24 hours of effective antibiotic therapy
- [x] 2. Watch for flushing, itching and low BP during the vancomycin infusion
- [x] 3. Keep the room quiet and dimly lit
- [ ] 4. Restrict fluids to 500 mL per day to prevent cerebral edema
- [ ] 5. Support the neck in flexion on two pillows to ease the neck pain
- [x] 6. Elevate the head of the bed to 30 degrees
- [ ] 7. Place her in a negative-pressure room and wear an N95 respirator for all care
- [x] 8. Implement seizure precautions
- [x] 9. Confirm that public health has been notified so close contacts can receive preventive antibiotics
- [x] 10. Check her capillary glucose while she receives dexamethasone

Submit

← Quit

← Previous

Next →

NOT COMPLETE

Case Study Screen 5 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results** | **Provider Orders**

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

Day 1 2215: Two sets of blood cultures drawn. Dexamethasone IV given at 2218, then ceftriaxone IV and vancomycin IV started at 2220. Droplet precautions in place.

Day 1 2310: Lumbar puncture done after the head CT; CSF sent to the laboratory.

Day 2 0545: Client had a generalized tonic-clonic seizure lasting 2 minutes: sudden stiffening of the whole body, then rhythmic jerking of all four limbs; no warning or focal movements beforehand. Now postictal: snoring respirations, opens eyes to pain only (GCS 8). Headache had been worsening overnight, with two episodes of vomiting. Pupils equal, 4 mm, sluggish.

The nurse has reviewed the Nurses' Notes from Day 2 0545, the Vital Signs from Day 2 0600 and the Provider Orders from Day 2 0550.

➤ For each action, click to specify whether it is **indicated** or **contraindicated** for the client at this time.

Nursing Action	Indicated	Contraindicated
Turn the client onto her side and suction the mouth as needed	○	○
Place a padded tongue blade between her teeth in case another seizure occurs	○	○
Apply oxygen and monitor her airway and breathing closely	○	○
Administer the levetiracetam loading dose	○	○
Prepare her for a repeat lumbar puncture to drain CSF and lower the pressure	○	○
Discontinue droplet precautions because she has received 8 hours of antibiotics	○	○
Prepare her for the STAT head CT	○	○

Note: Each row must have only 1 response option selected.

Submit

NURS 1017 Unit 7 Case Study 3, screen 5 (answer key, score 7/7)

Practice NCLEX RN Test

Question 5 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 5 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results** | **Provider Orders**

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

Day 1 2215: Two sets of blood cultures drawn. Dexamethasone IV given at 2218, then ceftriaxone IV and vancomycin IV started at 2220. Droplet precautions in place.

Day 1 2310: Lumbar puncture done after the head CT; CSF sent to the laboratory.

Day 2 0545: Client had a generalized tonic-clonic seizure lasting 2 minutes: sudden stiffening of the whole body, then rhythmic jerking of all four limbs; no warning or focal movements beforehand. Now postictal: snoring respirations, opens eyes to pain only (GCS 8). Headache had been worsening overnight, with two episodes of vomiting. Pupils equal, 4 mm, sluggish.

The nurse has reviewed the Nurses' Notes from Day 2 0545, the Vital Signs from Day 2 0600 and the Provider Orders from Day 2 0550.

➤ For each action, click to specify whether it is **indicated** or **contraindicated** for the client at this time.

Nursing Action	Indicated	Contraindicated
Turn the client onto her side and suction the mouth as needed	●	○
Place a padded tongue blade between her teeth in case another seizure occurs	○	●
Apply oxygen and monitor her airway and breathing closely	●	○
Administer the levetiracetam loading dose	●	○
Prepare her for a repeat lumbar puncture to drain CSF and lower the pressure	○	●
Discontinue droplet precautions because she has received 8 hours of antibiotics	○	●
Prepare her for the STAT head CT	●	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

NOT COMPLETE

Case Study Screen 6 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

Day 1 2215: Two sets of blood cultures drawn. Dexamethasone IV given at 2218, then ceftriaxone IV and vancomycin IV started at 2220. Droplet precautions in place.

Day 1 2310: Lumbar puncture done after the head CT; CSF sent to the laboratory.

Day 2 0545: Client had a generalized tonic-clonic seizure lasting 2 minutes: sudden stiffening of the whole body, then rhythmic jerking of all four limbs; no warning or focal movements beforehand. Now postictal: snoring respirations, opens eyes to pain only (GCS 8). Headache had been worsening overnight, with two episodes of vomiting. Pupils equal, 4 mm, sluggish.

Day 2 0730: External ventricular drain (EVD) inserted by neurosurgery; opening ICP 28 mm Hg. Draining clear, slightly cloudy CSF.

Day 12 1000: EVD removed on Day 8; repeat CT shows normal ventricles. Client alert and oriented, ambulating. No seizures since Day 2. Reports she "can't hear well" in her right ear. Dark, dry areas of skin on two toes of the left foot. Roommates and her boyfriend received ciprofloxacin on Day 2. Discharge planned for Day 14.

The nurse has reviewed the Diagnostic Results from Day 2, the Nurses' Notes from Day 2 0730 and Day 12 1000, and the Vital Signs and Laboratory Results from Day 12.

➤ For each finding, click to specify whether it shows that an **expected outcome has been met** or that it **requires follow-up**.

Finding	Expected outcome met	Requires follow-up
Alert and oriented, with no seizures since Day 2	○	○
Reduced hearing in the right ear	○	○
Normal ventricles on CT after removal of the drain	○	○
Dark, dry areas of skin on two toes	○	○
Platelet count 214 × 10^9/L and lactate 1.1 mmol/L	○	○
Close contacts received preventive antibiotics	○	○
Temperature 36.9° C and WBC 8.1 × 10^9/L	○	○

Note: Each row must have only 1 response option selected.

Submit

NURS 1017 Unit 7 Case Study 3, screen 6 (answer key, score 7/7)

Practice NCLEX RN Test

Question 6 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 6 of 6

The nurse in the emergency department is caring for a 19-year-old female university student brought in by her roommate.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results** | **Provider Orders**

Day 1 2200: Roommate states the client has had a fever, a severe headache and vomiting since this morning, and has become more confused over the last 2 hours. Client says, "My neck hurts when I bend it" and covers her eyes because the room lights hurt. Lives in a university residence; a friend on her floor had a "bad cold" last week. Unsure whether she received the school meningococcal vaccine. Drowsy, opens eyes to voice, oriented to person only (GCS 13: E3 V4 M6). Neck stiff with pain on flexion; positive Kernig and Brudzinski signs. Nonblanching petechial rash on the legs and trunk. Hands and feet cool; capillary refill 4 seconds. Pupils equal, 4 mm, reactive. Lungs clear. Abdomen soft, nontender. Last menstrual period 2 weeks ago.

Day 1 2215: Two sets of blood cultures drawn. Dexamethasone IV given at 2218, then ceftriaxone IV and vancomycin IV started at 2220. Droplet precautions in place.

Day 1 2310: Lumbar puncture done after the head CT; CSF sent to the laboratory.

Day 2 0545: Client had a generalized tonic-clonic seizure lasting 2 minutes: sudden stiffening of the whole body, then rhythmic jerking of all four limbs; no warning or focal movements beforehand. Now postictal: snoring respirations, opens eyes to pain only (GCS 8). Headache had been worsening overnight, with two episodes of vomiting. Pupils equal, 4 mm, sluggish.

Day 2 0730: External ventricular drain (EVD) inserted by neurosurgery; opening ICP 28 mm Hg. Draining clear, slightly cloudy CSF.

Day 12 1000: EVD removed on Day 8; repeat CT shows normal ventricles. Client alert and oriented, ambulating. No seizures since Day 2. Reports she "can't hear well" in her right ear. Dark, dry areas of skin on two toes of the left foot. Roommates and her boyfriend received ciprofloxacin on Day 2. Discharge planned for Day 14.

The nurse has reviewed the Diagnostic Results from Day 2, the Nurses' Notes from Day 2 0730 and Day 12 1000, and the Vital Signs and Laboratory Results from Day 12.

➤ For each finding, click to specify whether it shows that an **expected outcome has been met** or that it **requires follow-up**.

Finding	Expected outcome met	Requires follow-up
Alert and oriented, with no seizures since Day 2	●	○
Reduced hearing in the right ear	○	●
Normal ventricles on CT after removal of the drain	●	○
Dark, dry areas of skin on two toes	○	●
Platelet count 214 × 10^9/L and lactate 1.1 mmol/L	●	○
Close contacts received preventive antibiotics	●	○
Temperature 36.9° C and WBC 8.1 × 10^9/L	●	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Finish attempt →

NURS 1017 Unit 7 Case Study 4

57-year-old female client with a first seizure in adulthood from a glioblastoma: recognizing signs of increased ICP, focal deficits and seizure features, documenting the seizure type, vasogenic edema and dexamethasone adverse effects, a care plan that keeps her safe and prevents rises in ICP, status epilepticus after craniotomy, and corticosteroid effects at discharge.

Readiness problems: none

NURS 1017 Unit 7 Case Study 4, screen 1

Practice NCLEX RN Test

Question 1 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 1 of 6

The nurse reviews the Nurses' Notes and the Vital Signs. **Highlight** the findings that suggest increased intracranial pressure (ICP).

Nurses' Notes | **Vital Signs**

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 4, screen 1 (answer key, score 3/3)

Practice NCLEX RN Test

Question 1 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 1 of 6

The nurse reviews the Nurses' Notes and the Vital Signs. **Highlight** the findings that suggest increased intracranial pressure (ICP).

Nurses' Notes | **Vital Signs**

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 4, screen 2

Practice NCLEX RN Test

Question 2 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 2 of 6

The nurse in the emergency department is caring for a 57-year-old female client brought in by her husband after a seizure at home.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

The nurse has reviewed the Diagnostic Results and the Laboratory Results.

➤ For each finding, click to specify whether it is related to the **tumor's location** (a focal deficit), to **increased intracranial pressure (ICP)**, or to the **seizure**. Each finding may be related to more than one cause.

Finding	Tumor's location	Increased ICP	Seizure
Trouble finding words	☐	☐	☐
Headaches worst on waking and with coughing or bending	☐	☐	☐
Bitten tongue and urinary incontinence	☐	☐	☐
Weaker grip in the right hand	☐	☐	☐
Bilateral papilledema	☐	☐	☐
Irritable and forgetful, "not herself"	☐	☐	☐
Drowsy and slow to answer in the emergency department	☐	☐	☐

Note: Each column must have at least 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 4, screen 2 (answer key, score 8/8)

Practice NCLEX RN Test

Question 2 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 2 of 6

The nurse in the emergency department is caring for a 57-year-old female client brought in by her husband after a seizure at home.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

The nurse has reviewed the Diagnostic Results and the Laboratory Results.

➤ For each finding, click to specify whether it is related to the **tumor's location** (a focal deficit), to **increased intracranial pressure (ICP)**, or to the **seizure**. Each finding may be related to more than one cause.

Finding	Tumor's location	Increased ICP	Seizure
Trouble finding words	☑	☐	☐
Headaches worst on waking and with coughing or bending	☐	☑	☐
Bitten tongue and urinary incontinence	☐	☐	☑
Weaker grip in the right hand	☑	☐	☐
Bilateral papilledema	☐	☑	☐
Irritable and forgetful, "not herself"	☑	☐	☐
Drowsy and slow to answer in the emergency department	☐	☑	☑

Note: Each column must have at least 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 4, screen 3

Practice NCLEX RN Test

Question 3 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 3 of 6

The nurse in the emergency department is caring for a 57-year-old female client brought in by her husband after a seizure at home.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

Complete the following sentences by choosing from the lists of options.

The nurse documents the seizure the husband described as a Select... . The client's signs of increased ICP are mainly caused by Select... , which is expected to improve with Select... , so the nurse monitors her for Select... .

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 4, screen 3 (answer key, score 4/4)

Practice NCLEX RN Test

Question 3 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 3 of 6

The nurse in the emergency department is caring for a 57-year-old female client brought in by her husband after a seizure at home.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

Complete the following sentences by choosing from the lists of options.

The nurse documents the seizure the husband described as a [focal impaired awareness seizure that progressed to a bilateral tonic-clonic seizure] . The client's signs of increased ICP are mainly caused by [vasogenic edema] , which is expected to improve with [dexamethasone] , so the nurse monitors her for [hyperglycemia] .

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 4, screen 4

Practice NCLEX RN Test

Question 4 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 4 of 6

The nurse in the emergency department is caring for a 57-year-old female client brought in by her husband after a seizure at home.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

The nurse has reviewed the Provider Orders.

➤ The nurse plans care to keep the client safe and prevent increases in ICP before surgery. For each intervention, click to specify whether it is **indicated** or **contraindicated**.

Intervention	Indicated	Contraindicated
Keep the head of the bed at 30 degrees with her head in line with her body	○	○
Encourage her to cough forcefully every hour to keep her lungs clear	○	○
Give the stool softener and teach her not to strain or hold her breath during bowel movements	○	○
Keep the bed low with the side rails padded and up	○	○
Check her capillary glucose as ordered and report high readings	○	○
Lay her flat with her legs raised when she has a headache	○	○
Watch for drowsiness, dizziness, and mood changes after starting levetiracetam	○	○
Apply wrist restraints if she becomes restless	○	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 4, screen 4 (answer key, score 8/8)

Practice NCLEX RN Test

Question 4 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 4 of 6

The nurse in the emergency department is caring for a 57-year-old female client brought in by her husband after a seizure at home.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

The nurse has reviewed the Provider Orders.

➤ The nurse plans care to keep the client safe and prevent increases in ICP before surgery. For each intervention, click to specify whether it is **indicated** or **contraindicated**.

Intervention	Indicated	Contraindicated
Keep the head of the bed at 30 degrees with her head in line with her body	●	○
Encourage her to cough forcefully every hour to keep her lungs clear	○	●
Give the stool softener and teach her not to strain or hold her breath during bowel movements	●	○
Keep the bed low with the side rails padded and up	●	○
Check her capillary glucose as ordered and report high readings	●	○
Lay her flat with her legs raised when she has a headache	○	●
Watch for drowsiness, dizziness, and mood changes after starting levetiracetam	●	○
Apply wrist restraints if she becomes restless	○	●

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 4, screen 5

Practice NCLEX RN Test

Question 5 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 5 of 6

The nurse in the emergency department is caring for a 57-year-old female client brought in by her husband after a seizure at home.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results** | **Provider Orders**

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

Day 3 1500: Returned from craniotomy and subtotal resection of the left frontotemporal tumor. Awake, oriented, mild word-finding difficulty unchanged from before surgery. Dressing dry.

Day 4 0710: Called to the room by a nursing assistant: client is having a generalized tonic-clonic seizure that started at 0705 and is still continuing. Lips dusky; frothy saliva. Side rails up and padded; IV saline lock patent.

The nurse has reviewed the Nurses' Notes from Day 3 1500 and Day 4 0710 and the Provider Orders from Day 3 1500.

Which actions should the nurse take now? **Select all that apply.**

- [ ] 1. Withhold medication until the seizure has lasted 15 minutes
- [ ] 2. Apply oxygen and have suction ready
- [ ] 3. Check her capillary blood glucose
- [ ] 4. Monitor her breathing and level of sedation after the lorazepam is given
- [ ] 5. Insert a padded tongue blade between her teeth
- [ ] 6. Administer the prescribed IV lorazepam
- [ ] 7. Call for help and notify the provider
- [ ] 8. Turn the client onto her side
- [ ] 9. Restrain her arms and legs to prevent injury

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 4, screen 5 (answer key, score 6/6)

Practice NCLEX RN Test

Question 5 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 5 of 6

The nurse in the emergency department is caring for a 57-year-old female client brought in by her husband after a seizure at home.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results** | **Provider Orders**

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

Day 3 1500: Returned from craniotomy and subtotal resection of the left frontotemporal tumor. Awake, oriented, mild word-finding difficulty unchanged from before surgery. Dressing dry.

Day 4 0710: Called to the room by a nursing assistant: client is having a generalized tonic-clonic seizure that started at 0705 and is still continuing. Lips dusky; frothy saliva. Side rails up and padded; IV saline lock patent.

The nurse has reviewed the Nurses' Notes from Day 3 1500 and Day 4 0710 and the Provider Orders from Day 3 1500.

Which actions should the nurse take now? **Select all that apply.**

- [ ] 1. Withhold medication until the seizure has lasted 15 minutes
- [x] 2. Apply oxygen and have suction ready
- [x] 3. Check her capillary blood glucose
- [x] 4. Monitor her breathing and level of sedation after the lorazepam is given
- [ ] 5. Insert a padded tongue blade between her teeth
- [x] 6. Administer the prescribed IV lorazepam
- [x] 7. Call for help and notify the provider
- [x] 8. Turn the client onto her side
- [ ] 9. Restrain her arms and legs to prevent injury

Submit

← Quit

← Previous

Next →

Practice NCLEX RN Test

Question 6 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 6 of 6

The nurse in the emergency department is caring for a 57-year-old female client brought in by her husband after a seizure at home.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results** | **Provider Orders**

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

Day 3 1500: Returned from craniotomy and subtotal resection of the left frontotemporal tumor. Awake, oriented, mild word-finding difficulty unchanged from before surgery. Dressing dry.

Day 4 0710: Called to the room by a nursing assistant: client is having a generalized tonic-clonic seizure that started at 0705 and is still continuing. Lips dusky; frothy saliva. Side rails up and padded; IV saline lock patent.

Day 9 1000: No further seizures since Day 4 (levetiracetam increased to 1,000 mg every 12 hours). Headache 2/10, no vomiting. Incision clean and dry, staples intact. White patches on the tongue and inner cheeks that do not wipe off easily. Dexamethasone now 4 mg PO twice daily with a planned taper. Referred to the cancer centre for radiation and temozolomide. The nurse reviews the discharge teaching.

Day 9 1030: Client states: "I'll stop the dexamethasone once my headaches are gone." Also states: "I won't drive until my doctor tells me it's safe." Husband states: "If she has a seizure, I'll turn her on her side, time it, and call 911 if it lasts 5 minutes."

The nurse has reviewed the Diagnostic Results from Day 5, the Nurses' Notes from Day 9 1000 and 1030, and the Laboratory Results.

➤ For each finding, click to specify whether it shows that an **expected outcome has been met** or that it **requires follow-up**.

Finding	Expected outcome met	Requires follow-up
No seizures since Day 4	○	○
Capillary glucose 14.8 mmol/L before supper	○	○
Headache 2/10 with no vomiting	○	○
"I'll stop the dexamethasone once my headaches are gone."	○	○
White patches on the tongue and inner cheeks	○	○
"I won't drive until my doctor tells me it's safe."	○	○
Husband's plan for managing a seizure at home	○	○
Incision clean and dry, staples intact	○	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Finish attempt →

NURS 1017 Unit 7 Case Study 4, screen 6 (answer key, score 8/8)

Practice NCLEX RN Test

Question 6 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 6 of 6

The nurse in the emergency department is caring for a 57-year-old female client brought in by her husband after a seizure at home.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results** | **Provider Orders**

1910: Husband states they were watching television when she suddenly stopped talking and stared straight ahead, smacking her lips, with her right hand picking at her shirt; she did not answer him. About a minute later her whole body stiffened and then jerked for about 2 minutes. Bit her tongue; incontinent of urine. Over the past 6 weeks she has had headaches that are worst when she wakes up and when she coughs or bends over, and she vomited on waking twice this week without feeling nauseated. He says she has been "irritable and forgetful, not herself," and has trouble finding words. Now drowsy but rousable, oriented to person and place, slow to answer. Right hand grip slightly weaker than the left (4/5). Fundoscopic examination by the ED physician: bilateral papilledema. Point-of-care glucose 6.1 mmol/L.

Day 3 1500: Returned from craniotomy and subtotal resection of the left frontotemporal tumor. Awake, oriented, mild word-finding difficulty unchanged from before surgery. Dressing dry.

Day 4 0710: Called to the room by a nursing assistant: client is having a generalized tonic-clonic seizure that started at 0705 and is still continuing. Lips dusky; frothy saliva. Side rails up and padded; IV saline lock patent.

Day 9 1000: No further seizures since Day 4 (levetiracetam increased to 1,000 mg every 12 hours). Headache 2/10, no vomiting. Incision clean and dry, staples intact. White patches on the tongue and inner cheeks that do not wipe off easily. Dexamethasone now 4 mg PO twice daily with a planned taper. Referred to the cancer centre for radiation and temozolomide. The nurse reviews the discharge teaching.

Day 9 1030: Client states: "I'll stop the dexamethasone once my headaches are gone." Also states: "I won't drive until my doctor tells me it's safe." Husband states: "If she has a seizure, I'll turn her on her side, time it, and call 911 if it lasts 5 minutes."

The nurse has reviewed the Diagnostic Results from Day 5, the Nurses' Notes from Day 9 1000 and 1030, and the Laboratory Results.

➤ For each finding, click to specify whether it shows that an **expected outcome has been met** or that it **requires follow-up**.

Finding	Expected outcome met	Requires follow-up
No seizures since Day 4	●	○
Capillary glucose 14.8 mmol/L before supper	○	●
Headache 2/10 with no vomiting	●	○
"I'll stop the dexamethasone once my headaches are gone."	○	●
White patches on the tongue and inner cheeks	○	●
"I won't drive until my doctor tells me it's safe."	●	○
Husband's plan for managing a seizure at home	●	○
Incision clean and dry, staples intact	●	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Finish attempt →

NURS 1017 Unit 7 Case Study 5

22-year-old male client with a complete C6 spinal cord injury after diving into shallow water: recognizing breathing problems and neurogenic shock, spinal vs neurogenic vs hypovolemic shock, respiratory priority, nursing care after fusion (logrolling, suctioning, skin, bladder, vasopressor and anticoagulant monitoring), autonomic dysreflexia in rehabilitation, and teaching on prevention and lifelong self-care.

Readiness problems: none

NURS 1017 Unit 7 Case Study 5, screen 1

Practice NCLEX RN Test

Question 1 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 1 of 6

The nurse reviews the Nurses' Notes and the Vital Signs. **Highlight** the findings that require immediate follow-up.

Nurses' Notes | **Vital Signs**

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 5, screen 1 (answer key, score 7/7)

Practice NCLEX RN Test

Question 1 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 1 of 6

The nurse reviews the Nurses' Notes and the Vital Signs. **Highlight** the findings that require immediate follow-up.

Nurses' Notes | **Vital Signs**

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 5, screen 2

Practice NCLEX RN Test

Question 2 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 2 of 6

The nurse in the emergency department is caring for a 22-year-old male client who dove into shallow water at a lake.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

The nurse has reviewed the Diagnostic Results and the Laboratory Results.

➤ For each finding, click to specify whether it is consistent with **spinal shock**, **neurogenic shock**, or **hypovolemic shock**. Each finding may support more than one condition.

Finding	Spinal shock	Neurogenic shock	Hypovolemic shock
Flaccid paralysis and absent reflexes below the level of injury	☐	☐	☐
Hypotension	☐	☐	☐
Bradycardia	☐	☐	☐
Cool, clammy skin	☐	☐	☐
Loss of sympathetic nervous system outflow from an injury at T6 or above	☐	☐	☐
Warm, dry skin below the injury	☐	☐	☐
Tachycardia	☐	☐	☐
Loss of bladder and bowel reflexes, including anal sphincter tone	☐	☐	☐

Note: Each column must have at least 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 5, screen 2 (answer key, score 9/9)

Practice NCLEX RN Test

Question 2 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 2 of 6

The nurse in the emergency department is caring for a 22-year-old male client who dove into shallow water at a lake.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

The nurse has reviewed the Diagnostic Results and the Laboratory Results.

➤ For each finding, click to specify whether it is consistent with **spinal shock**, **neurogenic shock**, or **hypovolemic shock**. Each finding may support more than one condition.

Finding	Spinal shock	Neurogenic shock	Hypovolemic shock
Flaccid paralysis and absent reflexes below the level of injury	☑	☐	☐
Hypotension	☐	☑	☑
Bradycardia	☐	☑	☐
Cool, clammy skin	☐	☐	☑
Loss of sympathetic nervous system outflow from an injury at T6 or above	☐	☑	☐
Warm, dry skin below the injury	☐	☑	☐
Tachycardia	☐	☐	☑
Loss of bladder and bowel reflexes, including anal sphincter tone	☑	☐	☐

Note: Each column must have at least 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 5, screen 3

Practice NCLEX RN Test

Question 3 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 3 of 6

The nurse in the emergency department is caring for a 22-year-old male client who dove into shallow water at a lake.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results**

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

Day 1 1630: Vital capacity 1.1 L (predicted 4.8 L). Using neck muscles to breathe; says he feels "tired of breathing." Oxygen 4 L/min by nasal cannula applied.

The nurse has reviewed the Nurses' Notes from Day 1 1630 and the Vital Signs from Day 1 1630.

➤ Complete the following sentences by choosing from the lists of options.

The client's priority problem is Select... . His injury spares Select... but paralyzes Select... . His breathing may get worse over the next few days because Select... .

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 5, screen 3 (answer key, score 4/4)

Practice NCLEX RN Test

Question 3 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 3 of 6

The nurse in the emergency department is caring for a 22-year-old male client who dove into shallow water at a lake.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

Day 1 1630: Vital capacity 1.1 L (predicted 4.8 L). Using neck muscles to breathe; says he feels "tired of breathing." Oxygen 4 L/min by nasal cannula applied.

The nurse has reviewed the Nurses' Notes from Day 1 1630 and the Vital Signs from Day 1 1630.

➤ Complete the following sentences by choosing from the lists of options.

The client's priority problem is [risk for respiratory failure] . His injury spares [the phrenic nerve (C3–C5), which controls the diaphragm] but paralyzes [the intercostal and abdominal muscles] . His breathing may get worse over the next few days because [edema of the spinal cord can extend upward toward C3–C5] .

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 5, screen 4

Practice NCLEX RN Test

Question 4 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 4 of 6

The nurse in the emergency department is caring for a 22-year-old male client who dove into shallow water at a lake.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

Day 1 1630: Vital capacity 1.1 L (predicted 4.8 L). Using neck muscles to breathe; says he feels "tired of breathing." Oxygen 4 L/min by nasal cannula applied.

Day 2 0800: Underwent anterior cervical decompression and fusion of C5–C6 overnight; in a hard cervical collar. Remains on 4 L/min oxygen; vital capacity 1.3 L. Admitted to the ICU.

The nurse has reviewed the Nurses' Notes from Day 2 0800 and the Provider Orders.

➤ For each intervention, click to specify whether it is **indicated** or **contraindicated** for the client.

Intervention	Indicated	Contraindicated
Logroll the client with a second nurse holding the head in line with the body	○	○
Suction the airway for longer than 15 seconds to clear thick secretions	○	○
Measure the vital capacity and assess cough strength as ordered	○	○
Raise the head of the bed quickly to 90 degrees to ease his breathing	○	○
Check the norepinephrine IV site often for swelling, coolness, or pale skin	○	○
Remove the cervical collar to wash his neck while he lies alone	○	○
Turn him at least every 2 hours and inspect the skin over bony prominences	○	○
Monitor urine output through the indwelling catheter	○	○
Watch for bleeding gums, blood in the urine, and bruising at injection sites once enoxaparin starts	○	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 5, screen 4 (answer key, score 9/9)

Practice NCLEX RN Test

Question 4 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 4 of 6

The nurse in the emergency department is caring for a 22-year-old male client who dove into shallow water at a lake.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

Day 1 1630: Vital capacity 1.1 L (predicted 4.8 L). Using neck muscles to breathe; says he feels "tired of breathing." Oxygen 4 L/min by nasal cannula applied.

Day 2 0800: Underwent anterior cervical decompression and fusion of C5–C6 overnight; in a hard cervical collar. Remains on 4 L/min oxygen; vital capacity 1.3 L. Admitted to the ICU.

The nurse has reviewed the Nurses' Notes from Day 2 0800 and the Provider Orders.

➤ For each intervention, click to specify whether it is **indicated** or **contraindicated** for the client.

Intervention	Indicated	Contraindicated
Logroll the client with a second nurse holding the head in line with the body	●	○
Suction the airway for longer than 15 seconds to clear thick secretions	○	●
Measure the vital capacity and assess cough strength as ordered	●	○
Raise the head of the bed quickly to 90 degrees to ease his breathing	○	●
Check the norepinephrine IV site often for swelling, coolness, or pale skin	●	○
Remove the cervical collar to wash his neck while he lies alone	○	●
Turn him at least every 2 hours and inspect the skin over bony prominences	●	○
Monitor urine output through the indwelling catheter	●	○
Watch for bleeding gums, blood in the urine, and bruising at injection sites once enoxaparin starts	●	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

Practice NCLEX RN Test

Question 5 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 5 of 6

The nurse in the emergency department is caring for a 22-year-old male client who dove into shallow water at a lake.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

Day 1 1630: Vital capacity 1.1 L (predicted 4.8 L). Using neck muscles to breathe; says he feels "tired of breathing." Oxygen 4 L/min by nasal cannula applied.

Day 2 0800: Underwent anterior cervical decompression and fusion of C5–C6 overnight; in a hard cervical collar. Remains on 4 L/min oxygen; vital capacity 1.3 L. Admitted to the ICU.

Week 5 1400: Rehabilitation unit. Spinal shock resolved in Week 3; reflexes and spasticity in the legs have returned. Client lying in bed, reports a sudden pounding headache and blurred vision. Face and neck flushed and sweating; skin below the clavicles pale and cool with goosebumps. Nasal stuffiness. Wearing an abdominal binder and leg straps for the urine drainage bag. Indwelling catheter bag has had no new urine for 3 hours.

The nurse has reviewed the Nurses' Notes from Week 5 1400 and the Vital Signs from Week 5 1400.

Place the nurse's actions in the order in which they should be performed. Arrange from first to last.

UNORDERED OPTIONS	YOUR ORDERED STEPS
If urine does not drain, replace the catheter using lidocaine jelly	
Sit the client upright and lower his legs	
Check the catheter tubing for kinks or blockage	
Loosen the abdominal binder and the leg straps	
If the BP stays high after the bladder is	

→ ← ↑ ↓

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 5, screen 5 (answer key, score 1/1)

Practice NCLEX RN Test

Question 5 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 5 of 6

The nurse in the emergency department is caring for a 22-year-old male client who dove into shallow water at a lake.

Nurses' Notes | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

Day 1 1630: Vital capacity 1.1 L (predicted 4.8 L). Using neck muscles to breathe; says he feels "tired of breathing." Oxygen 4 L/min by nasal cannula applied.

Day 2 0800: Underwent anterior cervical decompression and fusion of C5–C6 overnight; in a hard cervical collar. Remains on 4 L/min oxygen; vital capacity 1.3 L. Admitted to the ICU.

Week 5 1400: Rehabilitation unit. Spinal shock resolved in Week 3; reflexes and spasticity in the legs have returned. Client lying in bed, reports a sudden pounding headache and blurred vision. Face and neck flushed and sweating; skin below the clavicles pale and cool with goosebumps. Nasal stuffiness. Wearing an abdominal binder and leg straps for the urine drainage bag. Indwelling catheter bag has had no new urine for 3 hours.

The nurse has reviewed the Nurses' Notes from Week 5 1400 and the Vital Signs from Week 5 1400.

Place the nurse's actions in the order in which they should be performed. Arrange from first to last.

UNORDERED OPTIONS	YOUR ORDERED STEPS
	1. Sit the client upright and lower his legs
	2. Loosen the abdominal binder and the leg straps
	3. Check the catheter tubing for kinks or blockage
	4. If urine does not drain, replace the catheter using lidocaine jelly
	5. If the BP stays high after the bladder is

Submit

← Quit

← Previous

Next →

NOT COMPLETE

Case Study Screen 6 of 6

The nurse in the emergency department is caring for a 22-year-old male client who dove into shallow water at a lake.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results** | **Provider Orders**

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

Day 1 1630: Vital capacity 1.1 L (predicted 4.8 L). Using neck muscles to breathe; says he feels "tired of breathing." Oxygen 4 L/min by nasal cannula applied.

Day 2 0800: Underwent anterior cervical decompression and fusion of C5–C6 overnight; in a hard cervical collar. Remains on 4 L/min oxygen; vital capacity 1.3 L. Admitted to the ICU.

Week 5 1400: Rehabilitation unit. Spinal shock resolved in Week 3; reflexes and spasticity in the legs have returned. Client lying in bed, reports a sudden pounding headache and blurred vision. Face and neck flushed and sweating; skin below the clavicles pale and cool with goosebumps. Nasal stuffiness. Wearing an abdominal binder and leg straps for the urine drainage bag. Indwelling catheter bag has had no new urine for 3 hours.

Week 5 1420: Tubing kinked under his thigh; straightened, 900 mL of urine drained. BP 128/76 mm Hg at 1415, 104/64 mm Hg at 1425. Headache resolving.

Week 10 1000: Discharge planning. The nurse reviews teaching with the client and his brother, who will help with his care at home.

The nurse has reviewed the Nurses' Notes from Week 5 1420 and Week 10 1000.

➤ For each statement, click to specify whether it indicates an **understanding** of the teaching or a **need for further teaching**.

Statement	Understanding	Needs further teaching
Client: "If I get a pounding headache and my face flushes, I'll sit up right away and check my catheter."	○	○
Brother: "When his blood pressure goes up, he should lie down flat."	○	○
Client: "Autonomic dysreflexia only happens in the first few months after an injury."	○	○
Client: "I'll carry my autonomic dysreflexia wallet card and show it to staff if I go to emergency."	○	○
Client: "I'll do my bowel program at the same time every day."	○	○
Brother: "Since he can't feel his legs, we don't need to check his skin."	○	○
Brother: "Next summer, everyone will check the water depth and go in feet first the first time."	○	○

Note: Each row must have only 1 response option selected.

Submit

NURS 1017 Unit 7 Case Study 5, screen 6 (answer key, score 7/7)

Practice NCLEX RN Test

Question 6 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 6 of 6

The nurse in the emergency department is caring for a 22-year-old male client who dove into shallow water at a lake.

Nurses' Notes | **Vital Signs** | **Diagnostic Results** | **Laboratory Results** | **Provider Orders**

Day 1 1430: Dove headfirst off a dock into about 1 metre of water at 1340 and struck his head on the bottom. Friends pulled him out of the water, keeping his head in line with his body. Arrived by ambulance with a cervical collar and spinal motion restriction. Alert and oriented × 4. States, "I can't feel or move my legs." Can shrug his shoulders and bend his elbows; cannot straighten his elbows or move his fingers. No sensation below the clavicles or in the arms and hands except the outer arms and thumbs. Legs flaccid with no reflexes; no anal sphincter tone. Breathing is diaphragmatic: the abdomen rises while the chest barely moves; speaks in short sentences; weak cough. Skin warm, dry and pink below the injury. 3-cm abrasion on the forehead.

Day 1 1630: Vital capacity 1.1 L (predicted 4.8 L). Using neck muscles to breathe; says he feels "tired of breathing." Oxygen 4 L/min by nasal cannula applied.

Day 2 0800: Underwent anterior cervical decompression and fusion of C5–C6 overnight; in a hard cervical collar. Remains on 4 L/min oxygen; vital capacity 1.3 L. Admitted to the ICU.

Week 5 1400: Rehabilitation unit. Spinal shock resolved in Week 3; reflexes and spasticity in the legs have returned. Client lying in bed, reports a sudden pounding headache and blurred vision. Face and neck flushed and sweating; skin below the clavicles pale and cool with goosebumps. Nasal stuffiness. Wearing an abdominal binder and leg straps for the urine drainage bag. Indwelling catheter bag has had no new urine for 3 hours.

Week 5 1420: Tubing kinked under his thigh; straightened, 900 mL of urine drained. BP 128/76 mm Hg at 1415, 104/64 mm Hg at 1425. Headache resolving.

Week 10 1000: Discharge planning. The nurse reviews teaching with the client and his brother, who will help with his care at home.

The nurse has reviewed the Nurses' Notes from Week 5 1420 and Week 10 1000.

➤ For each statement, click to specify whether it indicates an **understanding** of the teaching or a **need for further teaching**.

Statement	Understanding	Needs further teaching
Client: "If I get a pounding headache and my face flushes, I'll sit up right away and check my catheter."	●	○
Brother: "When his blood pressure goes up, he should lie down flat."	○	●
Client: "Autonomic dysreflexia only happens in the first few months after an injury."	○	●
Client: "I'll carry my autonomic dysreflexia wallet card and show it to staff if I go to emergency."	●	○
Client: "I'll do my bowel program at the same time every day."	●	○
Brother: "Since he can't feel his legs, we don't need to check his skin."	○	●
Brother: "Next summer, everyone will check the water depth and go in feet first the first time."	●	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Finish attempt →

NURS 1017 Unit 7 Case Study 6

76-year-old male client with Parkinson disease taking apixaban who develops delirium from a chronic subdural hematoma weeks after a fall: recognizing the hallmark signs of Parkinson disease (with ALS and MS as contrasts), delirium vs dementia and the types of dementia, delirium-safe and fall-safe nursing care, missed levodopa doses (parkinsonism-hyperpyrexia syndrome), and discharge teaching including levodopa adverse effects.

Readiness problems: none

NURS 1017 Unit 7 Case Study 6, screen 1

Practice NCLEX RN Test

Question 1 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 1 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | **History and Physical** | **Vital Signs**

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

The nurse reviews the client's history and assessment. Which of the following findings would be consistent with **Parkinson disease**? **Select all that apply.**

- [ ] 1. New weakness of the left arm with a downward drift
- [ ] 2. Inability to name the days of the week backward
- [ ] 3. Muscle twitching (fasciculations) and wasting
- [ ] 4. Masklike face and a soft, monotone voice
- [ ] 5. Resting tremor of the hand that stops with purposeful movement
- [ ] 6. Shuffling gait with reduced arm swing
- [ ] 7. A drop in BP from 138/78 mm Hg lying to 108/66 mm Hg standing
- [ ] 8. Cogwheel rigidity
- [ ] 9. Slow, small movements (bradykinesia)
- [ ] 10. Tremor that worsens as the hand approaches a target

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 6, screen 1 (answer key, score 6/6)

Practice NCLEX RN Test

Question 1 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 1 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | **History and Physical** | **Vital Signs**

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

The nurse reviews the client's history and assessment. Which of the following findings would be consistent with **Parkinson disease**? **Select all that apply.**

- [ ] 1. New weakness of the left arm with a downward drift
- [ ] 2. Inability to name the days of the week backward
- [ ] 3. Muscle twitching (fasciculations) and wasting
- [x] 4. Masklike face and a soft, monotone voice
- [x] 5. Resting tremor of the hand that stops with purposeful movement
- [x] 6. Shuffling gait with reduced arm swing
- [x] 7. A drop in BP from 138/78 mm Hg lying to 108/66 mm Hg standing
- [x] 8. Cogwheel rigidity
- [x] 9. Slow, small movements (bradykinesia)
- [ ] 10. Tremor that worsens as the hand approaches a target

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 6, screen 2

Practice NCLEX RN Test

Question 2 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 2 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | **History and Physical** | **Vital Signs**

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

➤ The client's wife asks whether her husband is "getting dementia." For each feature, click to specify whether it is more consistent with **delirium** or **dementia** (major neurocognitive disorder).

Feature	Delirium	Dementia
Develops over hours to days	○	○
Gradual, progressive decline over months to years	○	○
Inattention, such as being unable to name the days of the week backward	○	○
Alert, with attention preserved until the late stages	○	○
Level of consciousness and confusion fluctuate during the day	○	○
Caused by the progressive loss of neurons, as in Alzheimer disease	○	○
Often reversible when the underlying cause is treated	○	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 6, screen 2 (answer key, score 7/7)

Practice NCLEX RN Test

Question 2 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 2 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | **History and Physical** | **Vital Signs**

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

➤ The client's wife asks whether her husband is "getting dementia." For each feature, click to specify whether it is more consistent with **delirium** or **dementia** (major neurocognitive disorder).

Feature	Delirium	Dementia
Develops over hours to days	●	○
Gradual, progressive decline over months to years	○	●
Inattention, such as being unable to name the days of the week backward	●	○
Alert, with attention preserved until the late stages	○	●
Level of consciousness and confusion fluctuate during the day	●	○
Caused by the progressive loss of neurons, as in Alzheimer disease	○	●
Often reversible when the underlying cause is treated	●	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 6, screen 3

Practice NCLEX RN Test

Question 3 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 3 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | History and Physical | Vital Signs | Diagnostic Results | Laboratory Results

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

The nurse has reviewed the Diagnostic Results and the Laboratory Results.

Complete the following sentences by choosing from the lists of options.

The client's acute confusion is most likely Select... caused by Select... . This type of bleed comes from Select... , and his risk was increased by Select... .

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 6, screen 3 (answer key, score 4/4)

Practice NCLEX RN Test

Question 3 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 3 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | History and Physical | Vital Signs | Diagnostic Results | Laboratory Results

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

The nurse has reviewed the Diagnostic Results and the Laboratory Results.

➤ Complete the following sentences by choosing from the lists of options.

The client's acute confusion is most likely [delirium] caused by [a chronic subdural hematoma]. This type of bleed comes from [torn bridging veins], and his risk was increased by [brain atrophy, frequent falls, and anticoagulant therapy].

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 6, screen 4

Practice NCLEX RN Test

Question 4 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 4 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | History and Physical | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

The nurse has reviewed the Provider Orders.

➤ For each intervention, click to specify whether it is **indicated** or **contraindicated** for the client.

Intervention	Indicated	Contraindicated
Give the carbidopa-levodopa at the exact scheduled times, including while he is NPO	○	○
Give the PRN haloperidol if he becomes agitated	○	○
Reorient him with a clock and calendar, his glasses and hearing aids, and his wife's presence	○	○
Apply wrist restraints at night so he does not get out of bed	○	○
Have him sit and dangle his legs before standing, with assistance	○	○
Keep the room brightly lit and the television on all night to keep him oriented	○	○
Perform the ordered neurological checks, including pupils and arm strength	○	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 6, screen 4 (answer key, score 7/7)

Practice NCLEX RN Test

Question 4 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 4 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | History and Physical | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

The nurse has reviewed the Provider Orders.

➤ For each intervention, click to specify whether it is **indicated** or **contraindicated** for the client.

Intervention	Indicated	Contraindicated
Give the carbidopa-levodopa at the exact scheduled times, including while he is NPO	●	○
Give the PRN haloperidol if he becomes agitated	○	●
Reorient him with a clock and calendar, his glasses and hearing aids, and his wife's presence	●	○
Apply wrist restraints at night so he does not get out of bed	○	●
Have him sit and dangle his legs before standing, with assistance	●	○
Keep the room brightly lit and the television on all night to keep him oriented	○	●
Perform the ordered neurological checks, including pupils and arm strength	●	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Next →

Practice NCLEX RN Test

Question 5 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 5 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | History and Physical | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

Day 2 1400: Surgery postponed to 1600 because of an emergency case. The 0700 and 1100 carbidopa-levodopa doses were not given because the client was NPO. Client now rigid throughout and unable to turn in bed or swallow his saliva well; more confused; skin hot and sweaty. Pupils equal and reactive; left arm drift unchanged.

The nurse has reviewed the Nurses' Notes from Day 2 1400, the Vital Signs from Day 2 1400 and the Laboratory Results from Day 2 1415.

Which actions should the nurse take? **Select all that apply.**

- [ ] 1. Ask the provider how to give the carbidopa-levodopa now (for example, by nasogastric tube)
- [ ] 2. Give the PRN haloperidol for the increased confusion
- [ ] 3. Apply cooling measures
- [ ] 4. Encourage him to drink water to cool down
- [ ] 5. Monitor urine output and colour
- [ ] 6. Notify the provider of the missed doses and the client's condition
- [ ] 7. Continue to hold the carbidopa-levodopa until after surgery

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 6, screen 5 (answer key, score 4/4)

Practice NCLEX RN Test

Question 5 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 5 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | History and Physical | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

Day 2 1400: Surgery postponed to 1600 because of an emergency case. The 0700 and 1100 carbidopa-levodopa doses were not given because the client was NPO. Client now rigid throughout and unable to turn in bed or swallow his saliva well; more confused; skin hot and sweaty. Pupils equal and reactive; left arm drift unchanged.

The nurse has reviewed the Nurses' Notes from Day 2 1400, the Vital Signs from Day 2 1400 and the Laboratory Results from Day 2 1415.

Which actions should the nurse take? **Select all that apply.**

- [x] 1. Ask the provider how to give the carbidopa-levodopa now (for example, by nasogastric tube)
- [ ] 2. Give the PRN haloperidol for the increased confusion
- [x] 3. Apply cooling measures
- [ ] 4. Encourage him to drink water to cool down
- [x] 5. Monitor urine output and colour
- [x] 6. Notify the provider of the missed doses and the client's condition
- [ ] 7. Continue to hold the carbidopa-levodopa until after surgery

Submit

← Quit

← Previous

Next →

NURS 1017 Unit 7 Case Study 6, screen 6

Practice NCLEX RN Test

Question 6 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 6 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | History and Physical | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

Day 2 1400: Surgery postponed to 1600 because of an emergency case. The 0700 and 1100 carbidopa-levodopa doses were not given because the client was NPO. Client now rigid throughout and unable to turn in bed or swallow his saliva well; more confused; skin hot and sweaty. Pupils equal and reactive; left arm drift unchanged.

Day 3 0900: Levodopa restarted by NG tube at 1530 on Day 2; rigidity and fever settled overnight. Burr-hole drainage of the subdural hematoma done at 0730 today. CK falling.

Day 8 1000: Alert, oriented, attentive (names the months backward). Headache gone; left arm strength equal to the right. Taking carbidopa-levodopa on time by mouth. Discharge home tomorrow with a home care referral. Apixaban to stay on hold until the neurosurgery follow-up. The nurse reviews the discharge teaching with the client and his wife.

The nurse has reviewed the Nurses' Notes from Day 3 0900 and Day 8 1000 and the Vital Signs from Day 8 1000.

➤ For each statement, click to specify whether it indicates an **understanding** of the teaching or a **need for further teaching**.

Statement	Understanding	Needs further teaching
Wife: "His confusion cleared once the blood was drained, so it was delirium, not dementia."	○	○
Client: "I'll take my levodopa with my high-protein breakfast to settle my stomach."	○	○
Wife: "If he gets a new headache or becomes sleepy or confused again, I'll take him to emergency."	○	○
Client: "Now that my head is better, I can restart my apixaban tonight."	○	○
Wife: "His mother had Alzheimer disease, so his confusion means he has it too."	○	○
Wife: "I'll remove the throw rugs and keep a night light on."	○	○
Wife: "If he gets new jerky or writhing movements, sees things that aren't there, or gets dizzy when he stands, I'll tell his doctor."	○	○
Wife: "If he is in hospital again, I'll ask that his Parkinson pills be given on time."	○	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Finish attempt →

NURS 1017 Unit 7 Case Study 6, screen 6 (answer key, score 8/8)

Practice NCLEX RN Test

Question 6 of 6

Calculator

Question Navigator

NOT COMPLETE

Case Study Screen 6 of 6

The nurse in the emergency department is caring for a 76-year-old male client with Parkinson disease brought in by his wife.

Nurses' Notes | History and Physical | Vital Signs | Diagnostic Results | Laboratory Results | Provider Orders

Day 1 1400: Wife states that over the last 10 days he has become increasingly confused and sleepy and has had daily headaches. "Some hours he's his usual self, and other times he doesn't know where he is." Up at night, pulling at his clothes. Four weeks ago he fell in the kitchen and hit his head on a cupboard door, without losing consciousness. Before this he managed his own medications and finances. Assessment: drowsy, easily distracted, cannot name the days of the week backward; oriented to person only. Resting tremor of the right hand ("pill-rolling") that stops when he reaches for a cup; cogwheel rigidity in both arms; slow, small movements; shuffling gait with reduced arm swing (uses a walker); masklike face; soft, monotone voice. New mild weakness of the left arm with a downward drift. No muscle twitching or wasting. Last bowel movement 4 days ago.

Day 2 1400: Surgery postponed to 1600 because of an emergency case. The 0700 and 1100 carbidopa-levodopa doses were not given because the client was NPO. Client now rigid throughout and unable to turn in bed or swallow his saliva well; more confused; skin hot and sweaty. Pupils equal and reactive; left arm drift unchanged.

Day 3 0900: Levodopa restarted by NG tube at 1530 on Day 2; rigidity and fever settled overnight. Burr-hole drainage of the subdural hematoma done at 0730 today. CK falling.

Day 8 1000: Alert, oriented, attentive (names the months backward). Headache gone; left arm strength equal to the right. Taking carbidopa-levodopa on time by mouth. Discharge home tomorrow with a home care referral. Apixaban to stay on hold until the neurosurgery follow-up. The nurse reviews the discharge teaching with the client and his wife.

The nurse has reviewed the Nurses' Notes from Day 3 0900 and Day 8 1000 and the Vital Signs from Day 8 1000.

➤ For each statement, click to specify whether it indicates an **understanding** of the teaching or a **need for further teaching**.

Statement	Understanding	Needs further teaching
Wife: "His confusion cleared once the blood was drained, so it was delirium, not dementia."	●	○
Client: "I'll take my levodopa with my high-protein breakfast to settle my stomach."	○	●
Wife: "If he gets a new headache or becomes sleepy or confused again, I'll take him to emergency."	●	○
Client: "Now that my head is better, I can restart my apixaban tonight."	○	●
Wife: "His mother had Alzheimer disease, so his confusion means he has it too."	○	●
Wife: "I'll remove the throw rugs and keep a night light on."	●	○
Wife: "If he gets new jerky or writhing movements, sees things that aren't there, or gets dizzy when he stands, I'll tell his doctor."	●	○
Wife: "If he is in hospital again, I'll ask that his Parkinson pills be given on time."	●	○

Note: Each row must have only 1 response option selected.

Submit

← Quit

← Previous

Finish attempt →