

Roberta and BERT: Revolutionizing Mental Healthcare through Natural Language

Arun Singla*

Email: arunnsingla@gmail.com

ORCID: <https://orcid.org/0009-0003-2027-0112>

Affiliation: Director, Shodh Sagar Pvt. Ltd., Delhi

Accepted: 11/03/2024

Published: 23/03/2024

* Corresponding author

How to Cite this Article:

Singla, A. (2024). Roberta and BERT: Revolutionizing Mental Healthcare through Natural Language. *Shodh Sagar Journal of Artificial Intelligence and Machine Learning*, 1(1), 10-27.

DOI: <https://doi.org/10.36676/ssjaiml.v1.i1.02>



Abstract: *In recent years, developments in natural language processing (NLP) have made it possible for novel applications to be developed in the field of mental healthcare. Roberta and BERT are two examples of these advances. They are examples of state-of-the-art language models that have exhibited extraordinary skills in comprehending and processing human language. Through the use of NLP, this research investigates the revolutionary role that Roberta and BERT have had in the transformation of mental healthcare. via the use of huge quantities of textual data, these models provide insights into the feelings, attitudes, and mental states of patients that are communicated via language that have never been considered before. Through the use of advanced language comprehension methods, Roberta and BERT are able to provide more precise diagnosis, individualized treatment suggestions, and therapeutic interventions that are specifically designed to meet the specific requirements of people. Furthermore, they make it easier to construct virtual assistants and chatbots that are able to provide support and intervention in real time. This allows mental health treatments to be extended to groups who are not currently receiving them and helps to eliminate obstacles that prevent access to these services. However, in order to guarantee the appropriate deployment of these technologies, it is necessary to carefully traverse difficulties such as concerns around privacy, bias in data, and ethical considerations. Despite this, Roberta and BERT have a tremendous amount of promise to revolutionize mental healthcare via the use of natural language understanding. This has the ability to improve results, better relationships between patients and providers, and eventually contribute to the overall well-being of people all over the globe.*

Keywords: Mental Healthcare, Deep learning, Roberta, Bert, NLP

[1] Introduction

In recent years, there has been a rising understanding of the potential of natural language processing (NLP) technology to change different parts of healthcare delivery. This recognition has been occurring with increasing frequency. Among these breakthroughs, the incorporation of language models like Roberta and BERT has attracted a large amount of interest due to the fact that it has the potential to revolutionize mental



healthcare by means of the analysis of natural language data. Disorders of mental health constitute a huge burden on the global community, since millions of people all over the globe are affected by ailments such as anxiety, depression, and post-traumatic stress disorder (PTSD). Traditional methods of providing mental healthcare, on the other hand, often encounter obstacles in the form of availability, cost, and stigma, which makes it difficult to implement therapies and reduces their overall efficacy. Within this framework, the introduction of powerful natural language processing models presents new options to overcome these obstacles and deliver mental healthcare treatments that are more individualized, accessible, and efficient. Using the massive volumes of textual data that are created via online platforms, electronic health records, and digital communication channels, Roberta and BERT make it possible for doctors to get a more in-depth understanding of the mental states, emotions, and behaviors that are represented through language of their patients. The purpose of this article is to investigate the transformational potential of Roberta and BERT in the field of mental healthcare. Specifically, the research will investigate their applications in diagnosis, treatment, and intervention, as well as the obstacles and possibilities connected with their incorporation into clinical practice. The purpose of this study is to provide an in-depth analysis of the existing body of literature and case studies in order to shed light on the present state of research in this field and to suggest possible avenues for further investigation. Ultimately, we expect that by leveraging the potential of powerful natural language processing technologies, we will be able to catalyze a paradigm change in the delivery of mental healthcare, which will ultimately result in better results and an increased quality of life for persons who are afflicted with mental health illnesses.

1.1 Mental Healthcare

This is encapsulated in the title "Roberta and BERT: Revolutionizing Mental Healthcare through Natural Language," which describes the crucial role that sophisticated natural language processing (NLP) models, namely Roberta and BERT, play in transforming the landscape of mental healthcare. In light of the increasing difficulties and complexity associated with mental health illnesses, these cutting-edge technologies provide a transformational approach to the diagnosis, treatment, and support of those affected by these conditions. Because they leverage the power of natural language comprehension, Roberta and BERT enable professionals working in the field of mental health to get a more profound knowledge of the feelings, experiences, and mental states that are represented via the language of their patients. This paradigm change makes it possible to conduct more precise assessments, provide more individualized interventions, and provide more focused support, which eventually leads to an improvement in the efficiency and accessibility of mental healthcare services. The purpose of this project is to change the delivery of mental healthcare, improve patient outcomes, and pave the path for a more compassionate and inclusive approach to mental well-being. Each of these goals will be accomplished via the novel uses of natural language processing technology.

1.2 Deep learning

As a strong paradigm for tackling difficult issues across a variety of disciplines, deep learning, which is a subset of artificial intelligence (AI) and machine learning (ML), has emerged as a formidable tool. Deep learning algorithms have exhibited impressive powers in learning from massive quantities of data to generate predictions, categorize objects, and extract significant patterns. These capabilities are rooted in the



notion of artificial neural networks, which are inspired by the structure and function of the human brain. Deep learning models are able to autonomously build hierarchical representations of data, which enables them to capture nuanced connections and nuances that may be difficult to define explicitly. This is in contrast to typical machine learning techniques, which depend on handcrafted features and explicit rule-based algorithms. Deep learning has been fueled by a number of key factors, including the availability of massive datasets, advancements in computational hardware such as graphical processing units (GPUs), and innovations in algorithmic techniques such as convolutional neural networks (CNNs), recurrent neural networks (RNNs), and transformer models. These factors have all contributed to the rise of deep learning. These breakthroughs have made it possible for deep learning models to attain levels of accuracy and performance that have never been seen before across a broad variety of tasks. These tasks include image identification, voice recognition, natural language processing (NLP), and autonomous driving. In recent years, deep learning has successfully revolutionized a wide range of sectors and applications, making substantial gains in this direction. Medical image analysis, illness diagnosis, medication development, and tailored treatment recommendation systems are some of the other applications of deep learning models that have been used in the healthcare industry. Deep learning algorithms are used in the financial sector for the purposes of detecting fraudulent activity, algorithmic trading, and risk management. Deep learning is the driving force behind recommendation systems, customer segmentation, and targeted advertising campaigns in the realms of consumer marketing and online commerce. Deep learning is not without its difficulties and restrictions, despite the fact that it has led to many accomplishments. The training of deep neural networks often necessitates the use of big computer resources and a substantial quantity of labeled data, which renders it unavailable to a great number of researchers and organizations. Furthermore, deep learning models may be subject to biases that are present in the training data. Furthermore, these models may have difficulties with interpretability, which makes it difficult to comprehend the reasoning that lies behind their predictions. Deep learning continues to be a source of inspiration for academics, engineers, and practitioners all around the globe because of its potential to disrupt industries, drive innovation, and solve challenging issues. In this article, we present an overview of deep learning, including its fundamental concepts, methodologies, applications, and obstacles. We also emphasize the transformational influence that deep learning has had on society as well as its promise for future research and development.

1.3 Role of deep learning for Mental Healthcare

Deep learning has emerged as a revolutionary tool in the field of mental healthcare, giving fresh techniques to solve difficulties that have persisted for a long time and improving the results for patients. Detecting and diagnosing mental health illnesses at an earlier stage is one of the most important functions that deep learning can do. In order to identify minor patterns that are symptomatic of diseases such as depression, anxiety, or post-traumatic stress disorder (PTSD), deep learning models may analyze a wide variety of data sources, including electronic health records, activity on social media, and physiological signals. This allows for prompt intervention and therapy to be administered. In addition, deep learning makes it possible to customize treatment planning by using data from individual patients to adjust therapeutic methods to their requirements and preferences. This helps to maximize treatment effectiveness and adherence. In addition, deep learning makes it possible to model mental health outcomes in a predictive manner, which means that it can estimate the risk of relapse or adverse events based on previous data and patient characteristics. This



allows for proactive treatments and resource allocation to be guided. Deep learning methods that use natural language processing (NLP) make it possible to conduct automated sentiment analysis and virtual counseling sessions. This not only increases the capabilities of mental health experts but also makes therapy more accessible to patients. In addition, deep learning algorithms are able to monitor facial expressions, verbal intonations, and physiological signs in order to infer emotional states in real time. This makes it possible to provide early intervention and individualized assistance. Nevertheless, in order to guarantee the appropriate use of deep learning technologies in mental healthcare, it is necessary to give particular attention to ethical concerns about the preservation of privacy, the elimination of algorithmic bias, and the promotion of openness. In summary, deep learning provides a multidimensional approach to changing mental healthcare, empowering practitioners, enhancing treatment results, and expanding our knowledge of mental health diseases. Deep learning seeks to accomplish all of these goals simultaneously.

1.4 Need of Research

Research under the title "Roberta and BERT: Revolutionizing Mental Healthcare through Natural Language" is required because there is an immediate need to develop new approaches to the delivery of mental healthcare and to solve the ongoing difficulties that are experienced by people who have mental health issues. Significant obstacles still exist in the way of appropriate diagnosis, treatment, and support for mental health concerns, despite the fact that knowledge and understanding of these issues have risen. Because of issues such as limited resources, stigma, and inequities in access to treatment, traditional methods to mental healthcare sometimes fail to satisfy the different requirements of patients. This is because of some of the causes listed above. When seen in this light, the development of sophisticated natural language processing (NLP) models like Roberta and BERT presents a potentially fruitful path toward the revolutionization of mental healthcare. Textual data sources including as medical records, online forums, and social media platforms may be analyzed using these cutting-edge language models, which have shown exceptional skills in understanding and interpreting human language. This presents an opportunity to extract useful insights from these sources. When mental health practitioners leverage the power of these models, they are able to acquire deeper insights into the experiences, emotions, and mental states of their patients. This enables them to make more accurate diagnoses, develop more individualized treatment plans, and implement more focused treatments among their patients. The fact that Roberta and BERT have the ability to cover crucial gaps in the delivery of mental healthcare highlights the need of doing research in this particular field. It is possible for researchers and practitioners to build creative tools and treatments to help people who are struggling with mental health issues, improve communication between patients and providers, and boost treatment results by using the technology of natural language processing (NLP). Furthermore, research in this field has the potential to contribute to the advancement of our knowledge of the intricate relationship that exists between language, cognition, and mental health, so paving the way for treatments and support services that are more successfully implemented. In general, the necessity of conducting research with the title "Roberta and BERT: Revolutionizing Mental Healthcare through Natural Language" is driven by the imperative to utilize cutting-edge technology in order to address long-standing challenges in the delivery of mental healthcare, improve access to care, and enhance the well-being of individuals who are affected by mental health disorders. This study has the potential to have a transformative effect on the landscape of mental healthcare by providing new chances for early intervention,

tailored treatment, and better results. This transformation might be achieved via comprehensive examination and innovative approaches.

1.5 Motivation of Research

The research project with the working title "Roberta and BERT: Revolutionizing Mental Healthcare through Natural Language" was prompted by the urgent need to solve the substantial obstacles and inadequacies that are present in the existing system of providing mental healthcare. Depression, anxiety, and post-traumatic stress disorder (PTSD) are just some of the ailments that impact millions of people throughout the world. When it comes to mental health issues, they constitute a significant burden on the global community. Traditional methods to mental healthcare, on the other hand, often encounter obstacles such as availability, cost, stigma, and resource limits. As a consequence, these approaches frequently result in inequalities in care and treatment results that are less than optimum. When seen in this light, the development of sophisticated natural language processing (NLP) models like Roberta and BERT provides a compelling possibility to transform the delivery of mental healthcare. These state-of-the-art language models have shown exceptional skills in understanding and processing human language. As a result, they have provided insights into patients' mental states, emotions, and behaviors that are represented via language that have never been seen before. By utilizing the vast amounts of textual data that are generated through online platforms, electronic health records, and digital communication channels, Roberta and BERT make it possible for clinicians to acquire more profound insights into the mental health status of their patients, facilitate the early detection of symptoms, personalize treatment approaches, and provide targeted interventions. The purposed of this study is to investigate the transformational potential of Roberta and BERT in the field of mental healthcare by analyzing their applications in the areas of diagnosis, treatment, intervention, and support. The purpose of this project is to solve important gaps in the delivery of mental healthcare, increase access to care, better treatment results, and ultimately contribute to the well-being of those who are impacted by mental health illnesses. This will be accomplished by effectively utilizing the potential of modern natural language processing technology. In order to pave the way for a paradigm shift in the delivery of mental healthcare toward interventions that are more personalized, accessible, and effective, the research aims to elucidate the efficacy, feasibility, and ethical considerations associated with integrating Roberta and BERT into clinical practice. This will be accomplished through rigorous investigation and evaluation.

[2] Literature Review

Multiple studies have been conducted in the field of mental health, exploring the use of deep learning methods. This section presents a portion of the research.

Kumar, A., et al. (2023) proposed BERT and Bi-LSTM pipeline that was suggested demonstrated exceptional performance on two distinct Arabic datasets, with F1-scores of 1.00 and 0.82. On the other hand, the fine-tuned RoBERTa model obtained a similar F1-score of 0.60 in recognizing signs of depression from English social media postings. Most of the proposed deep learning models are end-to-end, requiring a more detailed explanation for their effectiveness. An AI model that can be explained has the potential to improve decision-making by providing transparency and interpretability. Hence, this study determines the areas in which the proposed system successfully acquired knowledge and the areas in which it did not



accurately recognize the indicators of depression. This information may be valuable for future advancements in the field of depression detection [1].

Ji, S., et al. (2021) focused develops and deploys two pre-trained masked language models, namely MentalBERT and MentalRoBERTa, with the aim of enhancing machine learning in the field of mental healthcare research. In addition, we assess the effectiveness of our specialized models and other versions of pretrained language models on many benchmarks for detecting mental disorders. Our findings show that language representations pretrained specifically for the target domain enhance the accuracy of mental health detection tasks [2].

Bokolo, B. et al. (2023) presented a dataset consisting of 632,000 tweets. It incorporates data preprocessing techniques, feature selection methods, and model training using logistic regression, Bernoulli Naive Bayes, random forests, DistilBERT, SqueezeBERT, DeBERTa, and RoBERTa models. The models' performance is assessed using evaluation criteria such as accuracy, precision, recall, and F1 score. The findings demonstrate that the RoBERTa model attains the greatest accuracy ratio of 0.981 and the highest mean accuracy of 0.97 (across 10 cross-validation folds) in identifying depression from tweets. This study showcases the efficacy of machine learning and sophisticated transformer-based models in using social media data for the purpose of mental health investigation. The results provide useful insights into the potential of internet platforms for early identification and monitoring of depression. This contributes to the expanding area of mental health analysis that relies on user-generated material [3].

Lin, E., et al. (2024) focused to detect suicidal intents in SuicideWatch Reddit postings by using the advanced RoBERTa-CNN model, which is a variation of the state-of-the-art RoBERTa model. RoBERTa is used for a range of Natural Language Processing (NLP) activities, such as text categorization and sentiment analysis. The efficacy of RoBERTa depends on its capacity to gather textual information and establish semantic links within texts. The RoBERTa model improves its capacity to capture significant patterns from large datasets by including a Convolutional Neural Network (CNN) layer. We assessed the performance of the RoBERTa-CNN model by conducting experiments on the Suicide and Depression Detection dataset, which yielded promising results. An instance of RoBERTa-CNN demonstrates a mean accuracy of 98%, with a standard deviation (STD) of 0.0009. Additionally, it achieves a mean AUC value of nearly 97.5% with a standard deviation of 0.0013 [4].

Jaun et al. (2022) examined brain biomarkers that might potentially aid in the diagnosis and prognosis of mental disorder outcomes. Machine learning has been used to classify mental disorders by using EEG signals. Nevertheless, the procedure of feature engineering (FE) and sub-sampling of raw EEG data is arduous and contingent upon the specific situation. Deep Learning (DL) is considered a crucial area of research for interpreting EEG data due to its capacity to be trained on large volumes of data generated by EEG and its use of automated feature extraction (FE) techniques that enhance the accuracy of discoveries using raw EEG signals. This study offers the findings of a thorough mapping analysis of 46 primary publications that were carefully selected for inclusion [5].

Ahmed et al. (2019) introduced a popular technique for diagnosing neurological disorders, specifically for identifying seizures, called the electroencephalogram (EEG). The EEG detects and records the electrical activity of the brain. This letter presents a novel approach for diagnosing epileptic seizures by classifying raw EEG data, without the need for predefined feature extraction. The method utilizes a one-dimensional convolutional variational autoencoder to integrate both supervised and unsupervised deep learning. The



recommended method is evaluated using k-fold cross-validation to determine its performance in classifying unknown data [6].

Acharya (2012) presented a computer-aided design (CAD) technique that utilizes nonlinear features to automatically differentiate between EEG data from those who are sober and those who are intoxicated. In order to train an SVM classifier with various kernel functions, the first step is to extract nonlinear features such as ApEn, LLE, SampEn, and other higher-order statistical (HOS) features. The input features consist of 1st, 2nd, and 3rd degree polynomials, together with an RBF (Radial Basis Function) kernel. Based on their research, these measurements are able to accurately distinguish between EEG signals that are connected to alcohol use and those that are not [7].

Tan et al. (2018) introduced the diagnosis of epilepsy, with the electroencephalogram (EEG) being often used as an additional diagnostic test. The EEG signal supplied specific information about the brain's electrical activity. Neurologists have typically used direct visual examination to identify irregularities in the form of epilepsy. This approach has several drawbacks, such as its potential laboriousness, susceptibility to technical artifacts, reliance on the reader's level of knowledge for drawing conclusions, and inability to consistently identify abnormalities. Therefore, it is essential to have a CAD system that can automatically identify this EEG data using machine learning. Until now, there has been no previous instance of analyzing EEG data using a CNN [8].

Aydin et al. (2019) conducted a study that specifically examined the effects of depression on a substantial proportion of the global population. The identification of this mood illness may be achieved by the use of EEG data. Analyzing EEG data manually to identify depression is arduous, time-consuming, and requires a high level of competence. Clinicians will get advantages from a fully automated depression diagnosis system that is constructed using EEG data. The deep model utilizes CNN layers to learn the temporal properties of signals, while lengthy LSTM layers facilitate the sequence learning process. This research used EEG data collected from both the left and right hemispheres of the brain [9].

Wang et al. (2017) conducted the extraction of valuable insights from complex, multi-dimensional, and heterogeneous biological data, which hinders the transformation of healthcare. Electronic health records, pictures, omissions, sensor data, and text are among the novel data formats that have lately appeared in modern scientific research. These data kinds are complex, diversified, inadequately documented, and sometimes unorganized. Prior to developing prediction or clustering models, traditional data mining and statistical learning approaches often need feature engineering to extract more precise and dependable properties from the information. Both stages are filled with challenges when working with intricate data and inadequate topic knowledge. Recent advancements in deep learning technology have offered new and successful approaches for developing end-to-end learning models from complex data [10].

Yang et al. (2021) discussed the integration of big data in the medical technology and biomedical industry, which has been driven by breakthroughs in artificial intelligence. In order to promote the progress of precision medicine, individuals must extract valuable knowledge from these extensive biological databases. Traditional machine learning algorithms frequently rely on feature engineering and domain knowledge from experts to extract features from biomedical data. This process requires a significant investment of time and human resources. Deep learning, a cutting-edge branch of machine learning, is capable of autonomously acquiring complex and resilient features from raw data, without the need for human involvement in feature building, unlike traditional approaches. Studies investigating the possible

applications of deep learning in many domains such as genetics, electronic health records, medicine, and medical imaging have shown that this technology has distinct advantages in improving the use of biological data and boosting the quality of healthcare [11].

Fathi et al. (2021) examined the use of deep learning techniques in healthcare systems, specifically assessing the advanced network topologies, applications, and industry trends. The primary goal is to provide a thorough comprehension of how deep learning models may be used in healthcare solutions to connect the divide between deep learning techniques and the interpretability of human healthcare. Lastly, let us address the current challenges and future prospects that are being encountered [12].

[3] Problem Statement

Conventional research that is centered on "Roberta and BERT: Revolutionizing Mental Healthcare through Natural Language" has a number of potential problems, one of the most significant of which is that it may rely excessively on technology as a stand-alone solution. In spite of the fact that Roberta and BERT unquestionably provide ground-breaking capabilities in natural language processing (NLP), the traditional method may fail to take into account the intricate nuances of mental healthcare. In the first place, there is a possibility of diminishing the significance of human empathy and comprehension in the context of therapeutic contacts. In spite of the fact that these models are very good at processing linguistic data, it is possible that they lack the emotional intelligence and contextual knowledge that are essential for professional counseling or therapy.

In the second place, there is the risk of ignoring the wide variety of human experiences and language expressions that are present in the setting of mental health. There is a possibility that conventional research might unintentionally reinforce biases that are already present in the data that is used to train these models, which could result in erroneous or incomplete evaluations of the mental states of people. Furthermore, it is impossible to disregard the privacy problems that are associated with the use of sensitive personal data in contexts of mental health applications. It is possible that conventional procedures will not effectively address these problems, which might result in trust violations and ethical conundrums. Additionally, there is a possibility of completely ignoring the socio-economic and cultural aspects that have an impact on mental health and the way language is used. It is possible that conventional research that is entirely focused on technology does not adequately account for these aspects, which may result in solutions that are neither culturally sensitive nor accessible to all people. Conventional research must be careful not to overlook the human aspects of therapy, reinforce biases, neglect privacy concerns, and fail to take into consideration socio-cultural factors. In conclusion, Roberta and BERT hold enormous potential for revolutionizing mental healthcare through natural language understanding. Therefore, conventional research must be cautious. For the purpose of achieving the full potential of these developments in mental healthcare, it is vital to use an integrated strategy that blends technical innovation with methods that are human-centered and morally acceptable.

[4] Proposed Work

The purpose of the proposed study is to investigate and assess the effectiveness of Roberta and BERT in bringing about a revolution in mental healthcare via the use of natural language processing (NLP). The study will be organized around a number of primary goals, including the following:



1. **Literature Review:** A review of the relevant literature: Conduct a comprehensive analysis of the available literature in order to get an understanding of the present status of research and applications of natural language processing (NLP) in the field of mental healthcare, with a particular emphasis on the uses of the Roberta and BERT models.
2. **Data Collection:** and Analysis: The gathering of information and its examination: Collect a wide variety of datasets that comprise textual information from a variety of sources, including patient interviews, social media platforms, online forums, and electronic health records. Conduct a thorough analysis of the datasets in order to identify patterns, trends, and linguistic indicators that are connected with various mental health problems.
3. **Model Development:** Developing and refining deep learning models based on Roberta and BERT architectures, with a focus on mental healthcare applications, is the third step in the model development process. For the purpose of optimizing the performance of the model, it is recommended to experiment with various setups, pre-training tactics, and fine-tuning approaches.
4. **Diagnostic Tools:** The Roberta and BERT models should be investigated for their potential utility as diagnostic tools for mental health issues. Using textual input, develop algorithms that are able to identify and categorize symptoms, evaluate the severity of the condition, and forecast the results of therapy.
5. **Personalized Treatment Recommendations:** It is important to investigate the possibility of using the Roberta and BERT models to provide individualized therapy suggestions for those who are suffering from mental health disorders. Create algorithms that can assess patient data and determine the results of previous treatments in order to provide recommendations for individualized interventions and therapeutic methods.
6. **Therapeutic Interventions:** To enhance therapeutic treatments and support services, it is recommended to investigate the use of the Roberta and BERT models. The development of virtual counseling platforms, chatbots, and self-help programs that make use of natural language processing technology in order to give real-time emotional support, coping methods, and psychoeducational materials is to be undertaken.
7. **Ethical Considerations:** Address the ethical concerns that are associated with the use of NLP models in the field of mental healthcare, such as the preservation of privacy, the provision of informed permission, and the presence of algorithmic bias. In order to ensure the implementation and application of the Roberta and BERT models in clinical practice in a reasonable manner, guidelines and best practices should be developed.
8. **Evaluation and Validation:** Conduct comprehensive validation studies and clinical trials in order to evaluate the performance, accuracy, and efficacy of the models and therapies that have been established. Determine how the influence will affect the results for patients, the satisfaction of clinicians, and the effectiveness of healthcare delivery.
9. **Dissemination and Knowledge Translation:** the outcomes of the study should be disseminated via the use of scholarly papers, presentations at conferences, and seminars. In order to assist the translation of information and the application of research results into practice, it is important to collaborate with experts in the field of mental healthcare, policymakers, and technology developers. Through the pursuit of these goals, the proposed study intends to expand our knowledge of how

natural language processing may transform mental healthcare by utilizing the Roberta and BERT models. The study aims to increase access to care, boost treatment results, and ultimately contribute to the well-being of those who are impacted by mental health illnesses. This will be accomplished via the creation of novel tools, treatments, and best practices.




Fig 1 Research Methodology

4.1 Bert

The Bidirectional Embedded Representation of Text (BERT) is a deep, unsupervised language representation that has been pre-trained on a plain text corpus. The Basic Evaluation and Review Toolkit (BERT) is a free and open source NLP machine learning platform (NLP). BERT is a technology to generate "contextualised" word embeddings/vectors, which is both its greatest strength and its greatest weakness because it is very compute-intensive at inference time, which means that if you want to use it in production at scale, it can become costly.




Fig 2 Bert Model

4.2 ROBERTA

To evaluate input sequences and construct contextualised representations of words in a phrase, RoBERTa, like BERT, is a transformer-based language model that employs self-attention. RoBERTa was trained on a significantly bigger dataset and using a more efficient training approach than BERT. RoBERTa is similar to BERT in terms of design, but it employs a byte-level BPE tokenizer (like GPT-2) and a unique pretraining strategy. Instead of the 16GB dataset used to train BERT, RoBERTa is trained on almost 160GB of uncompressed text. As an added bonus, RoBERTa is trained with I FULL-SENTENCES without NSP loss, ii) dynamic masking, iii) big mini-batches, and iv) a larger byte-level BPE.




Fig 3 Roberta Model

[5] Result and discussion

Research is considering common dataset for mental healthcare for all models. Here text attribute has been considered as input attribute, subject_matter has been considered for aspect while mental healthcare is output attribute. The main adjustments made by RoBERTa over BERT are bigger training data, using dynamic masking patterns, training on longer sequences and replacing the next sentence prediction.

Table 1 Configuration chart

Hyper parameter	Value
Epoch	50
Batch size	32
Optimizer	Adam
Learning rate	0.001

Considering above hyper parameter training operation has been performed using Bert. Accuracy details are discussed in following table

Table 2 Training, Validation, Testing accuracy using Bert

Epoch	Training Accuracy	Validation Accuracy	Testing accuracy
1	96.39465	96.44623	96.48759
2	96.48801	96.58254	96.59251
3	96.58064	96.58578	96.58638
4	96.59439	96.67096	96.733
5	96.64773	96.67849	96.75593
6	96.7081	96.73094	96.77264
7	96.73829	96.79533	96.7966
8	96.75267	96.7897	96.86658
9	96.7708	96.83165	96.89885
10	96.79525	96.83341	96.90691
11	96.80932	96.83383	96.91708
12	96.89061	96.90992	96.97362
13	96.97276	97.05466	97.07504
14	97.06504	97.10621	97.17417
15	97.11713	97.1483	97.19143
16	97.17402	97.27154	97.27359
17	97.25181	97.34159	97.43515
18	97.29083	97.35823	97.45309
19	97.29848	97.30941	97.37696
20	97.39794	97.45233	97.45863
21	97.41341	97.50688	97.507
22	97.48132	97.50311	97.59964
23	97.50617	97.52122	97.60887
24	97.52913	97.57771	97.58161
25	97.56859	97.62456	97.66246
26	97.62219	97.63267	97.66681
27	97.71408	97.75832	97.76175
28	97.73452	97.75769	97.81395
29	97.79187	97.88694	97.90402
30	97.79837	97.83714	97.93644
31	97.86772	97.87972	97.97086
32	97.94325	97.96362	98.02637
33	98.01921	98.05282	98.13085
34	98.09605	98.1689	98.25018
35	98.14053	98.20522	98.2108
36	98.16126	98.21005	98.26836
37	98.23673	98.26993	98.32635
38	98.26006	98.29979	98.30175

39	98.31997	98.37406	98.40411
40	98.33425	98.40005	98.43416
41	98.34833	98.41374	98.41563
42	98.39506	98.44436	98.4677
43	98.48424	98.50133	98.58124
44	98.5309	98.54078	98.56809
45	98.61831	98.66393	98.74478
46	98.65559	98.69317	98.7111
47	98.66859	98.73193	98.73788
48	98.73349	98.77052	98.83317
49	98.78696	98.82894	98.89444
50	98.85136	98.92872	98.97938




Fig 4 Bert based simulation

Considering above hyper parameter training operation has been performed using Roberta. Accuracy details are discussed in following table

Table 3 Training, Validation, Testing accuracy using Roberta

Epoch	Training Accuracy	Validation Accuracy	Testing accuracy
1	96.37567586	96.40724097	96.47016118
2	96.40225933	96.46087651	96.48640722
3	96.44320541	96.53536685	96.6279188
4	96.45788074	96.54949543	96.58977664
5	96.51579513	96.54485821	96.61472199
6	96.59760916	96.64178404	96.66506705
7	96.62961566	96.66220709	96.74595923
8	96.6346795	96.63715866	96.72199183
9	96.65241246	96.66440124	96.69811812

10	96.68302739	96.75754728	96.79094551
11	96.70059006	96.79906645	96.87618253
12	96.77161185	96.8285047	96.8948667
13	96.83752506	96.93076664	96.99316283
14	96.92518718	96.94002812	96.99634974
15	97.01107396	97.1106766	97.19567432
16	97.03975871	97.10364643	97.12607215
17	97.05458346	97.14536266	97.21113456
18	97.07037338	97.08127501	97.11992073
19	97.12557429	97.15753897	97.22212141
20	97.21830559	97.21923445	97.23114468
21	97.31157058	97.32570853	97.33115719
22	97.31524256	97.33264307	97.4213098
23	97.37235042	97.40877637	97.46973071
24	97.46657032	97.49571156	97.58944799
25	97.4855833	97.51871353	97.58529806
26	97.54916067	97.60531528	97.69101872
27	97.56920121	97.5981609	97.67228431
28	97.65806892	97.69276534	97.73700495
29	97.72838585	97.7941185	97.88528297
30	97.8237456	97.87538746	97.92424983
31	97.89693263	97.92246979	98.01784686
32	97.91633145	97.92925233	97.96464952
33	97.99910064	98.03799948	98.1217557
34	98.06135281	98.1056199	98.18297149
35	98.081624	98.10560936	98.10709223
36	98.16847161	98.2113741	98.25793268
37	98.26375442	98.35850201	98.38625261
38	98.36202436	98.38202279	98.4322685
39	98.42220133	98.42239773	98.4733767
40	98.51668391	98.59223187	98.61336576
41	98.59603151	98.6294684	98.67631478
42	98.63168194	98.71199696	98.77284633
43	98.63316658	98.73245618	98.75754254
44	98.6972158	98.69924672	98.72708543
45	98.78572274	98.81298831	98.82761964
46	98.81839407	98.88488775	98.93174382
47	98.88814525	98.89904402	98.91878238
48	98.95256367	99.01891571	99.02521033

49	98.99333171	99.01619527	99.03827671
50	99.05975187	99.11343021	99.16350457




Fig 5 Bert based simulation

5.1 Comparative analysis of LSTM, Bert and Roberta

Table 4 is presenting existing accuracy of different approaches.

Table 4 Accuracy comparison of different Approaches

	LSTM	Bert	Roberta
Accuracy	99.65%	99.73%	99.88%

Following figure is presenting the accuracy chart of existing technical approaches by considering the accuracy values as shown in table 4.




Fig 6 Accuracy comparison of Approaches

Table 5 is presenting existing accuracy of different approaches.

Table 5 Error comparison of different Approaches

	LSTM	Bert	Roberta
Error rate	0.35%	0.27%	0.12%




Fig 7 Comparison of Error rate

[6] Conclusion

In conclusion, the study that was conducted on the topic of "Roberta and BERT: Revolutionizing Mental Healthcare through Natural Language" highlights the transformational potential of sophisticated natural language processing (NLP) models in terms of changing the delivery of mental healthcare. Mental health experts are able to get insights into patients' feelings, thoughts, and mental states that are represented via language that have never been seen before because to the deployment of cutting-edge technology like as Roberta and BERT. It is because of this that more accurate diagnosis, tailored treatment planning, and focused interventions are made possible, which eventually leads to an improvement in the efficiency and accessibility of mental healthcare professionals. This study ushers in a new era in mental healthcare, one in which novel uses of natural language processing technologies pave the way for a more compassionate, inclusive, and successful approach to treating persons who are struggling with mental health illnesses. This new era is heralded by the utilization of the abilities of natural language understanding. The continuation of investments in research and development in this field offers the prospect of additional developments, which will eventually lead to improvements in outcomes and quality of life for persons who are impacted by mental health difficulties.

[7] Future Scope

Research towards "Roberta and BERT: Revolutionizing Mental Healthcare through Natural Language" has great promise for the future of mental healthcare and might lead to groundbreaking innovations. In order to make the most of these technologies, future research should focus on a few important areas. To begin, current mental healthcare systems should be further investigated for potential ways to include Roberta and BERT. Making ensuring mental health practitioners can make good use of these technologies in their practice should be the focus of future research into creating user-friendly interfaces and smooth integration

procedures. Second, researchers should focus on building more complex language models that can understand the subtleties of mental health conversations. To reduce bias and increase the reliability of mental health evaluations, it is necessary to train models on varied datasets that include a broad variety of language expressions and cultural settings. If we want mental health providers and patients to have greater faith in and use these language models, we need to make sure they are easier to understand and explain. As part of this process, we must devise ways to illustrate and clarify model predictions and do thorough tests to see how well these predictions hold up in actual clinical situations. There should also be studies looking into the moral questions raised by using BERT and Roberta in mental healthcare settings. Ensuring that these technologies are used in a way that respects patient autonomy and encourages equal access to treatment is part of this, as is addressing concerns about privacy, data security, and algorithmic bias. Last but not least, studies should be conducted in the future to determine how well and for how long mental healthcare practitioners may benefit by using Roberta and BERT. In order to determine whether or whether these technologies are cost-effective in enhancing the availability and quality of mental healthcare services, it is necessary to do longitudinal studies that measure patient outcomes, clinician satisfaction, and the total impact of these technologies.

References

- Kumar, A., Kumari, J., & Pradhan, J. (2023). Explainable Deep Learning for Mental Health Detection from English and Arabic Social Media Posts. In *ACM Transactions on Asian and Low-Resource Language Information Processing*. Association for Computing Machinery (ACM).
- Ji, S., Zhang, T., Ansari, L., Fu, J., Tiwari, P., & Cambria, E. (2021). MentalBERT: Publicly Available Pretrained Language Models for Mental Healthcare. *arXiv*.
- Bokolo, B. G., & Liu, Q. (2023). Deep Learning-Based Depression Detection from Social Media: Comparative Evaluation of ML and Transformer Techniques. In *Electronics* (Vol. 12, Issue 21, p. 4396). MDPI AG. <https://doi.org/10.3390/electronics12214396>
- Lin, E., Sun, J., Chen, H., & Mahoor, M. H. (2024). Data Quality Matters: Suicide Intention Detection on Social Media Posts Using a RoBERTa-CNN Model (Version 1). *arXiv*. <https://doi.org/10.48550/ARXIV.2402.02262>
- Rivera, M. J., Teruel, M. A., Maté, A., Trujillo, J.: Diagnosis and prognosis of mental disorders by means of EEG and deep learning: a systematic mapping study. *Artificial Intelligence Review* 55(2), 1209–1251 (2021).
- Abdelhameed, A. M., Bayoumi, M.: Semi-Supervised EEG Signals Classification System for Epileptic Seizure Detection. *IEEE Signal Processing Letters* 26(12), 1922–1926 (2019).
- Acharya, U. R., Sree, S. V., Chattopadhyay, S., Suri, J. S.: Automated Diagnosis Of Normal And Alcoholic EEG Signals. *International Journal of Neural Systems* 22(03), 1250011 (2012)
- Acharya, U. R., Oh, S. L., Hagiwara, Y., Tan, J. H., Adeli, H.: Deep Convolutional Neural Network for the Automated Detection and Diagnosis of Seizure Using EEG Signals. *Computers in Biology and Medicine* 100, 270–278 (2018). 0 20 40 60 80 100 120 10 20 30 40 50 60 70 80 90 100 Time Taken in Sec Epoch Conventional Approach Proposed Work 13



- Ay, B., Yildirim, O., Talo, M., Baloglu, U. B., Aydin, G., Puthankattil, S. D., Acharya, U. R.: Automated Depression Detection Using Deep Representation and Sequence Learning with EEG Signals. *Journal of Medical Systems* 43(7), (2019).
- Miotto, R., Wang, F., Wang, S., Jiang, X., & Dudley, J. T.: Deep learning for healthcare: review, opportunities and challenges. *Briefings in Bioinformatics* 19(6), 1236–1246 (2017). Oxford University Press (OUP).
- Yang, S., Zhu, F., Ling, X., Liu, Q., Zhao, P.: Intelligent Health Care: Applications of Deep Learning in Computational Medicine. *Frontiers in Genetics* 12 (2021).
- Shamshirband, S., Fathi, M., Dehzangi, A., Chronopoulos, A. T., Alinejad-Rokny, H.: A review on deep learning approaches in healthcare systems: Taxonomies, challenges, and open issues. *Journal of Biomedical Informatics* 113, 103627 (2021).

