

Prevalence of Chikungunya, Dengue and Zika viruses in blood donors: a systematic literature review and meta-analysis

Ángel Giménez-Richarte¹, Mabel Ortiz de Salazar¹, Cristina Arbona¹, María P. Giménez-Richarte², Miriam Collado¹, Pedro L. Fernández¹, Francisco Quiles¹, Carlos Clavijo¹, Pascual Marco^{3,4}, Jose-Manuel Ramos-Rincon⁴



¹Valencian Community Blood Transfusion Centre, Valencian Community, Spain;

²Medical student, Miguel Hernandez University of Elche, Alicante, Spain;

³Service of Haematology, General University Hospital of Alicante-ISABIAL, Alicante, Spain;

⁴Clinical Medicine Department, Miguel Hernandez University of Elche, Alicante, Spain

Background - Blood transfusion centres should understand the epidemiology of emerging diseases that are transmissible through the transfusion of blood components. The risk of transmission of arboviruses through this route has become apparent in recent years. The aim of our study is to summarise the reported prevalence (viraemic rate, seroprevalence and/or antigen detection) of Chikungunya (CHIKV), Dengue (DENV) and Zika (ZIKV) viruses in blood donors according to screening test used and world region.

Materials and methods - We conducted a systematic literature review and meta-analysis having searched for information in the main bibliographic databases (MEDLINE, Embase, and Scopus). The prevalence for each of the viruses was calculated according to the screening test used and geographic location.

Results - We included 18 records on CHIKV, 71 on DENV, and 27 on ZIKV. The highest prevalences of RNA for CHIKV were 1.9% in Puerto Rico (2014), 1.0% in Thailand (2009), and 1.0% in French Polynesia (2014-15). The highest prevalences of RNA for DENV were 5.5% in Saudi Arabia (2015-16), 2.3% in Madeira, Portugal (2012-13), and 0.6% in Brazil (2012). The highest prevalences of RNA for ZIKV were 2.8% in French Polynesia (2013-14), 2.7% in Brazil (2015-16), and 1.8% in Martinique (2016). Overall seroprevalence, as assessed by IgG antibodies, was 21.6% for CHIKV, 24.0% for DENV, and 5.1% for ZIKV.

Discussion - Our study shows a high proportion of donors who are viraemic and asymptomatic, especially during outbreaks, with prevalences surpassing 5% for DENV, 1% for CHIKV, and 2% for ZIKV. These data confirm a clear threat to blood transfusion safety. The elevated seroprevalence for these three arboviruses is also indicative of their wide circulation in populations, correlating with an increased risk of infected but asymptomatic donors. Health centres and institutions must address this threat, especially in tropical regions where the biggest outbreaks occur.

Keywords: *Chikungunya virus, Dengue virus, Zika virus, blood transfusion, blood safety.*

INTRODUCTION

Emerging and re-emerging viruses have materialised as the latest challenge to blood transfusion safety. In this sense, the World Health Organisation (WHO) has called for blood

Arrived: 31 March 2021

Revision accepted: 20 July 2021

Correspondence: Ángel Giménez Richarte
e-mail: gimenez_ang@gva.es

transfusion centres to be informed of the epidemiology of different emerging transfusion-transmitted infections and to evaluate the possible impact on donor selection criteria and the supply of blood products¹.

Among these emerging viruses, arboviruses are especially relevant because of their known or theoretical potential for transmission through blood transfusions². Within this group, Chikungunya virus (CHIKV), Dengue virus (DENV), and Zika virus (ZIKV) stand out for their high global incidence and the wide dissemination of their vector.

CHIKV is an alphavirus in the *Togaviridae* family, transmitted by *Aedes* mosquitoes (e.g. *A. albopictus*, *A. aegypti*). Following an incubation period of 1 to 12 days, the acute phase of infection by this virus is characterised by fever; severe, incapacitating arthralgia; and other non-specific symptoms. Some patients also develop chronic illness³. Since the virus was first isolated, periodic outbreaks have been reported in Africa, Asia, and islands in the Indian Ocean, while the first cases in the Americas were reported in 2013. Since then, different outbreaks have been reported across regions of South and Central America⁴. In Europe, several outbreaks have occurred since 2007⁵, including one in 2015 involving 693,489 suspected and 37,480 confirmed cases⁶. Although no cases of transfusion-related infections have been notified, organisations such as the American Association of Blood Banks have sounded the alarm on the theoretical potential given the high percentage of asymptomatic people infected (3% to 28%) and the high rates of viraemia that they have⁷. One case of iatrogenic CHIKV transmission was reported following an accidental needle puncture in France⁸.

For its part, DENV is a flavivirus in the *Flaviviridae* family. Four distinct serotypes have been documented: DEN-1, DEN-2, DEN-3, and DEN-4. Like CHIKV, DENV is transmitted by *Aedes* mosquitoes, usually *A. aegypti*. It is the main arbovirus worldwide in terms of mortality and morbidity; its incubation period is normally 4 to 7 days, although it can range from 3 to 10 days. The clinical classification of dengue divides cases into those with or without warning signs and severe dengue (including dengue shock syndrome)⁹. The first large epidemics date back to the 1870s¹⁰. Today, the disease is endemic in more than 100 countries from the WHO regions of Africa, the Americas, the Eastern Mediterranean, Southeast

Asia, and the Western Pacific; in 2015 alone, more than 3.2 million cases were notified across the Americas, Southeast Asia, and the Western Pacific¹¹. In Europe, 11 cases of local transmission were also reported in 2019¹². Since 2002, numerous cases of transfusion-transmitted infections have been described in Hong Kong, Singapore, Brazil, Pakistan, and Puerto Rico (USA)¹³⁻¹⁷.

ZIKV is another flavivirus from the *Flaviviridae* family. *Aedes* spp. mosquitoes such as *A. africanus*, *A. aegypti*, and *A. albopictus* are the vectors of transmission, and the incubation period can be anywhere from 2 to 12 days. Although 80% of infected people remain asymptomatic, an acute presentation with non-specific symptoms, such as fever, arthralgia, and exanthema, can occur. Infection has also been related to the appearance of microcephalia in neonates (congenital Zika syndrome) and to a Guillain-Barré-type neurological presentation¹⁸. For decades, little attention was paid to this virus, as it only provoked isolated cases in Southeast Asia and Africa. However, in 2007, a large epidemic outbreak was registered on Yap Island (Micronesia), and in 2015 and 2016, another large outbreak occurred in the Americas. In 2019, the first two cases of local transmission were reported in Europe (France)¹⁹. Moreover, transmission via transfusion of platelets has been reported in Brazil^{20,21}.

Upon performing a review of the available scientific literature on the prevalence of CHIKV, DENV, and ZIKV in blood donors, we identified only two systematic reviews: one by Liu *et al.*, with very restrictive inclusion criteria and ten included studies on ZIKV, and one by Eick *et al.*, with three included studies on the prevalence of ZIKV and 11 on the prevalence of DENV^{22,23}. We did not identify any similar papers on CHIKV. There is, therefore, a lack of literature giving a broad overview of the prevalence of these three arboviruses in blood donors.

The emergence of these viruses represents a real threat to obtaining blood components and has a direct impact on donor selection criteria and the stock of components. Following WHO recommendations, the aim of this study was to summarise the reported and published prevalence of CHIKV, DENV and ZIKV in blood donors according to the screening test used (viraemic rate, seroprevalence or antigen detection) and world region (geographical region and country).

We define the research question in a PICOS (population,

intervention, comparison, outcome, study) format. The population was blood donors, including conventional whole blood donors and those donating via apheresis, who were screened for the target viruses using any test and in any defined geographic region. The intervention was screening using different techniques to detect antibodies, antigens, or nucleic acids. Comparisons were not applicable to this question and the outcomes were reported and published prevalences of each virus according to the screening test used and the geographic region in which the screening was performed. Any primary studies were included.

MATERIALS AND METHODS

Design

A systematic literature review and meta-analysis were designed and conducted in accordance with the *Cochrane Handbook of Systematic Reviews of Interventions* and reported in line with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement (see *Online Supplementary Content, Table SI*). Although no published protocol is available, we collected and analysed data according to pre-specified outcomes and methods, performing meta-analyses, pooling the data obtained in the different studies, and establishing a single prevalence estimate for each virus. This information may be of interest from an epidemiological point of view and when considering measures with a possible impact on transfusion safety.

Sources of data

We conducted literature searches in MEDLINE, Embase, and Scopus using the University Miguel Hernández server.

Search strategy

We performed free text searches in the three bibliographic databases using the terms: “transfusion” AND “Dengue”, “transfusion” AND “Chikungunya”, “transfusion” AND “Zika”, “blood donation” AND “Dengue”, “blood donation” AND “Chikungunya”, “blood donation” AND “Zika”.

In MEDLINE, we also used Medical Subject Headings (MeSH): “blood transfusion” AND “Zika virus”, “blood donors” AND “Zika virus”, “blood transfusion” AND “Dengue”, “blood donors” AND “Dengue”, “blood transfusion” AND “Chikungunya virus”, “blood donors” AND “Chikungunya virus”.

Finally, in the Embase searches we used the Emtree thesaurus with the terms: “blood transfusion” AND “Zika virus”, “blood donor” AND “Zika virus”, “blood transfusion” AND “Chikungunya”, “blood donor” AND “Chikungunya”, “blood transfusion” AND “Dengue”, “blood donor” AND “Dengue”.

The records were entered into the Mendeley Desktop reference manager (Elsevier). The search was performed from the year of database inception to March 28, 2020. A weekly alert system was set up to update the search with any relevant results until August 7, 2020.

Study selection

First, we used the bibliographic reference manager to create folders containing records for each virus, eliminating duplicates. We then conducted an initial screening of titles and abstracts and retrieved the full text of all pre-selected records.

Eligible studies were publications in any language describing the prevalence of the virus in blood donor screening (both donors of conventional whole blood and those donating via different methods of apheresis). We included all studies (original articles, brief reports, letters to the editor, and conference papers) reporting the number of positive results as a proportion of total samples analysed, as long as the paper stated the type of test used for screening (serological tests, antigen tests or nucleic acid amplification tests [NAT]) and the geographic location of the study population. We excluded studies that involved people other than blood donors, such as patients, children, pregnant women, the general non-donor population, and other non-donor or unspecified populations.

A single review author selected all included articles, obviating the need for an analysis of interobserver concordance. We did perform an intraobserver concordance analysis, including in the review all records that were deemed to meet inclusion and exclusion criteria during two critical assessments of the full texts.

When a single record reported results for two different study populations, these were separated in the analysis if the participants' characteristics differed for important variables, for example geographic region (e.g. studies evaluating one population in Africa and another in Europe), or if the prevalence was substantially different by population (i.e. we separated populations sub-nationally if the differences in prevalence were relevant). If a single

population underwent screening using more than one type of test, separate analyses were performed for each. In the case of records with overlapping study populations, we selected the most relevant publication (i.e. with the largest number of screened donors).

Data extraction and analysis

A single review author extracted data on prevalence and populations from the studies included, directly entering the data into Comprehensive Meta-Analysis (CMA) software. A second review author double-checked that all data were entered correctly. We recalculated the prevalence for the three viruses, pooling all positive cases and donors screened for each to calculate an overall proportion of positive results in the blood donor screening for CHIKV, DENV, and ZIKV, according to the type of screening test used. We then stratified the results by geographic location or country as long as there were at least three included studies, the minimum number we considered capable of representing a geographic area. Moreover, when the number of publications and the nature of the screening test allowed it, we calculated the prevalence ratio according to whether or not the study had

taken place in an endemic region or during an epidemic outbreak. If so, we calculated the prevalence.

Results are expressed as prevalences with 95% confidence intervals and are presented with forest plots. We evaluated the heterogeneity of the studies for each screening test using the I^2 statistic. To calculate the confidence intervals, create the forest plots, and analyse the heterogeneity, we used CMA software, version 2 (Borenstein, Hedges, Higgins, & Rothstein, 2005).

Quality assessment

To evaluate the methodological quality of included records, we used the STROBE (STrengthening the Reporting of Observational studies in Epidemiology) checklists for cross-sectional studies and conference abstracts. No tools were applied to letters to the editor.

RESULTS

Figure 1 presents the PRISMA flow chart, describing the study selection process. Following full-text assessment, 18 studies on CHIKV²⁴⁻⁴¹, 71 on DENV^{24,26,27,31,33,36,38,40,42-104}, and 27 on ZIKV^{26,28,32,36,38,40,67,105-124} were included. *Online Supplementary Content, Tables SII-SIV*, describes the main characteristics of the studies included.




Figure 1 - PRISMA flow chart on selection of studies included for each virus covered by the literature review

Analysis of results

For each virus analysed, the prevalence varied according to the screening technique used, the geographic regions, and their characteristics.

Prevalence of Chikungunya virus

According to the assessment of IgG antibodies, the overall seroprevalence of CHIKV was 21.6% (95% CI: 20.6% to 22.5%). Several studies reported a seroprevalence of 0%, while one in Rwanda in 2015 found a seroprevalence of 63.0% (95% CI: 59.8% to 66.2%)²⁸. By regions, the highest value was in Africa (seroprevalence 37.8%, 95% CI: 36.2% to 39.4%). The prevalence ratio between studies performed

in an endemic area or during an outbreak and those in non-endemic regions was 24.4 (Table I).

The only study we identified on the seroprevalence of IgM antibodies against CHIKV in blood donors reported a seroprevalence of 5.5% (95% CI: 3.1% to 9.7%) (Table I).

Finally, NAT showed an overall prevalence of 0.5% (95% CI: 0.4% to 0.5%). The highest rates were in the screening in Puerto Rico in 2014 (prevalence 1.9%, 95% CI: 1.4% to 2.4%)³⁵. In populations living in endemic areas or going through epidemic outbreaks, the prevalence was 0.6% (95% CI: 0.6% to 0.7%), compared to 0% in non-endemic regions (Table I).

Table I - Global prevalence of Chikungunya virus in blood donors, by population

Study ID	Population	Screening test	Positive/ Total	Prevalence	95% CI	Forest plot	Relative weight %
IgG antibodies							
Sayama 2013	Laos 2012	ELISA (Not specified)	1/199	0.005	<0.001 0.035		0.1
Humphrey 2019	Egypt 2013-16	ELISA (Euroimmun)	11/199	0.055	0.031 0.097		1.2
Humphrey 2019	India 2013-16	ELISA (Euroimmun)	22/200	0.110	0.074 0.161		2.3
Humphrey 2019	Iran 2013-16	ELISA (Euroimmun)	0/113	0.000	<0.001 0.066		0.1
Humphrey 2019	Jordan 2013-16	ELISA (Euroimmun)	1/199	0.005	<0.001 0.035		0.1
Humphrey 2019	Lebanon 2013-16	ELISA (Euroimmun)	1/116	0.009	0.001 0.059		0.1
Humphrey 2019	Pakistan 2013-16	ELISA (Euroimmun)	3/200	0.015	0.005 0.046		0.4
Humphrey 2019	Palestine 2013-16	ELISA (Euroimmun)	6/200	0.030	0.014 0.065		0.7
Humphrey 2019	Philippines 2013-16	ELISA (Euroimmun)	21/199	0.177	0.118 0.256		2.0
Humphrey 2019	Qatar 2013-16	ELISA (Euroimmun)	7/200	0.035	0.017 0.072		0.8
Humphrey 2019	Sudan 2013-16	ELISA (Euroimmun)	5/97	0.052	0.020 0.120		0.6
Humphrey 2019	Syria 2013-16	ELISA (Euroimmun)	1/200	0.005	<0.001 0.035		0.1
Humphrey 2019	Yemen 2013-16	ELISA (Euroimmun)	4/149	0.027	0.010 0.069		0.5
Seruyange 2019	Rwanda 2015	ELISA (In house)	551/874	0.630	0.598 0.662		24.1
Seruyange 2019	Sweden 2015	ELISA (In house)	172/15	0.079	0.050 0.124		1.9
Moyen 2014	Congo 2011	ELISA (Not specified)	178/517	0.344	0.305 0.386		13.8
Clements 2019	Uganda 2006-07	ELISA (In house)	552/1744	0.317	0.295 0.339		44.6
Slavov 2018	Brazil 2015	ELISA (Abcam)	1/442	0.002	<0.001 0.016		0.1
Slavov 2018	Brazil 2016	ELISA (Abcam)	0/445	0.000	<0.001 0.017		0.1
Saba Villarreal 2018	Bolivia B. and S.C.* 2016-17	ELISA (Euroimmun)	87/168	0.518	0.443 0.592		5.0
Saba Villarreal 2018	Bolivia, others 2016-17	ELISA (Euroimmun)	15/281	0.053	0.032 0.087		1.7
Total			1484/6887	0.216	0.206 0.225		
Subgroups							
Southeast Asia			44/518	0.085	0.064 0.112		
West Asia			23/1377	0.017	0.011 0.025		
Africa			1297/3431	0.378	0.362 0.394		
America			103/1346	0.077	0.064 0.092		
Endemic or epidemic area			1426/3454	0.413	0.397 0.429		
Non-endemic or epidemic area			58/3433	0.017	0.013 0.022		
<i>Heterogeneity I² 97.70</i>							
IgM antibodies							
Sayama 2013	Laos 2012	ELISA (Not specified)	11/199	0.055	0.031 0.097		
NAT							
Sayama 2013	Laos 2012	Not specified	0/199	0.000	0.000 0.039		0.2
Appassakij 2014	Thailand 2009	In house	20/2000	0.010	0.007 0.015		5.8
Beau 2020	French Polynesia 2014-15	Altona D.	34/3433	0.010	0.007 0.014		9.9
Brouard 2008	Reunion Island 2005-07	Not specified	2/500	0.004	0.001 0.016		0.6
Benites 2019	Brazil 2015-16	Bio-Manguinhos	0/3737	0.000	0.000 0.002		0.2
Stramer 2019	USA 2011-12	Cobas test	0/10528	0.000	0.000 <0.001		0.2
Chiu 2015	Puerto Rico 2014	In house	3/557	0.005	0.002 0.017		0.9
Simmons 2016	Puerto Rico 2014	In house	161/26688	0.006	0.005 0.007		46.8
Simmons 2016	Puerto Rico 2014	In house	56/3007	0.017	0.014 0.024		16.1
Saã 2019	Puerto Rico 2015-16	Not specified	0/1186	0.000	0.000 0.007		0.2
Slavov 2018	Brazil 2015-16	Not specified	0/897	0.000	0.000 0.009		0.2
Sharma 2018	Brazil 2016	In house	0/676	0.000	0.000 0.012		0.2
Gallian 2017	Guadeloupe 2014-15	Altona D.	22/6189	0.004	0.002 0.005		6.4
Gallian 2017	Martinique 2014-15	Altona D.	43/10197	0.004	0.003 0.006		12.5
Sargento 2017	Portugal 2017	In house	0/110	0.000	0.000 0.068		0.2
Total			341/69904	0.005	0.004 0.005		
Americas			285/63662	0.005	0.004 0.005		
Puerto Rico			220/31438	0.007	0.006 0.008		
Brazil			0/5310	0.000	0.000 0.000		
Endemic or epidemic areas			341/54433	0.006	0.006 0.007		
Non-endemic or non-epidemic areas			0/15471	0.000	0.000 <0.001		
<i>Heterogeneity I² 88.03</i>							

*Beni and Santa Cruz regions

CI: confidence interval; NAT: nucleic acid test; ELISA: enzyme-linked immunosorbent assay

Prevalence of Dengue virus

Tests for IgG antibodies showed an overall seroprevalence of DENV of 24.0% (95% CI: 23.5% to 24.4%). Several studies reported a seroprevalence of more than 90% of screened individuals, for example in the Philippines, Puerto Rico, Brazil, Guadeloupe and Martinique, and the Dominican Republic^{27,81,90,98,100}. By geographic region, the Americas

stand out for the high seroprevalence of 61.3% (95% CI: 60.0% to 62.6%), followed by Africa at 22.0% (95% CI: 21.0% to 23.1%) and Southeast Asia at 20.4% (95% CI: 19.7% to 21.1%). Saudi Arabia and Brazil were the individual countries with the highest seroprevalence (36.0% and 32.5%, respectively). The prevalence ratio between endemic and non-endemic regions was 13.9 (Table II).

Table II - Global prevalence of IgG antibodies against Dengue virus in blood donors, by population

Study ID	Population	Screening test	Positive/ Total	Prevalence	95% CI	Forest plot			Relative weight %
						0.00	0.50	1.00	
Zeng 2018	China 2015	ELISA (Panbio)	34/819	0.042	0.030 0.058				0.9
Liao 2017	China 2014	ELISA (Panbio)	51/1500	0.034	0.026 0.045				1.3
Kwan 2017	China 2014	ELISA (Panbio)	86/3827	0.023	0.018 0.028				2.3
Gao 2017	China 2013-14	ELISA (Panbio)	7/1685	0.004	0.002 0.009				0.2
Ranjan 2016	India 2012	ELISA (Novatec)	116/200	0.580	0.511 0.647				1.3
Chhabra 2013	India	IC (J. Mitra & Co)	3/380	0.008	0.003 0.024				0.1
Jain 2019	India 2016	IC (R.L.)	55/369	0.149	0.116 0.189				1.3
Hossain 2003	Bangladesh 1966-1997	ELISA (Not specified)	1/184	0.005	0.001 0.038				<0.0
Sayama 2013	Laos 2012	ELISA (Not specified)	139/199	0.699	0.631 0.758				1.1
Harif 2014	Malaysia 2009-10	ELISA (Panbio)	151/360	0.419	0.370 0.471				2.4
Mohamad 2016	Malaysia 2015	ELISA (Not specified)	1/126	0.008	0.001 0.054				<0.0
Low 2015	Singapore 2009-10	ELISA (Panbio)	1885/3627	0.520	0.503 0.536				24.7
Humphrey 2019	Egypt 2013-16	ELISA (Novatec)	40/199	0.201	0.151 0.262				0.9
Humphrey 2019	India 2013-16	ELISA (Novatec)	125/200	0.625	0.556 0.689				1.3
Humphrey 2019	Iran 2013-16	ELISA (Novatec)	6/113	0.053	0.024 0.113				0.2
Humphrey 2019	Jordan 2013-16	ELISA (Novatec)	9/199	0.045	0.024 0.085				0.2
Humphrey 2019	Lebanon 2013-16	ELISA (Novatec)	6/116	0.052	0.023 0.110				0.2
Humphrey 2019	Pakistan 2013-16	ELISA (Novatec)	40/200	0.200	0.150 0.261				0.9
Humphrey 2019	Palestine 2013-16	ELISA (Novatec)	17/200	0.085	0.054 0.133				0.4
Humphrey 2019	Philippines 2013-16	ELISA (Novatec)	114/119	0.958	0.903 0.982				0.1
Humphrey 2019	Qatar 2013-16	ELISA (Novatec)	7/200	0.035	0.017 0.072				0.2
Humphrey 2019	Sudan 2013-16	ELISA (Novatec)	47/97	0.485	0.387 0.583				0.7
Humphrey 2019	Syria 2013-16	ELISA (Novatec)	26/200	0.130	0.090 0.184				0.6
Humphrey 2019	Yemen 2013-16	ELISA (Novatec)	36/149	0.242	0.180 0.317				0.8
Faddy 2013	Australia 2008-09	ELISA (Panbio)	323/3553	0.091	0.082 0.101				8.0
Faddy 2012	Australia 2008-09	ELISA (Panbio)	182/1799	0.101	0.088 0.116				4.5
Faddy 2015	Australia 2008-09	ELISA (Panbio)	31/457	0.068	0.048 0.095				0.8
Darcy 2001	Solomon Islands 1994-1995	ELISA (Panbio)	202/515	0.392	0.351 0.435				3.4
Aubry 2015	French Polynesia 2011-13	ELISA (Not specified)	476/593	0.803	0.769 0.833				2.6
Ergunay 2010	Turkey	ELISA (Euroimmun)	21/2435	0.009	0.006 0.013				0.6
Tezcan 2014	Turkey 2010-11	IC (S.D.)	153/920	0.166	0.144 0.192				3.5
Ashshi 2015	Saudi Arabia 2014	ELISA (Panbio)	7/100	0.070	0.034 0.140				0.2
Ashshi 2017	Saudi Arabia 2015-16	ELISA (Panbio)	355/910	0.390	0.359 0.422				5.9
Jamjoom 2016	Saudi Arabia	ELISA (Panbio)	68/184	0.370	0.303 0.442				1.2
Aghaie 2014	Iran 2012	ELISA (Panbio)	41/540	0.076	0.056 0.102				1.0
Larrieu 2014	Reunion Island 2008	ELISA (Not specified)	72/1825	0.040	0.031 0.049				1.9
Vairo 2014	Tanzania 2011	ELISA (Panbio)	253/500	0.506	0.462 0.550				3.4
Collembert 2006	Burkina Faso 2003-04	ELISA (Panbio)	62/191	0.325	0.262 0.394				1.1
Sawadogo 2019	Burkina Faso 2016	ELISA (Panbio)	721/1007	0.716	0.687 0.743				5.6
Noden 2014	Namibia 2011-12	ELISA (Panbio)	25/312	0.080	0.055 0.116				0.6
Clements 2019	Uganda 2006-07	ELISA (In house)	72/1744	0.041	0.033 0.052				1.9
Mohammed 2012	Puerto Rico 2006	ELISA (Not specified)	275/300	0.917	0.880 0.943				0.6
Slavov 2019	Brazil 2015-16	ELISA (Euroimmun)	20/475	0.042	0.027 0.064				0.5
Busch 2016	Brazil 2012	ELISA (F.D.)	453/498	0.910	0.881 0.932				1.1
Ribas-Silva 2012	Brazil	IC (I-R. D.)	3/213	0.014	0.005 0.043				0.1
De Almeida 2018	Brazil 2017	IC (K.B.)	6/298	0.020	0.009 0.044				0.2
Saba Villarroel 2018	Bolivia B. and S.C.* 2016-17	ELISA (Euroimmun)	155/168	0.923	0.871 0.955				0.3
Saba Villarroel 2018	Bolivia others 2016-17	ELISA (Euroimmun)	68/281	0.242	0.196 0.296				1.4
Rodriguez 2009	Mexico 2006-07	ELISA (Panbio)	472/800	0.590	0.556 0.624				5.3
L'Azu 2015	Guadeloupe & Martinique 2011	ELISA (Panbio)	732/783	0.935	0.915 0.950				1.3
Arcuri 2012	Argentina 2010-11	ELISA (Bisplot)	15/95	0.158	0.098 0.246				0.3
Arcuri 2012	Argentina 2010-11	ELISA (Bisplot)	6/286	0.021	0.010 0.046				0.2
Yamashiro 2004	Dominican Republic 2002	ELISA (D.A./F.T.)	987/1008	0.979	0.968 0.986				0.6
Golubic 2012	Croatia 1997-2007	ELISA (Not specified)	0/600	0.000	0.000 0.013				<0.0
Total			9258/38658	0.240	0.235 0.244				
Southeast Asia			2768/13595	0.204	0.197 0.211				
China			178/7831	0.023	0.019 0.026				
India			299/1149	0.260	0.235 0.286				
Oceania			1214/6917	0.176	0.167 0.185				
Australia			536/5809	0.092	0.085 0.100				
West Asia			792/6466	0.123	0.115 0.131				
Saudi Arabia			430/1194	0.360	0.333 0.387				
Africa			1292/5875	0.220	0.210 0.231				
Americas			3192/5205	0.613	0.600 0.626				
Brazil			482/1484	0.325	0.301 0.349				
Endemic or epidemic area			8952/26203	0.342	0.336 0.347				
Non-endemic or non-epidemic area			306/12455	0.025	0.022 0.027				
<i>Heterogeneity: I² 99.39</i>									

*Beni and Santa Cruz regions

CI: confidence interval; IC: immunochromatography;

Table III - Global prevalence of IgM antibodies against Dengue virus in blood donors, by population

Study ID	Population	Screening test	Positive/ Total	Prevalence	95% CI	Forest plot	Relative weight %
						0.00 0.10 0.20	
Zeng 2018	China 2015	ELISA (Panbio)	3/819	0.004	0.001 0.011		0.5
Chen 2015	China 2014	ELISA (Not specified)	71/3000	0.024	0.019 0.030		10.7
Gao 2017	China 2013-14	ELISA (Panbio)	1/685	0.004	0.007 0.008		0.9
Tsai 2018	Taiwan 2014	IC/ELISA (Not specified)	17/8000	0.002	0.001 0.003		2.6
Ranjan 2016	India 2012	ELISA (Novatec)	27/200	0.135	0.094 0.190		3.6
Chhabra 2013	India	IC (J. Mitra & Co)	2/380	0.005	0.001 0.021		0.3
Kulkarni 2018	India 2016-17	ELISA (Panbio)	31/520	0.060	0.042 0.084		4.5
Harif 2014	Malaysia 2009-10	ELISA (Panbio)	25/360	0.069	0.047 0.101		3.6
Mohamad 2016	Malaysia 2015	ELISA (Not specified)	0/126	0.000	0.000 0.060		0.1
Low 2015	Singapore 2009-10	ELISA (Panbio)	113/3995	0.028	0.024 0.034		16.9
Faddy 2013	Australia 2008-09	ELISA (Panbio)	24/10906	0.002	0.002 0.003		3.7
Faddy 2012	Australia 2008-09	ELISA (Panbio)	8/10024	<0.001	<0.001 0.002		1.2
Tezcan 2014	Turkey 2010-11	IC (Standar D.)	8/920	0.009	0.004 0.017		1.2
Ashshi 2015	Saudi Arabia 2014	ELISA (Panbio)	6/100	0.060	0.027 0.127		0.9
Ashshi 2017	Saudi Arabia 2015-16	ELISA (Panbio)	50/910	0.055	0.042 0.072		7.3
Slavov 2019	Brazil 2015-16	ELISA (Panbio)	32/475	0.067	0.048 0.094		4.6
Busch 2016	Brazil 2012	ELISA (Focus D.)	44/498	0.088	0.066 0.117		6.2
Ribas-Silva 2012	Brazil	IC (I-R. D.)	0/213	0.000	0.000 0.036		0.1
De Almeida 2018	Brazil 2017	IC (Katal B.)	5/298	0.017	0.007 0.040		0.8
De Carvalho 2010	Brazil 2008	ELISA (Panbio)	137/3000	0.046	0.039 0.054		20.1
Arellanos-Soto 2015	Mexico 2010-12	ELISA (Panbio)	53/2061	0.026	0.020 0.034		7.9
Rodriguez 2009	Mexico 2006-07	ELISA (Panbio)	16/800	0.020	0.012 0.032		2.4
Arcuri 2012	Argentina 2010-11	ELISA (Bispot)	0/286	0.000	0.000 0.027		0.1
Arcuri 2012	Argentina 2010-11	ELISA (Bispot)	0/95	0.000	0.000 0.078		0.1
Total			678/49671	0.014	0.013 0.015		
Southeast Asia			295/19085	0.016	0.014 0.017		
China			75/4504	0.017	0.013 0.020		
India			60/1100	0.055	0.041 0.068		
West Asia			64/1930	0.033	0.026 0.042		
Americas			287/7726	0.037	0.033 0.042		
Brazil			218/4484	0.049	0.042 0.055		

Heterogeneity P 96.67

CI: confidence interval; IC: immunochromatography; ELISA: enzyme-linked immunosorbent assay

Tests for IgM antibodies against DENV show a seroprevalence of 1.4% (95% CI: 1.3% to 1.5%), with individual studies reporting values ranging from 0% to 13.5% (95% CI: 1.3% to 1.5%); this top value was reported in greater Delhi (India) in 2012⁵⁰. By geographic region, the highest percentage of positive results was in the Americas, with a seroprevalence of 3.7% (95% CI: 3.3% to 4.2%). By country, the highest seroprevalence was reported in China, at 5.5% (95% CI: 4.1% to 6.8%). Most studies took place in regions where DENV is endemic and/or had epidemic outbreaks at the time (Table III).

The NAT showed an overall DENV viraemic rate of 0.2% (95% CI: 0.2% to 0.2%), with the highest results coming from Saudi Arabia in 2015 to 2016 (prevalence 5.5%, 95% CI: 4.2% to 7.2%)⁷¹. In the Americas, the prevalence was 0.2% (95% CI: 0.2% to 0.2%). The only three studies undertaken in non-endemic regions found a prevalence of 0.0% (Table IV). The highest prevalence was in Brazil, at 0.3% (95% CI: 0.3% to 0.3%).

Finally, several studies tested donors for the dengue NS1 antigen, which showed an overall prevalence of 0.2% (95% CI: 0.1% to 0.2%), with results in individual studies ranging from 0% to 5.3% (95% CI: 4.0% to 6.9%). These latter results came from Saudi Arabia in 2015 to 2016, in the screening

reported by Ashshi *et al.*⁷². By region, the Americas again led the ranking for the highest prevalence, with a pooled proportion of 0.1% (95% CI: 0.1% to 0.1%). All the studies took place in endemic regions or in areas with an epidemic outbreak (Table IV).

Prevalence of Zika virus

The overall seroprevalence of IgG antibodies against ZIKV was 5.1% (95% CI: 4.6% to 5.7%). The highest rate was in the donor screening programme in the Bolivian regions of Beni and Santa Cruz in 2016 to 2017, at 27.5% (95% CI: 22.8% to 32.8%)⁴⁰. By region, the highest seroprevalence was again in the Americas, at 7.4% (95% CI: 6.3% to 8.7%). The prevalence ratio between endemic and non-endemic regions was 9.0 (Table V).

NAT showed an overall prevalence of ZIKV of $0.7 \times 10^{-2}\%$ (95% CI: $0.7 \times 10^{-2}\%$ to $0.8 \times 10^{-2}\%$), varying from 0% to 2.8% (95% CI: 2.1% to 3.8%). The highest viraemic rate was recorded in a study in French Polynesia in 2013 to 2014²⁶. The country with the highest prevalence estimate for ZIKV was Brazil (0.5%, 95% CI: 0.4% to 0.7%). The studies in endemic populations, with local transmission or an epidemic outbreak, documented a prevalence of 0.1% (95% CI: 0.1% to 0.1%) (Table V).

Table IV - Global prevalence of Dengue virus in blood donors according to nucleic acid amplification and NS1 antigen, by population

Study ID	Population	Screening test	Positive/ Totals	Prevalence	95% CI		Forest plot		Relative weight %
NAT							0.00	0.05	0.10
Zeng 2018	China 2015	In house	0/1164	0.000	0.000	0.007			0.1
Liao 2017	China 2014	In house	2/3000	0.001	<0.001	0.003			0.4
Gao 2017	China 2013-14	In house	0/1685	0.000	0.000	0.005			0.1
Tsai 2018	Taiwan 2015	In house	1/8000	<0.001	0.000	0.001			0.2
Lin 2016	Taiwan 2015	LightMix Kit (T.B.)	16/6515	0.003	0.002	0.004			3.2
Lu 2018	Taiwan 2015	Procleix (Grifols)	21/5000	0.004	0.003	0.006			4.2
Ranjan 2016	India 2012	In house	0/200	0.000	0.000	0.039			0.1
Sayama 2013	Laos 2012	Not specified	0/199	0.000	0.000	0.039			0.1
Linnen 2008	Australia 2003	Procleix (Chiron)	0/5879	0.000	0.000	0.001			0.1
Linnen 2008	Brazil 2004-05	Procleix (Chiron)	9/2994	0.003	0.002	0.006			1.8
Linnen 2008	Honduras 2003	Procleix (Chiron)	3/4858	<0.001	<0.001	0.002			0.6
Rooks 2016	Australia 2008-09	Procleix (Hologic)	0/664	0.000	0.000	0.012			0.1
Rooks 2016	Australia 2012-13	Procleix (Hologic)	0/5518	0.000	0.000	0.001			0.1
Faddy 2015	Australia 2008-13	Not specified	0/6182	0.000	0.000	0.001			0.1
Beau 2020	French Polynesia 2013-18	RealStar (Altona)	5/34000	<0.001	<0.001	<0.001			1.0
Ashshi 2017	Saudi Arabia 2015-16	In house	50/910	0.055	0.042	0.072			9.4
Stramer 2019	USA 2015	Cobas test	0/10528	0.000	0.000	<0.001			0.1
Mohammed 2012	Puerto Rico 2005	NAT (Gen-Probe)	12/16521	<0.001	<0.001	0.001			2.4
Stramer 2013	Puerto Rico 2012-13	Not specified	114/49909	0.002	0.002	0.003			22.7
Stramer 2010	Puerto Rico 2007	NAT (Gen-Probe)	29/15350	0.002	0.001	0.003			5.8
Saá 2019	Puerto Rico 2015-16	Not specified	0/1186	0.000	0.000	0.007			0.1
Slavov 2019	Brazil 2015-16	In house	0/475	0.000	0.000	0.017			0.1
Slavov 2018	Brazil 2015-16	In house	1/631	0.002	<0.001	0.011			0.2
Sharma 2018	Brazil 2016	In house	0/676	0.000	0.000	0.012			0.1
Sabino 2013	Brazil 2012	Not specified	102/20132	0.005	0.004	0.006			20.3
Lavezzo 2010	Brazil 2006	In house	0/205	0.000	0.000	0.038			0.1
Dias 2012	Brazil 2010	In house	2/500	0.004	0.001	0.016			0.4
Busch 2016	Brazil 2012	Procleix (Hologic)	87/16241	0.005	0.004	0.007			17.3
Levi 2009	Brazil 2007-08	In house	0/23568	0.000	0.000	<0.001			0.1
De Almeida 2018	Brazil 2017	Not specified	0/298	0.000	0.000	0.026			0.1
Escoval 2013	Portugal 2012-13	Not specified	44/1948	0.023	0.017	0.030			8.6
Sargento 2017	Portugal	Not specified	0/110	0.000	0.000	0.068			0.1
Total			498/245046	0.002	0.002	0.002			
Southeast Asia			40/24599	0.002	0.002	0.001			
China			2/5849	<0.001	0.000	0.001			
Taiwan			38/19515	0.002	0.001	0.003			
Oceania			5/52243	<0.001	0.000	<0.001			
Australia			0/18243	0.000	0.000	0.000			
Americas			359/164072	0.002	0.002	0.002			
Puerto Rico			155/82966	0.002	0.002	0.002			
Brazil			201/65720	0.003	0.003	0.003			
Heterogeneity I ² 95.59									
NS1 antigen							0.00	0.05	0.10
Tsai 2018	Taiwan 2015	IC (Not specified)	0/8000	0.000	0.000	0.001			0.4
Lin 2016	Taiwan 2015	Platelia (Bio-Rad)	2/6515	<0.001	<0.001	0.001			1.5
Chhabra 2013	India	Not specified	0/380	0.000	0.000	0.021			0.4
Mangwana 2014	India 2013	Platelia (Bio-Rad)	0/1709	0.000	0.000	0.005			0.4
Jain 2019	India 2016	Not specified	2/369	0.005	0.001	0.021			1.5
Kulkarni 2018	India 2016-17	Microlisa (J. Mitra)	3/520	0.006	0.002	0.018			2.3
Mohamad 2016	Malaysia 2015	Not specified	0/126	0.000	0.000	0.060			0.4
Yusuf 2018	Malaysia 2016	Platelia (Bio-Rad)	0/374	0.000	0.000	0.021			0.4
Fellizar 2012	Philippines	Platelia (Bio-Rad)	3/158	0.019	0.006	0.057			2.3
Rooks 2016	Australia 2008-13	Platelia (Bio-Rad)	0/973	0.000	0.000	0.008			0.4
Flower 2011	Australia 2008-09	Platelia (Bio-Rad)	20/1087	0.018	0.012	0.028			15.0
Ashshi 2015	Saudi Arabia 2014	ELISA (Panbio)	1/100	0.010	0.001	0.068			0.8
Ashshi 2017	Saudi Arabia 2015-16	ELISA (Panbio)	48/910	0.053	0.040	0.069			34.7
Stramer 2011	Puerto Rico 2010	Platelia (Bio-Rad)	8/53019	<0.001	<0.001	0.000			6.1
Stramer 2011	Puerto Rico 2010	Platelia (Bio-Rad)	1/2837	<0.001	0.000	0.003			0.8
Stramer 2010	Puerto Rico 2007	Platelia (Bio-Rad)	3/4401	0.001	<0.001	0.002			2.3
Slavov 2019	Brazil 2015-16	Platelia (Bio-Rad)	0/475	0.000	0.000	0.017			0.4
Patafino 2009	Brazil 2007	Platelia (Bio-Rad)	1/4000	<0.001	0.000	0.002			0.8
De Carvalho 2010	Brazil 2008	Platelia (Bio-Rad)	39/3000	0.013	0.010	0.018			29.4
Total			131/88953	0.002	0.001	0.002			
Southeast Asia			10/18151	<0.001	<0.001	0.001			
India			5/2978	0.002	0.002	0.003			
Americas			52/67732	<0.001	0.001	0.001			
Puerto Rico			12/60257	<0.001	<0.001	<0.001			
Brazil			40/7475	0.005	0.004	0.007			
Heterogeneity I ² 95.59									

CI: confidence interval; IC: immunochromatography; NAT: nucleic acid test; ELISA: enzyme-linked immunosorbent assay

Table V - Global prevalence of Zika virus in blood donors, by population

Study ID	Population	Screening test	Positive/ Total	Prevalence	95% CI	Forest plot	Relative weight %
IgG antibodies							
Postorino 2019	Laos 2015	ELISA (Euroimmun)/VNT	16/359	0.045	0.028 0.072	0.00 0.20 0.40	5.1
Postorino 2019	Laos 2016	ELISA (Euroimmun)/VNT	68/687	0.099	0.079 0.124		20.3
Aubry 2015	French Polynesia 2011-13	Not specified	5/593	0.008	0.004 0.020		1.6
Gake 2017	Cameroon 2015	ELISA (Euroimmun)/VNT	53/1084	0.049	0.038 0.064		16.7
Nurtop 2020	Congo 2011	ELISA (Euroimmun)/VNT	7/386	0.018	0.009 0.038		2.3
Seruyange 2019	Rwanda 2015	ELISA (Euroimmun)/VNT	12/874	0.014	0.008 0.024		3.9
Seruyange 2019	Sweden	ELISA (Euroimmun)/VNT	0/215	0.000	0.000 0.036		0.2
Diarra 2020	Mali 2013	ELISA (Euroimmun)/VNT	47/637	0.074	0.056 0.097		14.4
Saba Villarroel 2018	Bolivia B. and S.C.* 2016-17	ELISA (Euroimmun)/VNT	84/305	0.275	0.228 0.328		20.1
Saba Villarroel 2018	Bolivia others 2016-17	ELISA (Euroimmun)/VNT	1/510	0.002	<0.000 0.014		0.3
Slavov 2019	Brazil 2010-15	ELISA (Euroimmun)/VNT	0/180	0.000	0.000 0.043		0.2
Slavov 2019	Brazil 2016	ELISA (Euroimmun)/VNT	19/320	0.059	0.038 0.091		5.9
Slavov 2019	Brazil 2017	ELISA (Euroimmun)/VNT	29/317	0.092	0.064 0.129		8.7
Diefenbach 2019	Brazil 2016-17	ELISA (Euroimmun)/VNT	1/182	0.006	<0.001 0.038		0.3
Total			342/6649	0.051	0.046 0.057		
Africa							
Americas							
Brazil							
Endemic or epidemic areas							
Non-endemic or non-epidemic areas							
Heterogeneity I² 95.28							
NAT							
Zheng 2019	China Shenzhen	Procleix (Grifols)	0/9309	0.000	0.000 <0.001	0.00 0.025 0.05	0.1
Lam 2017	Singapore 2016-17	Procleix (Not specified)	1/63144	<0.001	0.000 <0.001		0.2
Beau 2020	French Polynesia 2013-14	In house	42/1505	0.028	0.021 0.038		7.4
Slavov 2019	Brazil 2016	In house	0/475	0.000	0.000 0.017		0.1
Magnus 2018	Brazil 2016	Lancotti's & Pyke's	3/1857	0.002	<0.001 0.005		0.6
Benites 2019	Brazil 2015-2016	Multiplex (B-M. /F.)	4/3737	0.001	<0.001 0.003		0.7
Slavov 2017	Brazil 2015-16	In house	37/1393	0.027	0.019 0.036		6.6
Sharma 2018	Brazil 2016	Multiplex (Kit Biomol)	0/616	0.000	0.000 0.013		0.1
Gallian 2016	Martinique 2016	RealStar (Altona)	76/4129	0.018	0.015 0.023		13.6
Williamson 2018	Puerto Rico 2016	Cobas test (Roche)	339/52942	0.006	0.006 0.007		61.3
Saá 2019	Puerto Rico 2015-16	Cobas test (Roche)	1/1186	<0.001	<0.001 0.006		0.2
Pate 2017	USA 2016-17	Cobas test (Roche)	12/1776190	<0.001	<0.001 <0.001		2.2
Fedyk 2019	USA 2016-17	Cobas test (Roche)	4/92618	<0.001	<0.001 <0.001		0.7
Galel 2017	USA 2016-17	Cobas test (Roche)	14/358786	<0.001	<0.001 <0.001		2.6
Williamson 2017	USA 2016-17	Procleix (Hologic/Grifols)	10/933831	<0.001	<0.001 <0.001		1.8
Saá 2018	USA 2017	Procleix (Grifols)	9/4325889	<0.001	<0.001 <0.001		1.6
Covin 2017	USA 2016-17	Procleix (Hologic)	0/30493	0.000	0.000 <0.001		0.1
Diefenbach 2019	Brazil 2016-17	In house	0/500	0.000	0.000 0.016		0.1
Borena 2017	Austria 2016	RealStar (Altona)	0/1001	0.000	0.000 0.008		0.1
Total			552/7659601	<0.001	<0.001 <0.001		
Americas							
Brazil							
USA							
Endemic or epidemic area							
Non-endemic or non-epidemic areas							
Heterogeneity I² 99.23							

*Beni and Santa Cruz regions

CI: confidence interval; NAT: nucleic acid test; VNT: virus neutralization test

DISCUSSION

Since the advent of blood transfusions, patients' safety has been threatened by the transmission of infectious agents¹²⁵. Since the turn of the century, a high number of transfusion-transmitted arbovirus cases have been notified, in some instances ending in a fatal outcome for the patient². The chance that an asymptomatic but viraemic person donates blood is an important concern for transfusion safety and is a possibility for all of the three arboviruses studied. To understand the magnitude of the problem, it is essential to review the published literature reporting viraemic rates in blood donors. Our study updates, collates, and summarises all the notified and published data to date.

The viraemic rates of the three arboviruses in areas experiencing outbreaks were high according to NAT screening (from 1.9% for CHIKV to 5.5% for DENV and 2.8% for ZIKV)^{35,71,26}. Such donors are asymptomatic but infected, often with high levels of viraemia, so there is a real risk of transmission of these viruses via transfusion. NAT methods are expensive and complex, and they require a series of material and human resources that are not accessible in all settings. Health services in most countries do not make routine use of NAT assays capable of detecting these viruses during the donation process. No study in Africa used this screening technique. On the other hand, when NAT assays are used in areas with no outbreaks, the prevalence is practically zero. Consequently, it is

important to select the target population appropriately for these screening tests.

Assessing seroprevalence of different arboviruses is important for understanding population exposures in the past. High rates of exposure could be correlated to a greater number of infected and asymptomatic donors capable of transmitting the infection, so this could constitute a source of information on the magnitude of the problem. Moreover, as Liu *et al.* pointed out in their review, quantifying the seroprevalence of these viruses is of interest from an epidemiological point of view²². In some populations, blood donor screening is the only type of seroprevalence study that has been performed.

The seroprevalence of IgG antibodies against the three arboviruses was high, especially for CHIKV and DENV. In the case of CHIKV, we found the highest seroprevalence of IgG antibodies in sub-Saharan Africa, where periodic outbreaks have been recorded since the 1950s. Some of the most prominent occurred in the Republic of Tanzania in 1954, in the Democratic Republic of Congo in 1999 to 2000, and in Kenya in 2017. The high seroprevalence found in the regions of Beni and Santa Cruz (Bolivia)⁴⁰, areas with very specific climatic, environmental, and economic conditions, was also noteworthy. We found high seroprevalence rates for IgG against DENV in hyperendemic regions or where studies took place following extensive epidemic outbreaks. Fourteen studies reported seroprevalence rates of more than 50%, with several reporting rates over 90%. DENV has been producing epidemic outbreaks for more than 200 years. This long epidemiological trajectory has translated into its wider geographic dissemination and generally higher seroprevalence rates. The seroprevalence of IgG against ZIKV in blood donors is clearly the lowest for the three arboviruses studied, reflecting the very recent appearance of this virus, which has only caused significant outbreaks since about 2007. As with CHIKV, the highest seroprevalence was found in the Beni region of Bolivia, as well as in Laos and the São Paulo region in Brazil, where outbreaks have been registered since 2016^{40,106,114}. However, the seroprevalence in African blood donors is very low, indicating the limited transmission of the virus on this continent, in contrast to DENV and CHIKV. The areas in which seroprevalence

is highest have some similarities: a tropical climate with a clear, rainy season, abundant vegetation and water resources, and a low level of economic resources. All health centres and institutions should support efforts to reduce the risk of transmitting arboviruses through blood transfusions. A wide range of interventions could have an impact, from broad environmental policies directed at addressing the climate crisis or the use of water, agricultural, and forestry resources, to community-based environmental measures targeting vector control, improved conservation of wetlands and water resources, and improvements to health systems.

Blood transfusion centres also have a role to play: first, we should improve screening in potential blood donors using specific questions about the symptomatology of potential infections. It is also important that donors understand the symptoms of possible infections and are encouraged to report any they experience in the days and weeks following the donation. Secondly, it may be worth establishing a quarantine period for red blood cell concentrates, postponing their release until after the incubation period for infections has passed. Implementation of these measures requires adequate training among personnel working in donor selection or haemovigilance and co-responsibility among donors in terms of monitoring their own health. However, these measures would not enable identification of asymptomatic donors¹²⁶. The following measure would therefore be the suspension of blood donation collections in a region, as done during the 2007 CHIKV outbreak in Italy, although this measure is difficult to apply in low-resource areas¹²⁷. Donor screening (ideally using NAT) to detect a virus or its biomarkers is another possibility. When NAT is not available, one more affordable and accessible option of interest is point-of-care testing (immunoassay, reverse transcriptase polymerase chain reaction [RT-PCR], reverse-transcription loop-mediated amplification [RT-LAMP]), which has demonstrated an acceptable sensitivity and specificity for CHIKV, DENV, and ZIKV, respectively^{128,129,130}. Finally, where available, techniques for the deactivation of pathogens could also be applied, as these methods have proven effective against several different arboviruses^{131,132,133,134}.

Our study has several limitations, chief among which is the considerable heterogeneity of the diagnostic

tests used by different groups on the same virus (from commercial test kits to in-house techniques). These tests have different sensitivities and specificities. Moreover, in the case of DENV, different NAT assays could fail to detect some serotypes or genotypes in naïve populations. Most commercial techniques have an acceptable sensitivity for the four serotypes: MA assay Gen-Probe (limit of detection [LoD] 95% 14.9 copies/mL; specificity 99.91%), RealStar dengue RT-PCR assay Altona Diagnostic (sensitivity 83.2%, 95% CI: 77.6% to 89.1%), Cobas CHIKV/DENV test Roche Molecular Systems (LoD 95% 0.37 to 1.05 copies/mL, specificity 100%)^{104,135,33}. However, in-house techniques are more variable: some are capable of detecting all four serotypes with acceptable sensitivity, while others have been designed to detect only the serotype in circulation in the specific setting in which it is being used.

Another problem is cross-reactivity between different arboviruses. Although the highest prevalence of CHIKV was in Africa, most studies did not perform neutralisation tests or only performed them on a subsample of those yielding positive results. CHIKV shows cross-reactivity with other alphaviruses such as the O'nyong-nyong and Mayaro viruses. Clements *et al.* identified 552 donors with a positive result for CHIKV, but neutralisation tests were run in just 24; of these, 23 showed higher titres for O'nyong-nyong virus than for Chikungunya³¹. Thus, the results for prevalence of IgG antibodies against CHIKV should be interpreted with caution, especially in Africa, where other alphaviruses have been shown to circulate. Although DENV also shows some cross-reactivity with other flaviviruses, authors of the studies on this virus usually did perform neutralisation tests (normally to identify the DENV serotype). In the case of ZIKV, all the studies included virus neutralisation tests. We selected studies performed in the blood donor population in order to obtain data that are representative of that population. However, the results may not be applicable to the general population. Studies are not available in all geographic areas, and a substantial proportion have been in areas known to have high prevalence, which may lead to an overestimation of results. In addition, the between-study heterogeneity was quite high ($I^2 > 75\%$ in all cases). As a single review author selected the studies for inclusion, we cannot rule out the risk of selection bias. Moreover, some risk of publication bias is possible, as there may have

been unpublished studies finding negligent prevalence estimates. So, the external validity of the study may be limited by the real prevalence.

CONCLUSIONS

Our review has helped to elucidate the prevalence of CHIKV, DENV, and ZIKV in blood donors around the world, as determined by different screening tests. We have demonstrated that in regions where large epidemic outbreaks have occurred, the donor population has been widely exposed to the viruses, and the viraemic rate observed from donor screening may be high. This fact represents a threat to blood transfusion safety, so it is important that centres involved in these procedures understand the epidemiology driving the emergence of these transfusion-transmissible arbovirus infections.

Over the next few years, it is likely that the vector will expand into new settings, increasing the risk of outbreaks worldwide. The transmission of different arboviruses through transfusion will become a global threat. Institutions, authorities, blood transfusion centres, and blood banks should make efforts to design a clear path forward.

ACKNOWLEDGEMENTS

We express our thanks to Meggan Harris for her assistance in editing.

AUTHORSHIP CONTRIBUTIONS

All Authors contributed to the study design and final approval of manuscript.

The Authors declare no conflicts of interest.

REFERENCES

1. World Health Organization. Blood donor selection: guidelines on assessing donor suitability for blood donation. 2012. Available at: https://apps.who.int/iris/bitstream/handle/10665/76724/9789241548519_eng.pdf. accessed on 16/06/2020.
2. Petersen LR, Busch MP. Transfusion-transmitted arboviruses. *Vox Sang* 2010; **98**: 495-503.
3. American Association of Blood Banks. Chikungunya virus. *Transfusion* 2009; **49**(Suppl 2); 59S-61S.
4. Madariaga M, Ticona E, Resurreccion C. Chikungunya: bending over the Americas and the rest of the world. *Braz J Infect Dis* 2016; **20**: 91-8.
5. Autochthonous transmission of Chikungunya virus in EU/EEA, 2007–2017 [Internet]. European Centre for Disease Prevention and Control. Available at: <https://www.ecdc.europa.eu/en/all-topics-z/chikungunya-virus-disease/surveillance-and-disease-data/autochthonous-transmission>. Accessed on 16/06/2020.

6. Fiebre Chikungunya - Epidemiología y situación mundial [Internet]. Asociación de Médicos de Sanidad Exterior. Available at: <https://www.amse.es/informacion-epidemiologica/658-fiebre-chikungunya>. Accessed on 08/06/2021.
7. Chikungunya [Internet]. Centers for Disease Control and Prevention CDC Yellow Book Chapter 4. Available at: <https://wwwnc.cdc.gov/travel/yellowbook/2020/travel-related-infectious-diseases/chikungunya>. Accessed on 08/06/2021.
8. Parola P, De Lamballerie X, Jourdan J, et al. Novel Chikungunya virus variant in travelers returning from Indian Ocean islands. *Emerg Infect Dis* 2006; **12**: 1493-9.
9. Organización Panamericana de la Salud. Organización Mundial de la Salud. Dengue: Guías para el diagnóstico, tratamiento, prevención y control. La Paz, Bolivia. OPS/OMS 2010.
10. Gubler DJ. Dengue/dengue haemorrhagic fever: history and current status. *Novartis Found Symp* 2006; **277**: 3-16; discussion 16-22, 71-3, 251-3.
11. Dengue y Dengue grave [Internet]. World Health Organization 2020. Available at: <https://www.who.int/es/news-room/fact-sheets/detail/Dengue-and-severe-Dengue>. Accessed on 16/06/2021.
12. Autochthonous transmission of Dengue virus in EU/EEA, 2010/19 [Internet]. European Centre for Disease Prevention and Control. Available at: <https://www.ecdc.europa.eu/en/all-topics-z/Dengue/surveillance-and-disease-data/autochthonous-transmission-Dengue-virus-eueea>. Accessed on 16/06/2021.
13. Tambyah PA, Koay ES, Poon ML, et al. Transfusion-Transmitted Dengue Infection Study Group. Dengue hemorrhagic fever transmitted by blood transfusion. *N Engl J Med* 2008; **359**: 1526-7.
14. American Association of Blood Banks. Dengue viruses. *Transfusion* 2009; **49** (Suppl 2): 67S-9S.
15. Karim F, Nasir N, Moiz B. Transfusion transmitted Dengue: one donor infects two patients. *Transfus Apher Sci* 2017; **56**: 151-3.
16. Sabino EC, Loureiro P, Lopes ME, et al. Transfusion-transmitted dengue and associated clinical symptoms during the 2012 epidemic in Brazil. *J Infect Dis* 2016; **213**: 694-702.
17. Matos D, Tomashek KM, Perez-Padilla J, et al. Probable and possible transfusion-transmitted dengue associated with NS1 antigen-negative but RNA confirmed-positive red blood cells. *Transfusion*. 2016; **56**: 215-222.
18. Slavov SN, Otaguiri KK, Kashima S, et al. Overview of Zika virus (ZIKV) infection in regards to the Brazilian epidemic. *Braz J Med Biol Res* [Internet] 2016; **49**: e5420.
19. Epidemiological update: second case of locally acquired Zika virus disease in Hyères, France [Internet]. European Centre for Disease Prevention and Control. Available at: <https://www.ecdc.europa.eu/en/news-events/epidemiological-update-second-case-locally-acquired-zika-virus-disease-hyeres-france>. Accessed on 23/06/2021.
20. Barjas-Castro, ML, Angerami RN, Cunha MS, et al. Probable transfusion-transmitted Zika virus in Brazil. *Transfusion* 2016; **56**: 1684-8.
21. Motta LJ, Spencer BR, Cordeiro SG, et al. Evidence for transmission of Zika virus by platelet transfusion. *N Engl J Med* 2016; **375**: 1101-3.
22. Liu R, Wang X, Ma Y, et al. Prevalence of Zika virus in blood donations: a systematic review and meta-analysis. *BMC Infect Dis* 2019; **19**: 590.
23. Eick SM, Dale AP, McKay B, et al. Seroprevalence of Dengue and Zika virus in blood donations: a systematic review. *Transfus Med Rev* 2019; **33**: 35-42.
24. Sayama Y, Suksakhone C, Keokhamphoui C, et al. Prevalence of mosquito-borne viral infection (dengue fever, Japanese encephalitis, and chikungunya fever) among blood donors in Lao PDR. *Vox Sang* 2013; **105**: 98-9.
25. Appassakij H, Promwong C, Rujirojindakul P, et al. The risk of blood transfusion-associated Chikungunya fever during the 2009 epidemic in Songkhla Province, Thailand. *Transfusion* 2014; **54**: 1945-52.
26. Beau F, Lastère S, Mallet HP, et al. Impact on blood safety of the last arboviruses outbreaks in French Polynesia (2012-18). *Transfus Clin Biol* 2020; **27**: 4-9.
27. Humphrey JM, Al-Absi ES, Hamdan MM, et al. Dengue and chikungunya seroprevalence among Qatari nationals and immigrants residing in Qatar. *PLoS One* 2019; **14**: e0211574.
28. Seruyange E, Ljungberg K, Muvunyi CM, et al. Seroreactivity to Chikungunya and West Nile viruses in Rwandan blood donors. *Vector Borne Zoonotic Dis* 2019; **19**: 731-40.
29. Brouard C, Bernillon P, Quatresous I, et al. Estimated risk of Chikungunya viremic blood donation during an epidemic on Reunion Island in the Indian Ocean, 2005 to 2007. *Transfusion* 2008; **48**: 1333-41.
30. Moyer N, Thiberville SD, Pastorino B, et al. First reported chikungunya fever outbreak in the Republic of Congo, 2011. *PLoS One* 2014; **9**: e115938.
31. Clements TL, Rossi CA, Irish AK, et al. Chikungunya and O'nyong-nyong viruses in Uganda: implications for diagnostics. *Open Forum Infect Dis* 2019; **6**: ofz001.
32. Benites BD, Rocha D, Andrade E, et al. Zika virus and the safety of blood supply in Brazil: a retrospective epidemiological evaluation. *Am J Trop Med Hyg* 2019; **100**: 174-7.
33. Stramer SL, Stanley J, Nguyen ML, et al. Duplex nucleic acid test for the detection of chikungunya and dengue RNA viruses in blood donations. *Transfusion* 2019; **59**: 1283-90.
34. Chiu CY, Bres V, Yu G, et al. Genomic assays for identification of Chikungunya virus in blood donors, Puerto Rico, 2014. *Emerg Infect Dis* 2015; **21**: 1409-13.
35. Simmons G, Brès V, Lu K, et al. High incidence of Chikungunya virus and frequency of viremic blood donations during epidemic, Puerto Rico, USA, 2014. *Emerg Infect Dis* 2016; **22**: 1221-8.
36. Saa P, Chiu C, Grimm K, et al. Acute Zika virus infection in an asymptomatic blood donor at the onset of the Puerto Rico epidemic. *Transfusion* 2019; **59**: 3164-70.
37. Slavov SN, Otaguiri KK, Bianchini ML, et al. Seroprevalence of Chikungunya virus in blood donors from Northern and Southeastern Brazil. *Hematol Transfus Cell Ther* 2018; **40**: 358-62.
38. Sharma R, Costa Santos L, da Silva RA, et al. Surveillance of donated blood during the 2016 arbovirus outbreak in Brazil. *J Med Virol* 2018; **90**: 1406-10.
39. Gallian P, Leparç-Goffart I, Richard P, et al. Epidemiology of Chikungunya virus outbreaks in Guadeloupe and Martinique, 2014: an observational study in volunteer blood donors. *PLoS Negl Trop Dis* 2017; **11**: e0005254.
40. Saba Villarroel PM, Nurtop E, Pastorino B, et al. Zika virus epidemiology in Bolivia: a seroprevalence study in volunteer blood donors. *PLoS Negl Trop Dis* 2018; **12**: e0006239.
41. Sargento C, Silva E, Ferreira C, et al. Is chikungunya virus a blood transfusion problem in north Portugal? *Vox Sang* 2017; **112**: 168.
42. Zeng P, Liao Q, Gao Z, et al. Sero-prevalence and viremia status of dengue virus among asymptomatic blood donors post epidemic outbreak in Chinese Guangzhou in 2015. *Transfus Med* 2018; **28**: 468-9.
43. Liao Q, Shan Z, Wang M, et al. An evaluation of asymptomatic Dengue infections among blood donors during the 2014 Dengue outbreak in Guangzhou, China. *J Med Virol* 2017; **89**: 2037-40.
44. Chen JY, Liao Q, You RR, et al. Serum epidemiological investigation of dengue virus infection in blood donors from Guangzhou. *J Trop Med* 2015; **8**: 1014-6.
45. Kwan TH, Lee SS, Chan DPC, et al. Assessing the risk of dengue virus transmission in a non-endemic city surrounded by endemic and hyperendemic areas. *Int J Infect Dis* 2017; **55**: 99-101.
46. Gao Z, Zhang Y, Yang Y, et al. Dengue virus infections among blood donors in Guangxi of China, 2013-14. *Transfus Med* 2018; **28**: 236-42.
47. Tsai J-J, Lin P-C, Tsai C-Y, et al. Low frequency of asymptomatic dengue virus-infected donors in blood donor centers during the largest dengue outbreak in Taiwan. *PLoS One* 2018; **13**: e0205248.
48. Lin MH, Chen JW, Yeh CC. Lesson learned from the 2015 dengue outbreak in Taiwan. *Vox Sang* 2016; **111**: 74.
49. Lu C, Chen Y, Tsai M, et al. Contribution of alanine aminotransferase to prevent DENV transfusion infection among asymptomatic blood donors in Taiwan during 2015 dengue epidemic. *Vox Sang* 2018; **113**: 217.
50. Ranjan P, Natarajan V, Bajpai M, et al. High seroprevalence of dengue virus infection in blood donors from Delhi: a single centre study. *J Clin Diagn Res* 2016; **10**: DC08-DC10.
51. Chhabra C, David R, Singh Swaroop N, et al. Additional dengue testing helps in prevention transfusion transmitted infections-experience from a tertiary care hospital in North India. *Vox Sang* 2013; **105**: 165.
52. Mangwana S. Dengue viremia in blood donors in Northern India: challenges of emerging dengue outbreaks to blood transfusion safety. *Asian J Transfus Sci* 2015; **9**: 177-80.

53. Jain A, Jain S, Chowdhury N. Seroprevalence of dengue in blood donors in an outbreak: experience of a blood bank in north India. *Trop Doct* 2019; **49**: 212-5.
54. Kulkarni R, Tiraki D, Wani D, et al. Risk of transfusion-associated dengue: screening of blood donors from Pune, western India. *Transfusion* 2019; **59**: 458-62.
55. Hossain MA, Khatun M, Arjumand F, et al. Serologic evidence of dengue infection before onset of epidemic, Bangladesh. *Emerg Infect Dis* 2003; **9**: 1411-4.
56. Harif NF, Kader ZS, Joshi SR, et al. Seropositive status of dengue virus infection among blood donors in North Malaysia. *Asian J Transfus Sci* 2014; **8**: 64.
57. Mohamad NN. Prevalence of positive Dengue serology among the blood donors in Hospital University Sains Malaysia (HUSM): a preliminary study. *Malays J Pathol* 2016; **38**: 208.
58. Yusuf MFM, Fauzi HM, Noordin SS, et al. NS1 Dengue antigen among blood donors in two blood collection centers in North Malaysia. *Mal J Med Health Sci* 2018; **14**: 17-22.
59. Low SL, Lam S, Wong WY, et al. Dengue seroprevalence of healthy adults in Singapore: serosurvey among blood donors, 2009. *Am J Trop Med Hyg* 2015; **93**: 40-5.
60. Fellizar A. Dengue NS1 antigen detection among selected blood donors in a dengue-endemic province in Northern Philippines. *Vox Sang* 2012; **103**: 190-1.
61. Rooks K, Seed CR, Fryk JJ, et al. Mitigating the risk of transfusion-transmitted dengue in Australia. *J Blood Transfus* 2016; **2016**: 3059848.
62. Flower RLP, Fryk J, Hyland C, et al. Dengue fever viral exposure rates among Australian blood donors during local outbreaks. *Vox Sang* 2011; **101**: 96.
63. Faddy HM, Seed CR, Fryk JJ, et al. Implications of dengue outbreaks for blood supply, Australia. *Emerg Infect Dis* 2013; **19**: 787-9.
64. Faddy H, Fryk J, Hyland C, et al. Are Australian blood donors exposed to dengue virus during local outbreaks? *Vox Sang* 2012; **103**: 185.
65. Faddy HM, Fryk JJ, Seed CR, et al. The search for dengue virus in Australian blood donations during local outbreaks. *Vox Sang* 2015; **109**: 67.
66. Darcy A, Clothier H, Phillips D, et al. Solomon Islands dengue seroprevalence study—previous circulation of dengue confirmed. *P N G Med J* 2001; **44**: 43-7.
67. Aubry M, Finke J, Teissier A, et al. Seroprevalence of arboviruses among blood donors in French Polynesia, 2011/13. *Int J Infect Dis* 2015; **41**: 11-2.
68. Ergünay K, Saygan MB, Aydoğan S, et al. Investigation of dengue virus and yellow fever virus seropositivities in blood donors from Central/Northern Anatolia, Turkey. *Mikrobiyol Bul* 2010; **44**: 415-24.
69. Tezcan S, Kizildamar S, Ülger M, et al. Mersin İli Kan Donörlerinde Flavivirus Seroepidemiolojisi. *Mikrobiyol Bul* 2014; **48**: 606-17.
70. Ashshi AM. Serodetection of Dengue virus and its antibodies among blood donors in the western region of Saudi Arabia: a preliminary study. *Blood Transfus* 2015; **13**: 135-8.
71. Ashshi AM. The prevalence of dengue virus serotypes in asymptomatic blood donors reveals the emergence of serotype 4 in Saudi Arabia. *Viol J* 2017; **14**: 107.
72. Ashshi AM, Alghamdi S, El-Shemi AG, et al. Seroprevalence of asymptomatic dengue virus infection and its antibodies among healthy/eligible Saudi blood donors: findings from Holy Makkah City. *Virology (Auckl)* 2017; **8**: 1-5.
73. Jamjoom GA, Azhar EI, Kao MA, et al. Seroepidemiology of asymptomatic dengue virus infection in Jeddah, Saudi Arabia. *Virology (Auckl)* 2016; **7**: 1-7.
74. Aghaie A, Aaskov J, Chinikar S, et al. Frequency of dengue virus infection in blood donors in Sistan and Baluchestan province in Iran. *Transfus Apher Sci* 2014; **50**: 59-62.
75. Larrieu S, Alain Michault A, Polycarpe D, et al. Dengue outbreaks: a constant risk for Reunion Island. Results from a seroprevalence study among blood donors. *Trans R Soc Trop Med Hyg* 2014; **108**: 57-9.
76. Vairo F, Nicastrì E, Yussuf SM, et al. IgG against dengue virus in healthy blood donors, Zanzibar, Tanzania. *Emerg Infect Dis* 2014; **20**: 465-8.
77. Collenberg E, Ouedraogo T, Ganamé J, et al. Seroprevalence of six different viruses among pregnant women and blood donors in rural and urban Burkina Faso: a comparative analysis. *J Med Virol* 2006; **78**: 683-92.
78. Sawadogo S, Baguoya A, Youghbare F, et al. Seroprevalence and factors associated with IgG anti-DENV positivity in blood donors in Burkina Faso during the 2016 dengue outbreak and implications for blood supply. *Transfus Med* 2020; **30**: 37-45.
79. Noden BH, Musuu M, Aku-Akai L, et al. Risk assessment of flavivirus transmission in Namibia. *Acta Trop* 2014; **137**: 123-9.
80. Stramer SL, Foster GA, Boucher CB, et al. Comparing the yield of dengue viremic blood donations using NS1 antigen (AG) and NAT. *Transfusion* 2011; **51**: 4A.
81. Mohammed H, Tomashek KM, Stramer SL, et al. Prevalence of anti-dengue immunoglobulin G antibodies among American Red Cross blood donors in Puerto Rico, 2006. *Transfusion* 2012; **52**: 1652-6.
82. Mohammed H, Linnen JM, Muñoz-Jordán JL, et al. Dengue virus in blood donations, Puerto Rico, 2005. *Transfusion* 2008; **48**: 1348-54.
83. Stramer SL, Foster GA, Brodsky J, et al. Investigational dengue testing yields high rates of ribonucleic acid (RNA)-positive donors in Puerto Rico. *Transfusion* 2013; **53**: 215A.
84. Stramer SL, Linnen M, Carrick M, et al. Dengue donor viremia determined by RNA and NS1 antigen, and detection of dengue transfusion transmission during the 2007 dengue outbreak in Puerto Rico. *Vox Sang* 2010; **99**: 32.
85. Slavov SN, Cilião-Alves DC, Gonzaga FAC, et al. Dengue seroprevalence among asymptomatic blood donors during an epidemic outbreak in Central-West Brazil. *PLoS One* 2019; **14**: e0213793.
86. Slavov SN, Hespanhol MR, Ferreira AR, et al. Silent dengue virus circulation among asymptomatic blood donors from a hyperendemic Brazilian region. *Transfus Med* 2018; **28**: 465-7.
87. Sabino E, Loureiro P, Lopes M, et al. Dengue RNA among blood donors and recipients during large epidemics of DENV-4 in Rio de Janeiro and Recife, Brazil. *Vox Sang* 2013; **105**: 39.
88. Lavezzo LC, Dos Santos Santana V, Terzian AC, et al. Arboviruses in blood donors: a study in the Amazon region and in a small city with a dengue outbreak. *Transfus Med* 2010; **20**: 278-9.
89. Dias LL, Amarilla AA, Poloni TR, et al. Detection of dengue virus in sera of Brazilian blood donors. *Transfusion* 2012; **52**: 1667-71.
90. Busch MP, Sabino EC, Brambilla D, et al. Duration of dengue viremia in blood donors and relationships between donor viremia, infection incidence and clinical case reports during a large epidemic. *J Infect Dis* 2016; **214**: 49-54.
91. Ribas-Silva RC, Eid AA. Dengue antibodies in blood donors. *Rev Bras Hematol Hemoter* 2012; **34**: 193-5.
92. Levi J, Wendel SW, Rodrigues CL. Absence of dengue viremia in blood donors from São Paulo - Brazil. *Vox Sang* 2009; **97**: 124.
93. Ribas-Silva RC, Eid AA. Dengue antibodies in blood donors. *Rev Bras Hematol Hemoter* 2012; **34**: 193-5.
94. Patavino GM, Salles NA, Barreto AE, et al. Use of EIA antigen assay to screen blood donors for dengue virus. *Transfusion* 2009; **49**: 208A.
95. De Carvalho SC, Schatzmayr HS, Dos Santos F, et al. Dengue virus in blood donors from an endemic area in Brazil. *Vox Sang* 2010; **99**: 319.
96. Arellanos-Soto D, Cruz V, Mendoza-Tavera N, et al. Constant risk of dengue virus infection by blood transfusion in an endemic area in Mexico. *Transfus Med* 2015; **25**: 122-4.
97. Rodríguez D, Garza M, Chavarria AM, et al. Dengue virus antibodies in blood donors from an endemic area. *Transfus Med* 2009; **19**: 125-31.
98. L'Azou M, Jean-Marie J, Bessaud M, et al. Dengue seroprevalence in the French West Indies: a prospective study in adult blood donors. *Am J Trop Med Hyg* 2015; **92**: 1137-40.
99. Arcuri A, Salim J, Krakowiesky C, et al. Determination of IgG/IgM anti dengue virus antibodies in blood blanks of two different geographic areas of Argentina, the city of Buenos Aires vs the city of Santo Tome, Corrientes. *Vox Sang* 2012; **103**: 189.
100. Yamashiro T, Disla M, Petit A, et al. Seroprevalence of IgG specific for dengue virus among adults and children in Santo Domingo, Dominican Republic. *Am J Trop Med Hyg* 2004; **71**: 138-43.

101. Golubic D, Dobler G. Flaviviruses in the north-west Croatia. *Infektoloski Glasnik*. 2012; **32**: 153-7.
102. Escoval MA, Sousa G, Freitas B, et al. Dengue outbreak in Madeira Island (Portugal) blood safety measures. *Vox Sang* 2013; **105**: 192-3.
103. Sargento C, Ferreira C, Achando P, et al. Choosing a molecular test to screen dengue infection. *Vox Sang* 2017; **112**: 168-9.
104. Linnen JM, Vinelli E, Sabino EC, et al. Dengue viremia in blood donors from Honduras, Brazil, and Australia. *Transfusion* 2008; **48**: 1355-62.
105. Zheng X, Zeng J, Xu X, et al. A preliminary survey of Zika virus infection by nucleic acid test in the volunteer blood donor samples in Shenzhen China. *J Med Virol* 2020; **92**: 1326-9.
106. Pastorino B, Sengvilaipaseuth O, Chanthongthip A, et al. Low Zika virus seroprevalence in Vientiane, Laos, 2003-2015. *Am J Trop Med Hyg* 2019; **100**: 639-42.
107. Lam S, Chua SS, Stone M, et al. Universal Zika screening for blood donors in Singapore. *Transfusion* 2017; **57**: 27A.
108. Gake B, Vernet MA, Leparc-Goffart I, et al. Low seroprevalence of Zika virus in Cameroonian blood donors. *Braz J Infect Dis* 2017; **21**: 481-3.
109. Nurtop E, Moyén N, Dzia-Lepfoundzou A, et al. A report of Zika virus seroprevalence in Republic of the Congo. *Vector Borne Zoonotic Dis* 2020; **20**: 40-2.
110. Diarra I, Nurtop E, Sangaré A, et al. Zika virus circulation in Mali. *Emerg Infect Dis* 2020; **26**: 945-52.
111. Borena W, Hofer T, Stiasny K, et al. No molecular or serological evidence of Zikavirus infection among healthy blood donors living in or travelling to regions where *Aedes albopictus* circulates. *PLoS One* 2017; **12**: e0178175.
112. Slavov SN, Gonzaga FAC, Pimentel BMS, et al. Zika virus RNA surveillance in blood donors in the Federal District of Brazil during the 2016 outbreak. *Hematol Transfus Cell Ther* 2020; **42**: 394-6.
113. Magnus MM, Espósito DLA, da Costa VA, et al. Risk of Zika virus transmission by blood donations in Brazil. *Hematol Transfus Cell Ther* 2018; **40**: 250-254.
114. Slavov SN, Guaragna, Machado R, et al. Zika virus seroprevalence in blood donors from the Northeastern region of São Paulo State, Brazil, between 2015 and 2017. *J Infect* 2020; **80**: 111-115.
115. Slavov SN, Hespagnol MR, Rodrigues ES, et al. Zika virus RNA detection in asymptomatic blood donors during an outbreak in the northeast region of São Paulo State, Brazil, 2016. *Transfusion* 2017; **57**: 2897-901.
116. Gallian P, Cabié A, Richard P, et al. Zika virus in asymptomatic blood donors in Martinique. *Blood* 2017; **129**: 263-6.
117. Williamson P, Simmons G, Biggerstaff B, et al. Characterization of evolving viral and serological stages of ZIKV RNA positive blood donors and estimation of population incidence of infections during the Puerto Rico Zika epidemic, 2016. *Vox Sang* 2018; **113**: 81.
118. Pate LL, Williamson PC, Busch MP, et al. Detection of Zika virus RNA in United States blood donations using Cobas® Zika on the Cobas® 6800/8800 systems. *Transfusion* 2017; **57**: 26A.
119. Fedyk CG, Shahin G, Cap AP. Screening of donated blood products for Zika virus in US armed services blood program donors. *Transfusion* 2019; **59**: 54A-55A.
120. Galel SA, Williamson PC, Busch MP, et al. First Zika-positive donations in the continental United States. *Transfusion* 2017; **57**: 762-9.
121. Williamson P, Linnen J, Kessler D, et al. US blood donors with evidence of Zika virus infection outside areas of active transmission. *Vox Sang* 2017; **112**: 45.
122. Saá P, Proctor M, Foster G, et al. Investigational testing for Zika Virus among U.S. blood donors. *N Engl J Med* 2018; **378**: 1778-88.
123. Covin R, Nguyen K-AT. Zika virus donor screening-first year experience of a community blood center. *Transfusion* 2017; **57**: 205A-206A.
124. Diefenbach CF, Slavov SN, Kashima S, et al. Prevalence of Zika virus (ZIKV) in blood donors from a hemotherapy service of the southern region of Brazil. *Vox Sang* 2019; **14**: 157-62.
125. Perkins HA, Busch MP. Transfusion-associated infections: 50 years of relentless challenges and remarkable progress. *Transfusion* 2010; **50**: 2080-99.
126. Petersen LR, Epstein JS. Chikungunya virus: new risk to transfusion safety in the Americas. *Transfusion* 2014; **54**: 1911-5.
127. Liumbruno GM, Calteri D, Petropulacos K, et al. The Chikungunya epidemic in Italy and its repercussion on the blood system. *Blood Transfus* 2008; **6**: 199-210.
128. Arya SC, Agarwal N. Rapid point-of-care diagnosis of chikungunya virus infection. *Asian Pac J Trop Dis* 2011; **1**: 230-1.
129. Tsai JJ, Liu LT, Lin PC, et al. Validation of the Pockit dengue virus reagent set for rapid detection of dengue virus in human serum on a field-deployable PCR system. *J Clin Microbiol* 2018; **56**: e01865-17.
130. Mauk MG, Song J, Bau HH, et al. Point-of-care molecular test for Zika infection. *Clin Lab Int* 2017; **41**: 25-7.
131. Fryk JJ, Marks DC, Hobson-Peters J, et al. Dengue and chikungunya viruses in plasma are effectively inactivated after treatment with methylene blue and visible light. *Transfusion* 2016; **56**: 2278-85.
132. Wang Y, Ren K, Liao X, et al. Inactivation of Zika virus in plasma and derivatives by four different methods. *J Med Virol* 2019; **91**: 2059-65.
133. Faddy HM, Fryk JJ, Hall RA, et al. Inactivation of yellow fever virus in plasma after treatment with methylene blue and visible light and in platelet concentrates following treatment with ultraviolet C light. *Transfusion* 2019; **59**: 2223-7.
134. Papin JF, Floyd RA, Dittmer DP. Methylene blue photoinactivation abolishes West Nile virus infectivity in vivo. *Antiviral Res* 2005; **68**: 84-7.
135. Najioullah F, Viron F, Césaire R. Evaluation of four commercial real-time RT-PCR kits for the detection of dengue viruses in clinical samples. *Virol J* 2014; **11**: 164.