



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system

System
setting

Theory

Result

Summary

Exciton–Polaritons in Artificial Lattices and Electron Transport in Bose-Fermi Hybrid Systems

Meng Sun

Supervisors: Prof. Ivan Savenko Prof. Sergej Flach

University of Science & Technology
Institute of Basic Science, Center for the Theoretical Physics of Complex System

sunmeg.89@gmail.com

June 18, 2020 via Zoom

ibS Institute for Basic Science

pCS Center for Theoretical
Physics of Complex Systems

 UST

Overview

1 Exciton-Polaritons in Artificial lattices

- Introduction
- Phase selection and Solitons
- Localized condensates in Lieb lattice

2 Electron Transport in Bose-Fermi Hybrid Systems

- System setting
- Bloch–Grüneisen approach for resistivity
- Result

3 Summary

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system

System
setting

Theory

Result

Summary

Exciton–Polaritons in Artificial Lattices

Introduction

Introduction

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system


System
setting

Theory

Result

Summary

$$|Polariton\rangle = \alpha |Photon\rangle + \beta |Exciton\rangle$$



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system


System
setting

Theory

Result

Summary

$$|Polariton\rangle = \alpha |Photon\rangle + \beta |Exciton\rangle$$



A strong coupling between exciton and photon in microcavities

$$\begin{aligned} H_{pol} &= E_{cav} \hat{a}_k^\dagger \hat{a}_k + E_{exc} \hat{b}_k^\dagger \hat{b}_k \\ &+ \frac{\Omega}{2} (\hat{a}_k^\dagger \hat{b}_k + \hat{a}_k \hat{b}_k^\dagger) \end{aligned}$$

Introduction

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates



Hybrid
system

System
setting

Theory
Result

Summary

$$|Polariton\rangle = \alpha |Photon\rangle + \beta |Exciton\rangle$$



A strong coupling between exciton and photon in microcavities

$$\begin{aligned} H_{pol} &= E_{cav} \hat{a}_k^\dagger \hat{a}_k + E_{exc} \hat{b}_k^\dagger \hat{b}_k \\ &+ \frac{\Omega}{2} (\hat{a}_k^\dagger \hat{b}_k + \hat{a}_k \hat{b}_k^\dagger) \end{aligned}$$

Introduction

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun



EPs in
Artificial
lattices

Introduction

Phase
selectionLocalized
condensatesHybrid
systemSystem
settingTheory
Result

Summary

$$|Polariton\rangle = \alpha |Photon\rangle + \beta |Exciton\rangle$$



A strong coupling between exciton and photon in microcavities

■ High critical temperature T_c

$$\begin{aligned} H_{pol} &= E_{cav} \hat{a}_k^\dagger \hat{a}_k + E_{exc} \hat{b}_k^\dagger \hat{b}_k \\ &+ \frac{\Omega}{2} (\hat{a}_k^\dagger \hat{b}_k + \hat{a}_k \hat{b}_k^\dagger) \end{aligned}$$

Introduction

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates



Hybrid
system

System
setting

Theory
Result

Summary

$$|Polariton\rangle = \alpha |Photon\rangle + \beta |Exciton\rangle$$



A strong coupling between exciton and photon in microcavities

$$\begin{aligned} H_{pol} = & E_{cav} \hat{a}_k^\dagger \hat{a}_k + E_{exc} \hat{b}_k^\dagger \hat{b}_k \\ & + \frac{\Omega}{2} (\hat{a}_k^\dagger \hat{b}_k + \hat{a}_k \hat{b}_k^\dagger) \end{aligned}$$

- High critical temperature T_c
- Nice coherence properties

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates



Hybrid
system

System
setting

Theory
Result

Summary

$$|Polariton\rangle = \alpha |Photon\rangle + \beta |Exciton\rangle$$



A strong coupling between exciton and photon in microcavities

$$\begin{aligned} H_{pol} = & E_{cav} \hat{a}_k^\dagger \hat{a}_k + E_{exc} \hat{b}_k^\dagger \hat{b}_k \\ & + \frac{\Omega}{2} (\hat{a}_k^\dagger \hat{b}_k + \hat{a}_k \hat{b}_k^\dagger) \end{aligned}$$

- High critical temperature T_c
- Nice coherence properties
- Strong Nonlinear interaction

Introduction

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates



Hybrid
system

System
setting

Theory
Result

Summary

$$|Polariton\rangle = \alpha |Photon\rangle + \beta |Exciton\rangle$$



A strong coupling between exciton and photon in microcavities

$$\begin{aligned} H_{pol} = & E_{cav} \hat{a}_k^\dagger \hat{a}_k + E_{exc} \hat{b}_k^\dagger \hat{b}_k \\ & + \frac{\Omega}{2} (\hat{a}_k^\dagger \hat{b}_k + \hat{a}_k \hat{b}_k^\dagger) \end{aligned}$$

- High critical temperature T_c
- Nice coherence properties
- Strong Nonlinear interaction
- Driven-dissipative system

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun



EPs in
Artificial
lattices

Introduction

Phase
selectionLocalized
condensatesHybrid
systemSystem
settingTheory
Result

Summary

$$|Polariton\rangle = \alpha |Photon\rangle + \beta |Exciton\rangle$$



A strong coupling between exciton and photon in microcavities

$$\begin{aligned} H_{pol} = & E_{cav} \hat{a}_k^\dagger \hat{a}_k + E_{exc} \hat{b}_k^\dagger \hat{b}_k \\ & + \frac{\Omega}{2} (\hat{a}_k^\dagger \hat{b}_k + \hat{a}_k \hat{b}_k^\dagger) \end{aligned}$$

- High critical temperature T_c
- Nice coherence properties
- Strong Nonlinear interaction
- Driven-dissipative system
- Detunable

Detuning

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices


Introduction

Phase
selectionLocalized
condensatesHybrid
systemSystem
setting

Theory

Result

Summary

Detune parameter: $\delta = E_{cav}(k=0) - E_{exc}(k=0)$ 

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase selection

Localized
condensates

Hybrid system

System setting

Theory

Result

Summary

With the large scale of de Broglie wavelength, exciton-polaritons can be manipulated by micro-scale potentials.

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates







Hybrid
system

System
setting

Theory

Result

Summary



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
lattices

Introduction

Phase selection

Localized
condensates

Hybrid system

System setting

Theory

Summary

With the large scale of de Broglie wavelength, exciton-polaritons can be manipulated by micro-scale potentials.

Works Include:

- Multivalley condensation [1]
 - Phase selection and solitons [2]
 - Flat band condensation [3, 4]
 - Topological insulator [5]

Works Include:

- Multivalley condensation [1]
 - Phase selection and solitons [2]
 - Flat band condensation [3, 4]
 - Topological insulator [5]

The driven-dissipative Gross–Pitaevskii equation

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}, t) &= \left[-\frac{\hbar^2}{2m} \nabla^2 + V(\mathbf{r}) + \alpha |\psi(\mathbf{r}, t)|^2 - \frac{i\gamma}{2} \right] \psi(\mathbf{r}, t) \\ &\quad + \frac{i}{2} R \eta_R(\mathbf{r}, t) \psi(\mathbf{r}, t) + iP(\mathbf{r}, t), \\ \frac{\partial}{\partial t} \eta_R(\mathbf{r}, t) &= I(\mathbf{r}, t) - R \eta_R(\mathbf{r}, t) |\psi(\mathbf{r}, t)|^2 - \gamma_R \eta_R(\mathbf{r}, t). \end{aligned}$$

EPs in
Artificial
Lattices &
Hybrid
Systems
M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system


System
setting

Theory

Result

Summary

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}, t) = \left[-\frac{\hbar^2}{2m} \nabla^2 + V(\mathbf{r}) + \alpha |\psi(\mathbf{r}, t)|^2 \right] \psi(\mathbf{r}, t)$$



EPs in
Artificial
Lattices &
Hybrid
Systems
M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system


System
setting

Theory

Result

Summary

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}, t) = \left[-\frac{\hbar^2}{2m} \nabla^2 + V(\mathbf{r}) + \alpha |\psi(\mathbf{r}, t)|^2 - \frac{i\gamma}{2} \right] \psi(\mathbf{r}, t)$$



EPs in
Artificial
Lattices &
Hybrid
Systems
M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system


System
setting

Theory

Result

Summary

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}, t) = \left[-\frac{\hbar^2}{2m} \nabla^2 + V(\mathbf{r}) + \alpha |\psi(\mathbf{r}, t)|^2 - \frac{i\gamma}{2} \right] \psi(\mathbf{r}, t) + iP(\mathbf{r}, t)$$



EPs in
Artificial
Lattices &
Hybrid
Systems
M. Sun

EPs in
Artificial
lattices

Introduction
Phase
selection

Localized
condensates

Hybrid
system

System
setting

Theory


Result

Summary

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}, t) = \left[-\frac{\hbar^2}{2m} \nabla^2 + V(\mathbf{r}) + \alpha |\psi(\mathbf{r}, t)|^2 - \frac{i\gamma}{2} \right] \psi(\mathbf{r}, t)$$

$$+ \frac{i}{2} R\eta_R(\mathbf{r}, t) \psi(\mathbf{r}, t) + iP(\mathbf{r}, t),$$

$$\frac{\partial}{\partial t} \eta_R(\mathbf{r}, t) = I(\mathbf{r}, t) - R\eta_R(\mathbf{r}, t) |\psi(\mathbf{r}, t)|^2 - \gamma_R \eta_R(\mathbf{r}, t).$$



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system

System
setting

Theory

Result

Summary

Exciton–Polaritons in Artificial Lattices

Phase Selection and Solitons [2]

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
lattices

Introduction

Phase selection

Localized
condensates

Hybrid system





System setting

Theory

Result

$$i\hbar\partial_t\psi = -\frac{\hbar^2}{2m^*}\partial_x^2\psi + V(x)\psi + (\alpha - i\beta)|\psi|^2\psi$$

$$i\hbar\partial_t\psi = -\frac{\hbar^2}{2m^*}\partial_x^2\psi + V(x)\psi + (\alpha - i\beta)|\psi|^2\psi$$

(a) $\Lambda\Lambda$ -type(b) VV -type

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
lattices

Introduction

Phase selection

Localized condensates

Hybrid system

System setting

Theory

Result

ΛΛ-type

$$\frac{\alpha}{\beta} = 0$$

$$\frac{\alpha}{\beta} = 2$$

$$\frac{\alpha}{\beta} = 6$$

Summary

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
lattices

Introduction
Phase
selsection

Localized
condensates

Hybrid
system

System
setting

Theory

Result



Summary

ΛΛ-type

$$\frac{\alpha}{\beta} = 0$$

$$\frac{\alpha}{\beta} = 2$$

$$\frac{\alpha}{\beta} = 6$$



Solitons

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction
Phase
selection



Localized
condensates

Hybrid
system

System
setting

Theory
Result

Summary



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system

System
setting

Theory

Result

Summary

Exciton–Polaritons in Artificial Lattices

Localized condensates in Lieb lattice [3]

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates


Hybrid
system

System
setting

Theory

Result

Summary



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system



System
setting

Theory

Result

Summary

$$\hat{H} = \begin{pmatrix} -\frac{\nabla^2}{2m_c} + V(\mathbf{r}) & \Omega \\ \Omega & \delta - \frac{i}{2\tau_x} - \frac{\nabla^2}{2m_x} + \alpha_x |\chi|^2 \end{pmatrix}$$



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system

System
setting

Theory

Result

Summary

$$P(\mathbf{r}, t) = P_0 \frac{(x \pm iy)^2}{R^2} \exp\left[-\frac{r^2}{R^2} - i\omega_0 t\right] \theta(t)\theta(t_p - t).$$

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
lattices

Introduction

Phase
selsection

Localized
condensates

Hybrid
system


System
setting

Theory

Result

Summary

$$P(\mathbf{r}, t) = P_0 \frac{(x \pm iy)^2}{R^2} \exp\left[-\frac{r^2}{R^2} - i\omega_0 t\right] \theta(t)\theta(t_p - t).$$



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system


System
setting

Theory

Result

Summary

$$i \begin{pmatrix} \dot{\varphi} \\ \dot{\chi} \end{pmatrix} = \hat{H} \begin{pmatrix} \varphi \\ \chi \end{pmatrix} + \begin{pmatrix} iP(\mathbf{r}, t) \\ 0 \end{pmatrix}.$$



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system


System
setting

Theory

Result

Summary

$$i \begin{pmatrix} \dot{\varphi} \\ \dot{\chi} \end{pmatrix} = \hat{H} \begin{pmatrix} \varphi \\ \chi \end{pmatrix} + \begin{pmatrix} iP(\mathbf{r}, t) \\ 0 \end{pmatrix}.$$



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
lattices

Introduction

Phase
selsection

Localized
condensates

Hybrid
system


System
setting

Theory

Result

Summary

$$i \begin{pmatrix} \dot{\varphi} \\ \dot{\chi} \end{pmatrix} = \hat{H} \begin{pmatrix} \varphi \\ \chi \end{pmatrix} + \frac{icn_r}{2} \begin{pmatrix} 0 \\ \chi \end{pmatrix} + \begin{pmatrix} iP(\mathbf{r}, t) \\ 0 \end{pmatrix},$$
$$\frac{\partial n_r}{\partial t} = I - \tau_r^{-1} n_r - c|\chi|^2 n_r.$$



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system

System
setting

Theory
Result

Summary

Electron Transport in Bose-Fermi Hybrid Systems

Bogolon–Mediated Electron Scattering [6, 7]

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase selection

Localized
condensates

Hybrid system

System setting

Theory

Result

- A layer of fermions (graphene or normal metal)
 - A layer of bosons in condensate (indirect-exciton)
 - Coulomb interaction between fermion and boson layers

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system


System
setting

Theory

Result

Summary

- A layer of fermions (graphene or normal metal)
- A layer of bosons in condensate (indirect-exciton)
- Coulomb interaction between fermion and boson layers



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
lattices

Introduction
Phase
selsection

Localized
condensates

Hybrid
system


System
setting

Theory

Result

Summary

$$g(r) = \frac{e_0^2}{4\pi\epsilon} \left(\frac{1}{r_{ee}} - \frac{1}{r_{eh}} \right)$$
$$g(k) = \frac{e_0^2}{2k\epsilon} e^{-kl} \left(1 - e^{-kd} \right)$$



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
lattices

Introduction

Phase selection

Localized condensates

Hybrid system

System
setting

Theory

Result

Electron-exciton interaction

$$V = \int d\mathbf{r} \int d\mathbf{R} \Psi_r^\dagger \Psi_r g(\mathbf{r} - \mathbf{R}) \Phi_R^\dagger \Phi_R$$

Summary

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifica
lattices

Introduction

Phase selection

Localized condensates

Hybrid system

System setting

Theory

Result

Electron-exciton interaction

$$V = \int d\mathbf{r} \int d\mathbf{R} \Psi_{\mathbf{r}}^{\dagger} \Psi_{\mathbf{r}} g(\mathbf{r} - \mathbf{R}) \Phi_{\mathbf{R}}^{\dagger} \Phi_{\mathbf{R}}$$

Excitons in BEC

$$\Phi_R = \sqrt{n_c} + \phi_R$$

Summary

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction
Phase
selection

Localized
condensates

Hybrid
system

System
setting

Theory

Result

Summary

Electron-exciton interaction

$$V = \int d\mathbf{r} \int d\mathbf{R} \Psi_{\mathbf{r}}^{\dagger} \Psi_{\mathbf{r}} g(\mathbf{r} - \mathbf{R}) \Phi_{\mathbf{R}}^{\dagger} \Phi_{\mathbf{R}}$$

Excitons in BEC

$$\Phi_{\mathbf{R}} = \sqrt{n_c} + \phi_{\mathbf{R}}$$

$$\begin{aligned} V_1 &= \sqrt{n_c} \int d\mathbf{r} \Psi_{\mathbf{r}}^{\dagger} \Psi_{\mathbf{r}} \int d\mathbf{R} g(\mathbf{r} - \mathbf{R}) [\phi_{\mathbf{R}}^{\dagger} + \phi_{\mathbf{R}}] \\ V_2 &= \int d\mathbf{r} \Psi_{\mathbf{r}}^{\dagger} \Psi_{\mathbf{r}} \int d\mathbf{R} g(\mathbf{r} - \mathbf{R}) \phi_{\mathbf{R}}^{\dagger} \phi_{\mathbf{R}} \end{aligned}$$

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
lattices

Introduction
Phase
selsection

Localized
condensates

Hybrid
system

System
setting

Theory

Result

Summary

Bogoliubov transformation

$$\phi_p = u_p b_p + v_p b_{-p}^\dagger, \quad \phi_p^\dagger = u_p b_p^\dagger + v_p b_{-p}$$

$$u_p^2 = 1 + v_p^2 = \frac{1}{2} \left(1 + \sqrt{1 + \frac{M^2 s^4}{\omega_p^2}} \right), \quad u_p v_p = -\frac{Ms^2}{2\omega_p}$$

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
lattices

Introduction
Phase
selsection

Localized
condensates

Hybrid
system

System
setting

Theory
Result

Summary

Bogoliubov transformation

$$\phi_p = u_p b_p + v_p b_{-p}^\dagger, \quad \phi_p^\dagger = u_p b_p^\dagger + v_p b_{-p}$$

$$u_p^2 = 1 + v_p^2 = \frac{1}{2} \left(1 + \sqrt{1 + \frac{M^2 s^4}{\omega_p^2}} \right), \quad u_p v_p = -\frac{Ms^2}{2\omega_p}$$

One- and two-bogolon interaction

$$V^{(1)} = \frac{\sqrt{n_c}}{L} \sum_{\mathbf{k}, \mathbf{p}} g_p \left[(v_p + u_{-\mathbf{p}}) b_{-\mathbf{p}}^\dagger + (u_p + v_{-\mathbf{p}}) b_{\mathbf{p}} \right] c_{\mathbf{k}+\mathbf{p}}^\dagger c_{\mathbf{k}}$$

$$V^{(2)} = \frac{1}{L^2} \sum_{\mathbf{k}, \mathbf{p}, \mathbf{q}} g_p \left[u_{\mathbf{q}-\mathbf{p}} u_{\mathbf{q}} b_{\mathbf{q}-\mathbf{p}}^\dagger b_{\mathbf{q}} + u_{\mathbf{q}-\mathbf{p}} v_{\mathbf{q}} b_{\mathbf{q}-\mathbf{p}}^\dagger b_{-\mathbf{q}}^\dagger \right.$$

$$+ \left. v_{\mathbf{q}-\mathbf{p}} u_{\mathbf{q}} b_{-\mathbf{q}+\mathbf{p}} b_{\mathbf{q}} + v_{\mathbf{q}-\mathbf{p}} v_{\mathbf{q}} b_{-\mathbf{q}+\mathbf{p}} b_{-\mathbf{q}}^\dagger \right] c_{\mathbf{k}+\mathbf{p}}^\dagger c_{\mathbf{k}}$$

EPs in
Artificial
Lattices &
Hybrid
Systems
M. Sun

EPs in
Artifical
lattices

Introduction
Phase
selsection
Localized
condensates


Hybrid
system

System
setting

Theory

Result

Summary



Boltzmann equation

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
lattices

Introduction

Phase selection

Localized
condensates

Hybrid system

System setting

Theory

Result

Summary

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction
Phase
selection
Localized
condensates

Hybrid
system

System
setting

Theory
Result

Summary

Boltzmann equation

$$e_0 \mathbf{E} \frac{\partial f}{\hbar \partial \mathbf{p}} = I\{f\}$$

Collision integral

$$\begin{aligned} I\{f\} = & -\frac{1}{\hbar} \int \frac{d\mathbf{q} d\mathbf{p}'}{(2\pi)^2} |V_q^{(1)}|^2 \\ & \times \left[N_q f_p (1 - f_{p'}) \delta(\varepsilon_p - \varepsilon_{p'} + \hbar\omega_q) \delta(\mathbf{p} - \mathbf{p}' + \mathbf{q}) \right. \\ & + (N_q + 1) f_p (1 - f_{p'}) \delta(\varepsilon_p - \varepsilon_{p'} - \hbar\omega_q) \delta(\mathbf{p} - \mathbf{p}' - \mathbf{q}) \\ & - N_q f_{p'} (1 - f_p) \delta(\varepsilon_{p'} - \varepsilon_p + \hbar\omega_q) \delta(\mathbf{p}' - \mathbf{p} + \mathbf{q}) \\ & \left. - (N_q + 1) f_{p'} (1 - f_p) \delta(\varepsilon_{p'} - \varepsilon_p - \hbar\omega_q) \delta(\mathbf{p}' - \mathbf{p} - \mathbf{q}) \right] \end{aligned}$$

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system

System
setting

Theory

Result

Summary

Boltzmann equation

$$e_0 \mathbf{E} \frac{\partial f}{\hbar \partial \mathbf{p}} = I\{f\}$$

Weak electric field

$$f = f^0(\varepsilon_p) + \left(\frac{\partial f^0}{\partial \varepsilon_p} \right) \phi_p$$

$$\phi_p = e_0 E_x \tau(\varepsilon_p) \frac{\partial \varepsilon_p}{\hbar \partial p_x}$$

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system

System
setting

Theory

Result

Summary

Boltzmann equation

$$e_0 \mathbf{E} \frac{\partial f}{\hbar \partial \mathbf{p}} = I\{f\}$$

Weak electric field

$$f = f^0(\varepsilon_p) + \left(\frac{\partial f^0}{\partial \varepsilon_p} \right) \phi_p$$

$$\phi_p = e_0 E_x \tau(\varepsilon_p) \frac{\partial \varepsilon_p}{\hbar \partial p_x}$$

Resistivity

$$\rho^{-1} = e_0^2 D(\varepsilon_f) \frac{v_f^2}{2} \langle \tau \rangle$$

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selectionLocalized
condensatesHybrid
systemSystem
setting


Theory

Result

Summary

$$\rho \propto T^4 \quad (T \ll T_{BG}), \quad \rho \propto T^1 \quad (T \gg T_{BG})$$

$$T_{BG} = \frac{2\hbar s k_f}{k_B}$$




EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifical
latticesIntroduction
Phase
selsectionLocalized
condensatesHybrid
systemSystem
settingTheory
Result

Summary



EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifica
lattices

Introduction

Phase

selection

Localized
condensates

Hybrid
systems

System
setting

Theory

Result

Phase selection

- Condensate: $0\text{-phase} \Leftrightarrow \pi\text{-phase}$
 - Dark soliton

Lieb lattice

- Exciton-polariton dispersion in Lieb lattice
 - Excite the CLS

Hybrid system

- Temperature dependent resistivity
 - Two-bogolon dominant in parabolic case

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artifica
lattices

Introduction

Phase

selection

Hybrid system

System settings

Theory

Result

Summary

- [1] M. Sun, I. G. Savenko, H. Flayac, and T. C. H. Liew.
Multivalley engineering in semiconductor microcavities.
Scientific Reports, 7:45243 EP –, 04 2017.
 - [2] Sukjin Yoon, Meng Sun, Yuri G. Rubo, and Ivan G. Savenko.
Phase selection and intermittency of exciton-polariton condensates in one-dimensional periodic structures.
Phys. Rev. A, 100:023609, Aug 2019.
 - [3] Meng Sun, I. G. Savenko, S. Flach, and Y. G. Rubo.
Excitation of localized condensates in the flat band of the exciton-polariton lieb lattice.
Phys. Rev. B, 98:161204, Oct 2018.
 - [4] Dogyun Ko, Meng Sun, Alexei Andreeanov, Y. G. Rubo, and I. G. Savenko.
Partial quantum revivals of localized condensates in distorted lattices.
Opt. Lett., 45(6):1571–1574, Mar 2020.
 - [5] M. Sun, D. Ko, D. Leykam, V. M. Kovalev, and I. G. Savenko.
Exciton-polariton topological insulator with an array of magnetic dots.
Phys. Rev. Applied, 12:064028, Dec 2019.
 - [6] Meng Sun, K. H. A. Villegas, V. M. Kovalev, and I. G. Savenko.
Bogolon-mediated electron scattering in graphene in hybrid bose-fermi systems.
Phys. Rev. B, 99:115408, Mar 2019.
 - [7] K. H. A. Villegas, Meng Sun, V. M. Kovalev, and I. G. Savenko.
Unconventional bloch-grüneisen scattering in hybrid bose-fermi systems.
Phys. Rev. Lett., 123:095301, Aug 2019.

EPs in
Artificial
Lattices &
Hybrid
Systems

M. Sun

EPs in
Artificial
lattices

Introduction

Phase
selection

Localized
condensates

Hybrid
system

System
setting

Theory

Result

Summary



