

Name: Key PID: _____

TA: _____ Sec. No: _____ Sec. Time: _____

Math 18

Exam 2 v. 0: Practice

March 2, 2018

Turn off and put away your cell phone and other electronic devices.

You may use one hand-written standard sheet of notes, but calculators, notes, and other assistance are prohibited during this exam.

Read each question carefully, and answer each question completely.

Show all of your work; no credit will be given for unsupported answers.

Write your solutions clearly and legibly; no credit will be given for illegible solutions.

If any question is not clear, ask for clarification.

1. Compute the determinant

$$\begin{vmatrix} -2 & -1 & -1 & 0 \\ 4 & 3 & 3 & 2 \\ -4 & -6 & -7 & 0 \\ -1 & -5 & -9 & 0 \end{vmatrix}$$

We'll use cofactor expansion for the 4th column:

The desired determinant is equal to:

$$2 \cdot (-1)^{2+4} \det \begin{bmatrix} -2 & -1 & -1 \\ -4 & -6 & -7 \\ -1 & -5 & -9 \end{bmatrix}$$

$$= 2 \left(-2 \begin{vmatrix} -6 & -7 \\ -5 & -9 \end{vmatrix} + \begin{vmatrix} -4 & -7 \\ -1 & -9 \end{vmatrix} - \begin{vmatrix} -4 & -6 \\ -1 & -5 \end{vmatrix} \right)$$

$$= 2 \left[-2(54 - 35) + (36 - 7) - (20 - 6) \right]$$

$$= 2(-2 \cdot 19 + 29 - 14)$$

$$= 2(-23)$$

$$= -46.$$

2. Let $B = \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}, \begin{pmatrix} -2 \\ -5 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ 5 \\ -1 \end{pmatrix} \right\}$.

(a) Explain why B is a basis of \mathbb{R}^3 .

$$\begin{bmatrix} 1 & -2 & 2 \\ 2 & -5 & 5 \\ 1 & 2 & -1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 2 \\ 0 & -1 & 1 \\ 0 & 4 & -3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 2 \\ 0 & -1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

\Rightarrow these vectors are linearly independent, and the matrix formed by them is invertible \Rightarrow they form a basis for \mathbb{R}^3 .

(b) Let $x = \begin{pmatrix} -3 \\ -8 \\ -2 \end{pmatrix}$. Find the coordinate vector of x relative to B .

We need to find x_1, x_2 , and x_3 such that

$$\begin{bmatrix} -3 \\ -8 \\ -2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} -2 \\ -5 \\ 2 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ 5 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 & -2 & 2 \\ 2 & -5 & 5 \\ 1 & 2 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$\left[\begin{array}{ccc|c} 1 & -2 & 2 & -3 \\ 2 & -5 & 5 & -8 \\ 1 & 2 & -1 & -2 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & -2 & 2 & -3 \\ 0 & -1 & 1 & -2 \\ 0 & 4 & -3 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & -2 & 2 & -3 \\ 0 & -1 & 1 & -2 \\ 0 & 0 & 1 & -7 \end{array} \right]$$

$$\rightarrow \left[\begin{array}{ccc|c} 1 & -2 & 2 & -3 \\ 0 & 1 & 0 & -5 \\ 0 & 0 & 1 & -7 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -5 \\ 0 & 0 & 1 & -7 \end{array} \right]$$

$$\Rightarrow x_1 = 1 \quad x_2 = -5, \quad x_3 = -7.$$

$$\therefore \left[\begin{pmatrix} -3 \\ -8 \\ -2 \end{pmatrix} \right]_B = \begin{bmatrix} 1 \\ -5 \\ -7 \end{bmatrix}.$$

3. The matrix

$$A = \begin{pmatrix} | & | & & | \\ \mathbf{a}_1 & \mathbf{a}_2 & \dots & \mathbf{a}_6 \\ | & | & & | \end{pmatrix} = \begin{pmatrix} 1 & 0 & -2 & 4 & -2 & -4 \\ 1 & 0 & 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 5 & 0 & -1 \\ 2 & -2 & 3 & -6 & 2 & 6 \end{pmatrix}$$

wrong matrix.
But it's still ok!
Suppose we only know RREF(A)

is row equivalent to

$$B = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & -1 \\ 0 & 1 & 0 & 4 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 & 1 & -2 \end{pmatrix}$$

Let H be the subspace of \mathbb{R}^4 spanned by $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$ and K be the subspace of \mathbb{R}^4 spanned by $\mathbf{a}_4, \mathbf{a}_5, \mathbf{a}_6$. Find a basis for each of the following subspaces.

(a) Col A, the column space of A.

pivots are columns 1, 2, 3, and 5.

$\Rightarrow \{\vec{a}_1, \vec{a}_2, \vec{a}_3, \vec{a}_5\}$ form a basis for Col(A).

(b) Nul A, the null space of A.


We can use B to find Nul(A). Free variables are x_4 and x_6 .

Solution of $A\vec{x} = \vec{0}$ is

$$\Rightarrow \vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = x_4 \begin{bmatrix} -1 \\ -4 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_6 \begin{bmatrix} 1 \\ 0 \\ -4 \\ 0 \\ 2 \\ 1 \end{bmatrix}$$


these two vectors form a basis for Nul(A)

(c) $H + K$, the sum of subspaces H and K .



has not learned this yet.

(d) $H \cap K$, the intersection of H and K .



4. Let A be a 7×10 matrix. We also know that $\text{Nul } A$ can be written as the linear span of b_1, b_2, b_3, b_4 , but we do not know if b_1, b_2, b_3, b_4 are linearly independent or not.

(a) Find all possible value(s) of $\dim \text{Nul } A$, the dimension of the null space of A .

$$\text{Nul}(A) = \text{span}\{\vec{b}_1, \vec{b}_2, \vec{b}_3, \vec{b}_4\}. \Rightarrow \dim \text{Nul}(A) \leq 4.$$

Since A is of size 7×10 , A has no more than 7 pivots.

$$\Rightarrow \text{rank}(A) \leq 7. \Rightarrow 10 - \text{nullity}(A) \leq 7.$$

$$\text{nullity}(A) \geq 3.$$

$$\Rightarrow \dim \text{Nul}(A) = 3 \text{ or } 4.$$

(b) Find all possible value(s) of $\text{rank } A$, the rank of A .

From part a), we know that $\text{rank}(A) \leq 7$,
Since $\dim \text{Nul}(A) \leq 4$,

$$10 - \text{rank}(A) \leq 4.$$

$$\text{rank}(A) \geq 6.$$

$$\Rightarrow \text{rank}(A) = 6 \text{ or } 7.$$

Midterm 2vS

Math 18, Section A

May 19, 2017

Time Limit: 50 Minutes

Name (Print): _____

PID: _____

- **DO NOT** begin working, or even open this packet, until instructed to do so.
- You should be in your assigned seat, unless instructed otherwise by Ed or one of the TAs.
- Enter all requested information on the top of this page, and put your name on the top of every page, in case the pages become separated.
- You may use a two-sided page of notes on this exam.
- You may **not** use your books, additional notes, or any calculator on this exam.

You are required to show your work on each problem on this exam. The following rules apply:

- **Organize your work**, in a reasonably neat and coherent way, in the space provided. Work scattered all over the page without a clear ordering will receive very little credit.
- **Mysterious or unsupported answers will not receive full credit.** Unless otherwise directed in the statement of the problem, a correct answer, unsupported by calculations, explanation, or algebraic work will receive little or no credit. An incorrect answer supported by substantially correct calculations and explanations might still receive partial credit.

Question	Points
1	10
2	10
3	10
4	10
Total:	40

DO NOT turn this page until instructed to do so

1. Let $A = \begin{bmatrix} 1 & 0 & 3 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}$ and let a, b, c, d be some numbers.

(a) (7 points) Find A^{-1} .

$$\left[\begin{array}{cccc|cccc} 1 & 0 & 3 & 0 & 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_3 \leftrightarrow R_4} \left[\begin{array}{cccc|cccc} 1 & 0 & 3 & 0 & 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \right]$$

$$\xrightarrow{1/2 R_2} \left[\begin{array}{cccc|cccc} 1 & 0 & 3 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1/2 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \right] \xrightarrow{R_3 - R_2} \left[\begin{array}{cccc|cccc} 1 & 0 & 3 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1/2 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & -1/2 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \right]$$

$$\xrightarrow{R_1 - 3R_3} \left[\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 1 & 3/2 & 0 & -3 \\ 0 & 1 & 0 & 0 & 0 & 1/2 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & -1/2 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \right] \Rightarrow A^{-1} = \begin{bmatrix} 1 & 3/2 & 0 & -3 \\ 0 & 1/2 & 0 & 0 \\ 0 & -1/2 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

(b) (3 points) Solve the equation $A\vec{x} = (a, b, c, d)$. You may answer in terms of a, b, c and d .

Since A is invertible,

$$\vec{x} = A^{-1} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} a + 3/2b - 3d \\ \frac{1}{2}b \\ -\frac{1}{2}b + d \\ c \end{bmatrix}$$

2. (10 points) Let \mathbb{P}_2 be the vector space of polynomials with degree at most 2, and let $\mathcal{B} = \{1+x, 1+x^2, x+x^2\}$. \mathcal{B} is a basis for \mathbb{P}_2 (and you don't need to prove this). Let $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ satisfy $T([a_0 + a_1x + a_2x^2]_{\mathcal{B}}) = (a_0, a_1, a_2)$. Find the matrix corresponding to T .

We need to find $T(\vec{e}_1)$, $T(\vec{e}_2)$, $T(\vec{e}_3)$.

We need to find a polynomial $a_0 + a_1x + a_2x^2$ such that

$$[a_0 + a_1x + a_2x^2]_{\mathcal{B}} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}.$$

$$\Rightarrow [1+x]_{\mathcal{B}} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}.$$

$$\Rightarrow T\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}\right) = T([1+x]_{\mathcal{B}}) = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}.$$

similarly, $[1+x^2]_{\mathcal{B}} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$

$$T\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}\right) = T([1+x^2]_{\mathcal{B}}) = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}.$$

and $[x+x^2]_{\mathcal{B}} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$

$$\Rightarrow T\left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right) = T([x+x^2]_{\mathcal{B}}) = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}.$$

\therefore The standard matrix of T is.

$$[T(\vec{e}_1) \quad T(\vec{e}_2) \quad T(\vec{e}_3)] = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

3. (10 points) Let A be a matrix which is row-equivalent to $\begin{bmatrix} 1 & 2 & 0 & 3 \\ 0 & 0 & 5 & 6 \\ 0 & 0 & 0 & 0 \end{bmatrix}$, and let $T: \mathbb{R}^3 \rightarrow \mathbb{R}^7$

be a linear transformation such that $T(\vec{x}) = \vec{0}$ if and only if $\vec{x} = A\vec{y}$ for some $\vec{y} \in \mathbb{R}^4$. Find the dimension of the range of T .

$$\text{Since } T(\vec{x}) = \vec{0} \iff \vec{x} = A\vec{y} \text{ for some } \vec{y} \in \mathbb{R}^4,$$

$$\text{Nul}(T) = \text{Col}(A).$$

$$\Rightarrow \dim \text{Nul}(T) = \dim \text{Col}(A).$$

since A RREF(A) has two pivots, $\dim \text{Col}(A) = 2$.

$$\Rightarrow \dim \text{Nul}(T) = 2.$$


Since,

$$\dim \text{Nul}(T) + \dim \text{Range}(T) = 3.$$

$$2 + \dim \text{Range}(T) = 3$$

$$\dim \text{Range}(T) = 1.$$

4. (a) (5 points) Let P be a parallelepiped in \mathbb{R}^3 . Suppose that one of its vertices is $(0, 1, 0)$, and the three vertices adjacent to that one are $(0, 1, 2)$, $(1, 1, 1)$ and $(1, 0, -1)$. Find the volume of P .



Volume will not be on the midterm.

- (b) (5 points) Let A and B be invertible matrices with $\det A = 4$ and $\det B = 2$. Find $\det(A^2 B^{-1})$.

$$\begin{aligned}\det(A^2 B^{-1}) &= \det(A^2) \det(B^{-1}) \\&= \det(A \cdot A) \frac{1}{\det(B)} \\&= \det(A) \det(A) \cdot \frac{1}{2} \\&= 4^2 \cdot \frac{1}{2} \\&= 8.\end{aligned}$$

Midterm 2vP

Math 18, Section A

May 19, 2017

Time Limit: 50 Minutes

Name (Print): _____

PID: _____

- **DO NOT** begin working, or even open this packet, until instructed to do so.
- You should be in your assigned seat, unless instructed otherwise by Ed or one of the TAs.
- Enter all requested information on the top of this page, and put your name on the top of every page, in case the pages become separated.
- You may use a two-sided page of notes on this exam.
- You may **not** use your books, additional notes, or any calculator on this exam.

You are required to show your work on each problem on this exam. The following rules apply:

- **Organize your work**, in a reasonably neat and coherent way, in the space provided. Work scattered all over the page without a clear ordering will receive very little credit.
- **Mysterious or unsupported answers will not receive full credit.** Unless otherwise directed in the statement of the problem, a correct answer, unsupported by calculations, explanation, or algebraic work will receive little or no credit. An incorrect answer supported by substantially correct calculations and explanations might still receive partial credit.

Question	Points
1	10
2	10
3	10
4	10
Total:	40

DO NOT turn this page until instructed to do so

1. (10 points) Let I be the 3×3 identity matrix and let

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix}$$

Find a matrix M satisfying $MA = B + I$.

Since $\det A = 1$, A is invertible.

$$\Rightarrow MA A^{-1} = (B + I) A^{-1}.$$

$$M = (B + I) A^{-1}.$$

\Rightarrow we need to find A^{-1} .

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & -2 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right] \quad \underbrace{\begin{bmatrix} 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}}_{A^{-1}}$$

$$\Rightarrow A^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$M = \left(\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & -1 & 3 \end{bmatrix}.$$

2. (10 points) Let $T: \mathbb{R}^{1903} \rightarrow \mathbb{R}^4$ be a linear transformation with range

$$\text{Range } T = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

Find $\dim \text{Nul } T$.

$$\dim \text{Nul}(T) + \dim \text{range}(T) = 1903.$$

We can find $\dim \text{range}(T)$ by

$$\begin{bmatrix} 1 & 0 & 1 & 1 \\ 1 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\uparrow \quad \uparrow$
 pivots.

$$\Rightarrow \dim \text{range}(T) = 2.$$

$$\therefore \dim \text{Nul}(T) = 1901.$$

3. (10 points) Let \mathbb{P} be the space of polynomials in x (with the usual addition and scalar multiplication) and let $H = \text{span}\{x, x^2, x^5\} \subset \mathbb{P}$. Then $\mathcal{B} = \{x, x^2, x^5\}$ is a basis for H , and $\mathcal{C} = \{x + x^5, x^2, x\}$ is also a basis for H . Find a matrix A satisfying the following condition:

- For any $p(x) \in H$,

$$A[p(x)]_{\mathcal{B}} = [p(x)]_{\mathcal{C}}$$

This question is "change of bases" (Section 4.7).
We have not learned it yet!

4. (10 points) Suppose that S is a region of \mathbb{R}^3 whose volume is 3. Let $A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 4 & 6 \\ 0 & 5 & 3 \end{bmatrix}$ and let $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the linear transformation $T(\vec{x}) = A^{2017}\vec{x}$. Find the volume of $T(S)$.

We will not cover volume for our midterm!