

Câu 1:

```
SELECT sinhvien_id, SUM(diem_anh) AS max_anh FROM tbl_diem GROUP BY  
sinhvien_id HAVING max_anh =
```

```
(  
    SELECT MAX(sum_anh)  
    FROM (  
        SELECT SUM(diem_anh) AS sum_anh  
        FROM tbl_diem  
        GROUP BY sinhvien_id  
    )  
    AS tbl_sum  
)
```

câu 2:

```
SELECT tbl_sinhvien.email, SUM(diem_van) AS min_van FROM tbl_diem,tbl_sinhvien  
WHERE tbl_sinhvien.id = tbl_diem.sinhvien_id GROUP BY sinhvien_id HAVING min_van =
```

```
(  
    SELECT MIN(sum_van)  
    FROM (  
        SELECT SUM(diem_van) AS sum_van  
        FROM tbl_diem  
        GROUP BY sinhvien_id  
    )  
    AS tbl_sum  
)
```

Câu 3:

```
SELECT tbl_lop.ten_lop,tbl_sinhvien.id, SUM(diem_anh) AS max_anh  
FROM tbl_diem,tbl_sinhvien,tbl_lop  
WHERE tbl_sinhvien.id = tbl_diem.sinhvien_id AND tbl_sinhvien.lop_id = tbl_lop.id  
GROUP BY sinhvien_id  
HAVING max_anh =
```

```
(  
    SELECT MAX(sum_anh)  
    FROM (  
        SELECT SUM(diem_anh) AS sum_anh  
        FROM tbl_diem  
        GROUP BY sinhvien_id  
    )  
    AS tbl_sum  
)
```