

1.1 Brief review of differentiation

The function $f : R \rightarrow R$ is differentiable at the point $x \in \mathbb{R}$ if the limit

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists, in which case the derivative $f'(x)$ is defined as

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

The function $f(x)$ is called differentiable if it is differentiable at all points x .

Theorem 1.1. (Product Rule.)

The product $f(x)g(x)$ of two differentiable functions $f(x)$ and $g(x)$ is differentiable, and

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x)$$

Theorem 1.2. (Quotient Rule.)

The quotient $\frac{f(x)}{g(x)}$ of two differentiable functions $f(x)$ and $g(x)$ is differentiable at every point x where the function $\frac{f(x)}{g(x)}$ is well defined, and

$$\left(\frac{f(x)}{g(x)} \right)' = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

Theorem 1.3. (Chain Rule.)

The composite function $(g \circ f) = g(f(x))$ of two differentiable functions $f(x)$ and $g(x)$ is differentiable at every point x where $g(f(x))$ is well defined and

$$(g(f(x)))' = g'(f(x))f'(x)$$

Example: Chain Rule is often used for power functions, exponential functions, and logarithmic functions.

$$\frac{d}{dx}((f(x))^n) = n(f(x))^{n-1}f'(x)$$

$$\frac{d}{dx}(e^{f(x)}) = e^{f(x)}f'(x)$$

$$\frac{d}{dx}(\ln(f(x))) = \frac{f'(x)}{\ln(f(x))}$$

Lemma 1.1.

Let $f : [a, b] \rightarrow [c, d]$ be a differentiable function, and assume that $f(x)$ has an inverse function denoted by $f^{-1}(x)$ with $f^{-1} : [c, d] \rightarrow [a, b]$. The function $f^{-1}(x)$ is differentiable at every point $x \in [c, d]$ where $f'(f^{-1}(x)) \neq 0$ and

$$(f^{-1}(x))' = \frac{1}{f'(f^{-1}(x))}$$

$$\left(\frac{1}{f(x)} \right)' = \frac{1}{f'(f^{-1}(x))}$$

Reuse the Chain Rule

$$(g(f(z)))' = g'(f(z))f'(z)$$

With $g = f^{-1}$, we have

$$(f^{-1}(f(z)))' = (f^{-1})'(f(z))f'(z)$$

$$z' = (f^{-1})'(f(z))f'(z)$$

$$1 = (f^{-1})'(f(z))f'(z)$$

Let $z = f^{-1}(x)$, then $f(z) = f(f^{-1}(x)) = x$ and

$$1 = (f^{-1})'(x)f'(f^{-1}(x))$$

$$\begin{aligned}\frac{1}{f'(f^{-1}(x))} &= (f^{-1})'(x) \\ &= (f^{-1}(x))'\end{aligned}$$

Examples:

$$\begin{aligned}\frac{d}{dx}(xe^{3x^2-1}) &= x'e^{3x^2-1} + x(e^{3x^2-1})' \\ &= e^{3x^2-1} + x(3x^2-1)'(e^{3x^2-1}) \\ &= e^{3x^2-1} + 6x^2(e^{3x^2-1}) \\ &= (1+6x^2)(e^{3x^2-1})\end{aligned}$$

$$\begin{aligned}\frac{d}{dx}\left(\frac{\sqrt{3x^2-1}}{\sqrt{3x^2-1}+4}\right) &= \frac{d}{dx}\left(\frac{\sqrt{3x^2-1}+4-4}{\sqrt{3x^2-1}+4}\right) \\ &= \frac{d}{dx}\left(1 - \frac{4}{\sqrt{3x^2-1}+4}\right) \\ &= \left(\frac{4}{\sqrt{3x^2-1}+4}\right)'\end{aligned}$$

—

$$\frac{d}{dx}\left(\frac{\sqrt{3x^2-1}}{\sqrt{3x^2-1}+4}\right) = \frac{d}{dx}\left(1 - \frac{4}{\sqrt{3x^2-1}+4}\right)$$

Apply Quotient Rule:

$$\frac{d}{dx}\left(\frac{f(x)}{g(x)}\right) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

With $f(x) = -4$, $g(x) = \sqrt{3x^2-1}+4$:

$$f'(x) = 0$$

$$\begin{aligned} g'(x) &= \left(\sqrt{3x^2 - 1}\right)' \\ &= \frac{1}{2} \cdot (3x^2 - 1)^{-\frac{1}{2}} \cdot 6x \\ &= \frac{3x}{\sqrt{3x^2 - 1}} \end{aligned}$$

$$(g(x))^2 = \left(\sqrt{3x^2 - 1} + 4\right)^2$$

$$\begin{aligned} \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2} &= \frac{0 \cdot \left(\sqrt{3x^2 - 1} + 4\right) + 4\left(3x \div \sqrt{3x^2 - 1}\right)}{\left(\sqrt{3x^2 - 1} + 4\right)^2} \\ &= \frac{\left(12x \div \sqrt{3x^2 - 1}\right)}{\left(\sqrt{3x^2 - 1} + 4\right)^2} \\ &= \frac{12x}{\sqrt{3x^2 - 1}\left(\sqrt{3x^2 - 1} + 4\right)^2} \end{aligned}$$

—

To calculate:

$$\frac{d}{dx} \left(\frac{e^{x^2} + 1}{x - 1} \right)$$

Apply Quotient Rule:

$$\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

With

$$\begin{aligned} f(x) &= e^{x^2} + 1 \\ f'(x) &= (x^2)' e^{x^2} \\ &= 2xe^{x^2} \\ g(x) &= x - 1 \\ g'(x) &= 1 \\ f'(x)g(x) - f(x)g'(x) &= 2xe^{x^2}(x - 1) - (e^{x^2} + 1) \\ (g(x))^2 &= (x - 1)^2 \end{aligned}$$

We have

$$\frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2} = \frac{2xe^{x^2}(x - 1) - (e^{x^2} + 1)}{(x - 1)^2}$$

—

$$\begin{aligned}
\frac{d}{dx} \left(\ln \left(\frac{x}{2x^2 + 1} \right) \right) &= \frac{d}{dx} (\ln(x) - \ln(2x^2 + 1)) \\
&= \ln(x)' - \ln(2x^2 + 1)' \\
&= \frac{1}{x} - \frac{(2x^2 + 1)'}{2x^2 + 1} \\
&= \frac{1}{x} - \frac{4x}{2x^2 + 1} \\
&= \frac{2x^2 + 1}{x(2x^2 + 1)} - \frac{4x^2}{x(2x^2 + 1)} \\
&= \frac{-2x^2 + 1}{x(2x^2 + 1)}
\end{aligned}$$