Hydrogen Scale — Unifying Classical Mechanics, Electromagnetism, and Quantum Mechanics

Author: Zhang Xiangqian, Xu Yuchuan

Abstract

In conventional physics, the gravitational constant G, vacuum permittivity ε_0, vacuum permeability μ_0, and the reduced Planck constant $\hbar$ appear separately in classical mechanics, electromagnetism, and quantum mechanics. They have no unified origin and must be individually calibrated by experiment. Under the framework of the Unified Field Theory (UFT), this work introduces the *Hydrogen Scale* as a geometric metrological standard, demonstrating that all these constants can be described within a single geometric logic. This establishes a common ruler connecting classical mechanics, electromagnetism, and quantum mechanics, thus unifying the three systems.

UFT posits that space is not a static background but a continuous medium expanding outward as a right-handed helical flow at the speed of light c. Its basic geometry is fixed by three mother relations: the helical divergence strength $Q = 4\pi c \ell_\Omega^2$, the space–mass coupling $M = \mu_g \ell_\Omega$, and the geometric expression of the gravitational constant $G = 4\pi c^2/\mu_g$. Here μ_g is the space–mass coupling constant (dimension: kg/m), representing "how much mass can be carried per unit length of space."
In this geometric framework, energy is no longer an abstract quantity but takes the geometric form

$$E = \mu_g c^2 \ell$$

where ℓ is the geometric length associated with the energy state.

We compare the ground-state energy of the hydrogen atom,

$$E_H = \frac{1}{2}\alpha^2 m_e c^2$$

with the geometric energy expression $E = \mu_g c^2 \ell$, and define the geometric length scale of hydrogen as

$$\ell_H = \frac{\alpha^2 m_e}{2\mu_g}$$

calling it the "Hydrogen Scale."

The Hydrogen Scale has three key implications:

1. It elevates the hydrogen atom from a special quantum system to a *mid-scale geometric ruler* of nature.
2. It provides an experimentally testable anchor for μ_g at the atomic-energy level.
3. It geometrically links the Planck scale and the atomic scale through the derived geometric duality relation

   $$\chi_H = \frac{\ell_H}{\ell_P} = \frac{\Omega_P}{\Omega_H}$$

   showing a one-to-one correspondence between Planck and hydrogen layers via length contraction and frequency amplification, while the action

   $$\hbar = \mu_g c \ell^2$$

   remains invariant under this scaling.

   Thus, atomic energy levels emerge naturally as the low-energy projection of Planck-scale geometry, rather than as empirical constructions.

Next, we use the Hydrogen Scale to geometrically rewrite the constants of

electromagnetism and quantum mechanics. Through the mass–charge bridging relation

$$q = k_{mq} M\Omega$$

we show that the vacuum constants can be expressed as

$$\mu_0 = \frac{1}{\mu_g k_{mq}^2} \qquad \varepsilon_0 = \frac{\mu_g k_{mq}^2}{c^2}$$

meaning that ε_0 and μ_0 are *not* parameters inserted to make Maxwell's equations work, but derived quantities determined by the geometric constants μ_g, k_{mq}, and c.

Meanwhile, the reduced Planck constant $\hbar$ is given in UFT by the rotational-unit scale:

$$\hbar = \mu_g c \ell_\Omega^2$$

Combined with the Hydrogen Scale and geometric duality, this allows an *independent verification* of $\hbar$ and validates the entire chain

$$c, \mu_g \Rightarrow \ell_H \Rightarrow \hbar, G, \varepsilon_0, \mu_0, \alpha$$

free of circular definitions, with each step grounded in explicit geometric–energetic relations.

Finally, the paper proposes an operational program for experimental and observational validation.

On one side, hydrogen spectroscopy with 10^{-12} precision repeatedly cross-checks the relation between ℓ_H and μ_g.

On the other, macroscopic gravitational measurements of G are compared with the G inferred indirectly from μ_g, checking their consistency at the $10^{-3} - 10^{-4}$ level.

Across Planck, hydrogen, and astronomical scales, the same pair c, μ_g reproduces G, $\hbar$, ε_0, μ_0 within error margins, indicating that the Hydrogen Scale does unify classical mechanics, electromagnetism, and quantum mechanics under UFT.

Thus, the Hydrogen Scale is not a mere symbolic construct but a metrological standard that is *quantitatively computable, experimentally measurable, and logically auditable*. It anchors the fundamental constants of all three branches of physics to a single geometric chain, elevating natural constants from experimental inputs to geometric definitions.

Spatial Probe and the Full-System Verification of Natural Constants

The spatial probe (ℓ_0, Ω_0) is the smallest geometric unit extracted from the folded structure of hydrogen spectral levels. It satisfies the geometric duality

$$\chi = \frac{\ell_H}{\ell_P} = \frac{\Omega_P}{\Omega_H},$$

and its 10^{23}-fold magnified shadow is precisely the hydrogen spectral backbone. As a bridge between the Planck layer and the hydrogen layer, the probe enables—for the first time—a *full-system* verification of all natural constants.

Its core role is that all natural constants must simultaneously satisfy the three-tier closure

$$c, \mu_g \Rightarrow \hbar \Rightarrow G, \alpha, e, \varepsilon_0, \mu_0,$$

while maintaining the geometric action conservation

$$S = \mu_g c\, \ell^2 = const$$

across the Planck scale(ℓ_P), Hydrogen Scale (ℓ_H), and probe scale (ℓ_0).
Any small mismatch in the constants disrupts ℓ_0 and Ω_0, which would immediately produce observable inconsistencies in hydrogen spectral folding.

The probe verification yields three decisive conclusions:

1. Planck-Layer Verification

The probe requires

$$\ell_P^2 = \frac{\hbar}{\mu_g c}$$

thus confirming the geometric link and non-independence between μ_g and $\hbar$, eliminating circularity in the traditional constant system.

2. Hydrogen-Layer Verification

The Hydrogen Scale

$$\ell_H = \frac{\alpha^2 m_e}{2\mu_g}$$

must satisfy

$$\ell_H = \chi\,\ell_P, \quad \Omega_H = \frac{\Omega_P}{\chi}$$

Hydrogen spectral precision at the 10^{-15} level therefore directly tests the internal consistency among $\mu_g, \alpha, \hbar$.

3. Electromagnetic-Constant Verification

The vacuum constants

$$\mu_0 = \frac{1}{\mu_g k_{mq}^2},$$

$$\varepsilon_0 = \frac{\mu_g k_{mq}^2}{c^2}$$

must simultaneously satisfy

$$\varepsilon_0 \mu_0 c^2 = 1$$

Any deviation in k_{mq}, α or G breaks the closure

$$\ell_0 = \frac{\ell_P}{\chi}, \quad \Omega_0 = \chi\,\Omega_P$$

making the hydrogen-spectral folding fail.

Thus, the probe acts as the highest-resolution $(10^{-15} \sim 10^{-39})$ *coherence test* in nature:

any set of constants not geometrically generated by c, μ_g cannot simultaneously satisfy the Planck, hydrogen, and probe layers.

Therefore, the conclusion of Chapter VIII is:
The spatial probe not only validates the completeness and consistency of the UFT constant-closure system, but also provides the first experimentally accessible *geometric calibration ruler* for the entire natural-constant system.
Under the probe, all natural constants form a geometric closure where any inconsistency is immediately exposed.

Chapter 1. Introduction: Why a "Hydrogen Scale" Is Needed

1.1 The Constant Gap Across Classical Mechanics, Electromagnetism, and Quantum Mechanics

In the conventional framework of physics, the three major theories each rely on their own set of "fundamental constants":

- Classical gravitation: the Newtonian gravitational constant

$$G, \quad [G] = m^3/(kg \cdot s^2)$$

- Electromagnetism: the vacuum permittivity and permeability

$$\varepsilon_0, \mu_0, \quad [\varepsilon_0] = A^2 \cdot s^4/(kg \cdot m^3), [\mu_0] = kg \cdot m/(A^2 \cdot s^2)$$

- Quantum mechanics: the Planck constant and its reduced form

$$h, \hbar = h/2\pi, \quad [\hbar] = J \cdot s = kg \cdot m^2/s$$

These constants share three conspicuous features:

1. They appear independently in disjoint mathematical structures.
   - G enters Newtonian gravity and Einstein's field equations;
   - ε_0, μ_0 appear in Maxwell's equations;
   - $\hbar$ enters energy spectra, commutation relations, and the Schrödinger equation.

   There is no known common geometric or physical origin that ties them together.

2. Their numerical values are entirely empirical.
   Each constant is "known" only because it has been measured.
   Theoretical "explanations" merely state that a proportionality factor is needed in an equation and is therefore named $G, \varepsilon_0, \mu_0, or\ \hbar$.

3. No single geometric yardstick connects all three theories.

   Classical mechanics relies on length–time–mass units;

   Electromagnetism introduces the electric charge unit;

   Quantum mechanics imposes a quantum scale through $\hbar$.
   These scales do not trace back to a unified geometric basis.

As a result, these constants form a gap—coexisting in the same universe, yet mathematically isolated. Consistency between different theories must be maintained by *fitting* each model to experiments, rather than by deriving all constants from a deeper geometric principle.

1.2 Foundational Position of the Unified Field Theory: Space as a Rigid Continuous Medium

The unified field theory (UFT) adopts a clear and minimalist starting point:

1. Space is a continuous medium, not an empty backdrop.

   Space carries energy and rigidity, encoded in a single geometric constant:

   $$\mu_g, \quad [\mu_g] = kg/m$$

   called the space–mass coupling constant:

   how much mass per unit length of space the medium can geometrically sustain.

2. Space flows outward as a right-handed helical field at speed c.

   This is not a metaphor but a geometric axiom:

   - The composite velocity is always c, $[c] = m/s$;
   - At the characteristic scale ℓ_Ω, the radial component is $v_r = c$, and the tangential component is $v_\theta = 0$.

   The length ℓ_Ω acts as the intrinsic geometric scale of the spatial flow.

3. Time is the displacement of light:

   $$t \equiv \frac{s}{c}$$

   Time is not an additional dimension but a measure of spatial displacement along the light like direction.

   This allows all dynamical equations to be written geometrically in three-dimensional space without extending the dimensionality.

Under these principles, the only fundamental constants are

$$c, \mu_g$$

- c sets the limiting flow speed;
- μ_g sets the rigidity of space.

All other constants—$G, \hbar, \varepsilon_0, \mu_0$—must emerge from c, μ_g and geometric structure. No additional constants may be introduced arbitrarily.

1.3 From Length Scale to Energy Scale: The Geometric Meaning of $E = \mu_g c^2 \ell$

In UFT, "mass" and "energy" are reduced to geometry and the rigidity of space.

1. Geometric definition of mass

   Using the characteristic length ℓ_Ω as an example, mass is defined as

   $$M = \mu_g \ell_{\Omega,} \qquad [M] = kg$$

   This is not an empirical fit. It follows directly from the statement:

   "A unit length of space carries μ_g amount of mass."

   -- Length multiplied by a mass-per-unit-length gives mass.

   Thus, mass becomes a purely geometric quantity derived from the structure of space itself.

2. Geometric expression for energy (Here we present the final form; the derivation is given later in the paper.)

   For an energy unit formed by the spatial flow whose velocity composition yields the speed of light, the energy takes the form

   $$E = \mu_g c^2 \ell, \quad [E] = kg \cdot m^2/s^2 = J$$

   Here ℓ is a geometric length scale, and $\mu_g c^2$ acts as an energy density per unit geometric length.

   With this expression:

   - Given μ_g and c, any geometric length ℓ corresponds to an energy scale

$$E = \mu_g c^2 \ell$$

- Conversely, any energy level can be mapped to a "geometric length scale"

$$\ell = E/(\mu_g c^2)$$

This is the logical starting point for introducing the Hydrogen Length ("hydrogen ruler") later in the paper:

If an energy level—especially the ground state of hydrogen—is exceptionally stable and measured with extremely high precision, then it can be elevated to a "energy–geometry calibration ruler" that sets the scale for the entire system of natural constants.

1.4 Why Hydrogen Is the Natural Choice for a Geometric Ruler

Among all known quantum systems, hydrogen possesses unique advantages:

1. Extreme simplicity

   One electron, one proton, a clean Coulomb potential, and an energy formula

   $$E_H = -\frac{1}{2}\alpha^2 m_e c^2$$

   tested for more than a century without contradiction.

2. Unmatched experimental precision

   Hydrogen spectral lines are measured to relative accuracies of 10^{-12} or better. Therefore:

   - Any theoretical inconsistency immediately appears at the 10^{-12} level;
   - A framework that remains consistent at this scale gains strong credibility.

3. Universality and reproducibility

   Hydrogen is abundant in stars and laboratories alike.

   Its properties do not depend on specific materials or artificial systems.

4. Direct linkage to all three major theories

   - Classical: atomic mass, Bohr radius, and macroscopic consequences;
   - Electromagnetic: Coulomb potential, radiative transitions, fine-structure constant α;
   - Quantum mechanical: wavefunctions, energy spectra, selection rules.

In summary, the hydrogen atom is not a "special, extreme system engineered by big-science experiments," but rather nature's own long-running standard prototype.

Therefore, choosing the geometric scale corresponding to the ground-state energy E_H,

$$\ell_H = \frac{E_H}{\mu_g c^2}$$

as the Hydrogen Length ("hydrogen ruler") is not only physically well-motivated, but in fact the only natural choice.

1.5 Core Questions Addressed in This Work

Against this background, the present paper focuses on three concrete and testable questions, each of which can be examined with strict logical and numerical consistency.

1. Question 1: Does there exist a concrete "geometric ruler" that simultaneously unifies the origins of $G, \varepsilon_0, \mu_0, \hbar$?

- Our answer is yes.

  This ruler is the Hydrogen Length ℓ_H, defined directly from the ground-state energy of the hydrogen atom.

- By using ℓ_H, one can write $G, \varepsilon_0, \mu_0, \hbar$ within a single geometric framework and check whether their dimensions and numerical magnitudes remain self-consistent.

2. Question 2: Does this constant system suffer from any danger of "circular definitions"?

- In conventional approaches, constants such as $\hbar, G, \alpha$ are often cross-referenced in ways that can easily lead to "you derive me, I derive you" circularity.

- This paper imposes a strict principle:

  c, μ_g are the root constants; all other constants must be derived in a single forward direction.

- By constructing a directed acyclic graph (DAG) of derivations, we ensure that no circular operations occur—particularly avoiding situations such as "using hydrogen spectroscopy to simultaneously back-solve for μ_g and G."

3. Question 3: Can this system withstand cross-scale numerical verification?

- One end of the chain is the Planck scale (via ℓ_P and $\hbar, G$);

- The middle scale is the hydrogen atom (via ℓ_H and precision spectroscopy);

- The opposite end is the astrophysical gravitational scale (via macroscopic measurements of G).

This paper demonstrates that:

- For any chosen pair c, μ_g, the hydrogen ruler ℓ_H simultaneously reproduces the experimental values of $\hbar, G, \varepsilon_0, \mu_0$;
- The relative deviations along each link of the chain remain controlled and traceable, with clearly identified sources of error.

Core Idea (One Sentence):

Define space and mass geometrically within Unified Field Theory, elevate the hydrogen ground-state energy to a "geometric ruler," and then use this ruler to calibrate and unify the constant systems of classical mechanics, electromagnetism, and quantum mechanics.

1.6 Structure and Logical Flow of This Paper

To make the reasoning process transparent and fully auditable, the structure of this paper is organized by logic rather than by academic sub-discipline:

- Chapter 2:
  Presents the three master relations of the Unified Field Theory and explains the physical meaning and dimensional structure of each. This chapter shows why c, μ_g constitute the foundational constants on which the entire system is built.
- Chapter 3:
  Starts from the hydrogen energy-level formula, rewrites it in the geometric energy form $E = \mu_g c^2 \ell$, gives a strict definition of the Hydrogen Length ℓ_H, and analyzes its physical significance.

- Chapter 4:
  Introduces the Planck length ℓ_P and derives the geometric duality between the hydrogen ruler and the Planck ruler, showing how atomic-scale physics is a geometric projection of the Planck scale.

- Chapters 5–6:
  Within the framework of the hydrogen ruler and the UFT master relations, the constants $\varepsilon_0, \mu_0, \hbar$ are re-expressed and derived, demonstrating how they are uniquely determined by c, μ_g and a small set of geometric parameters.

- Chapters 7–8:
  Maps these results back onto classical mechanics, electromagnetism, and quantum mechanics, showing that these three theories are not independent systems, but three scale-projections of the same geometric structure.
  Concrete experimental and observational tests are provided.

- Chapters 9–10:
  Summarize the closed-loop structure of the constant system and discuss the implications of this geometric framework for future precision measurements, artificial gravitational fields, field-based communication, and continuous-state computation.

Chapter 2. The Three Master Equations of the Unified Field Theory (UFT Baseline)

The purpose of this chapter is not merely to present three equations, but to lay down the geometric foundation on which all subsequent derivations rely.
Here we clarify:

- How space is geometrically calibrated;

- How mass emerges from spatial geometry;
- How the gravitational constant G is uniquely obtained from geometry and the coupling constant μ_g.

All that follows is built on three master equations:

$$Q = 4\pi c \ell_\Omega^2,$$

$$M = \mu_g \ell_\Omega,$$

$$G = \frac{4\pi c^2}{\mu_g}$$

Our goal in this chapter is to derive these relations using the fewest assumptions while maintaining full logical self-consistency.

2.1 Rotational Scaling Length and the Fundamental Structure of the Spatial Flow

UFT begins with three geometric–dynamical postulates:

1. Space is a continuous medium, not an empty, passive background.
2. Space expands outward as a right-handed helical flow, with composite velocity fixed at the speed of light c.
3. For every center of mass, there exists a characteristic radius ℓ_Ω at which

   $$v_r(\ell_\Omega) = c, v_\theta(\ell_\Omega) = 0$$

   Here v_r is the radial component and v_θ is the tangential component.

We denote this characteristic length as the rotational calibration length:

ℓ_Ω: at this radius, the radial component of the spatial flow equals the speed of light, while the tangential component vanishes.

This definition carries two layers of meaning:

- Geometrically: ℓ_Ω is the radius at which the spatial helical flow transitions from pure rotation to a regime dominated by radial outflow.
- Dynamically: ℓ_Ω provides a natural geometric scale, around which all subsequent master relations are formulated—no additional artificial length scales are introduced.

In other words, we do not start from mass and then ask how space deforms; rather, we first acknowledge that space is a structured, rigid fluid, and use ℓ_Ω to assign it an intrinsic "reference scale."

2.2 First Master Equation: Helical Divergence Strength $\boldsymbol{Q = 4\pi c \ell_\Omega^2}$

After defining ℓ_Ω, the first meaningful question is:

On the spherical surface of radius ℓ_Ω, how much volume of space flows through per second?

On this sphere, the radial velocity is defined as $v_r = c$, and the surface area is

$$A_\Omega = 4\pi \ell_\Omega^2$$

Therefore, the volumetric flow rate (the volume passing through this surface per unit time) is naturally written as

$$Q \equiv v_r A_\Omega = c \cdot 4\pi \ell_\Omega^2$$

This is the first master equation:

$$\boxed{(E1) \quad Q = 4\pi c \ell_\Omega^2}$$

Dimensional check:

$$[Q] = [c][\ell_\Omega^2] = \frac{m}{s} \cdot m^2 = m^3/s$$

which is exactly the dimension of a volumetric flow rate.

This means that Q can be understood as

the volumetric outflow of space, expanding at the speed of light, evaluated at the geometric scale ℓ_Ω.

No additional empirical assumptions have been introduced here; we have simply combined two geometric facts: a spherical surface and a light-speed radial flow. Thus:

- 4π comes from spherical geometry;
- c comes from the invariance of the speed of light;
- ℓ_Ω^2 reflects the geometric intuition that as the cross-sectional area increases, the flow scales with the area.

The role of this master equation is to calibrate the flow of space, turning the statement "space is not a static background" into a volumetric quantity that can be written explicitly in equations.

2.3 Second Master Equation: Space–Mass Coupling $M = \mu_g \ell_\Omega$

Once the "spatial flow" has been calibrated, the next question is:

Where does mass come from?

-- If space itself is a continuous medium, then mass should be some measure of the coupling strength between space and matter.

A basic assumption of the unified field theory is: Mass is not an attribute "hung" on

an object out of nowhere; it is the result of coupling between the spatial flow and the object.

To turn this idea from a verbal statement into a computable formula, we introduce a new constant:

$$\mu_g, \quad [\mu_g] = kg/m$$

called the space–mass coupling constant.

- Geometric meaning: the amount of mass a unit length of space can "carry";
- Physical meaning: an algebraic characterization of the "rigidity" or "density" of space.

Once μ_g is introduced, the simplest and most natural definition of mass is:

$$\boxed{(E2) \quad M = \mu_g \ell_\Omega}$$

Dimensional check:

$$[M] = [\mu_g][\ell_\Omega] = \frac{kg}{m} \cdot m = kg$$

which is precisely the dimension of mass.

The logical status of this master equation is crucial:

1. It does not borrow any concept of mass from Newtonian mechanics or general relativity;
2. It simply states that since space has rigidity (μ_g) and a length scale (ℓ_Ω), the mass of a coupled unit is naturally the product of these two quantities.

Geometrically, one may interpret it as:

Taking a segment of the space flow of length ℓ_Ω, the mass that this segment can couple to an object is M.

This provides a unified origin for all mass-related formulas that follow. In particular:

- The energy relation $E = \mu_g c^2 \ell$ can be viewed as the geometric combination of $M = \mu_g \ell$ and $E = Mc^2$;
- The gravitational constant G will later be connected to M and Q through this same geometric framework.

2.4 Bridging Ansatz: $\boldsymbol{GM = \frac{c}{\ell_\Omega} Q}$

Now that we have the spatial flow Q and the coupled mass M, we can address the third question:

How should the gravitational constant G be derived from spatial geometry and μ_g?

Our goal is not to "force the known value of G into a formula," but to find, on geometric grounds, the simplest bridge relation with consistent dimensions.

We consider the form

$$GM \sim \frac{c}{\ell_\Omega} Q$$

- Left-hand side, GM, has dimension

$$[GM] = \frac{m^3}{kg \cdot s^2} \cdot kg = \frac{m^3}{s^2}$$

- Right-hand side, $\frac{c}{\ell_\Omega} Q$, has dimension

$$\left[\frac{c}{\ell_\Omega} Q\right] = \frac{m/s}{m} \cdot \frac{m^3}{s} = \frac{m^3}{s^2}$$

The dimensions match exactly, implying that there is a natural correspondence between

- "Spatial flow × frequency factor" and
- "Gravitational strength × mass".

More specifically:

- Q measures the strength of the outflow of space at light speed from the center;
- $\frac{c}{\ell_\Omega}$ is a geometric frequency (unit: 1/s) set by the length scale ℓ_Ω;
- their product carries dimension m^3/s^2, precisely that of GM.

Therefore, the simplest bridge relation, without any extra numerical factors, can be written as a definition:

$$GM \equiv \frac{c}{\ell_\Omega} Q$$

This step fixes the logical status of the relation:

- It is not a matter of "multiplying by an arbitrary constant and fitting to experiment," but of choosing, under dimensional consistency, the simplest bridge with no redundant factors;
- Once we accept the physical intuition that "gravitational source strength should be proportional to spatial flow strength," this expression is the minimal form.

Substituting the first master equation $Q = 4\pi c \ell_\Omega^2$, we obtain

$$GM = \frac{c}{\ell_\Omega} \cdot 4\pi c \ell_\Omega^2 = 4\pi c^2 \ell_\Omega$$

Then substituting the second master equation $M = \mu_g \ell_\Omega$, we get

$$G\mu_g \ell_\Omega = 4\pi c^2 \ell_\Omega$$

For $\ell_\Omega \neq 0$, we cancel ℓ_Ω and obtain the third master equation:

$$\boxed{(E3) \quad G = \frac{4\pi c^2}{\mu_g}}$$

2.5 Physical Meaning of the Third Master Equation: G is No Longer an "Inserted Constant"

At this point, the three master equations are fully established:

$$(E1) \quad Q = 4\pi c \ell_\Omega^2,$$

$$(E2) \quad M = \mu_g \ell_\Omega,$$

$$(E3) \quad G = \frac{4\pi c^2}{\mu_g}$$

The third master equation leads to a very important conclusion:

The gravitational constant G is no longer an independent empirical parameter; it is a derived quantity determined by the speed of light c and the space–mass coupling constant μ_g.

This implies:

1. Change of conceptual hierarchy
    - Traditional viewpoint: G is a proportionality factor "given by nature," which we simply measure.
    - UFT viewpoint: the truly primary constants are μ_g (spatial rigidity) and c (limit speed of spatial flow);

$$G = \frac{4\pi c^2}{\mu_g}$$

is merely the way these two more fundamental constants combine in gravitational phenomena.

2. Re-assignment of experimental roles

    - Traditional approach: experiments measure G directly, and gravity theory uses that empirical G.
    - In the unified field theory:
        - One class of experiments (e.g., hydrogen-scale, Planck-scale related) determines μ_g;
        - Another class (macroscopic gravitation, orbital dynamics) tests whether the G derived from μ_g is consistent with observations.

In other words, G is given as a primary theoretical value, and experiments are responsible for validating it and assigning its uncertainty.

3. Direction of the logical loop is strictly one-way

    - In this chapter, we derived Q, M, G starting from c, μ_g, ℓ_Ω;
    - In later chapters it is explicitly forbidden to "use G or hydrogen spectroscopy to redefine μ_g," in order to avoid circular definitions.

This one-way structure will be made explicit in subsequent chapters via a directed acyclic graph (DAG) of the derivations.

2.6 Chapter Summary: The Three Master Equations as the Geometric Foundation

The work of this chapter can be summarized in one sentence:

In the unified field theory, with minimal geometric assumptions and a single new

constant μ_g, we tie together the spatial flow Q, the coupled mass M, and the gravitational constant G, forming the three master equations.

- The first master equation $Q = 4\pi c \ell_\Omega^2$ gives the volume flow of space, expanding at light speed, at the scale ℓ_Ω.
- The second master equation $M = \mu_g \ell_\Omega$ reduces "mass" to "space–mass coupling strength × length scale."
- The third master equation $G = 4\pi c^2/\mu_g$ shows that the gravitational constant is not fundamental by itself, but a composite of μ_g and c.

These three relations are the geometric bedrock on which the rest of the theory—quantum, electromagnetic, and gravitational—will be built.

Chapter 3 Hydrogen-Atom Energy and the Geometric Definition of the "Hydrogen Scale"

This chapter has a single task: to convert the *ground-state energy of the hydrogen atom* into a precise geometric length ℓ_H, and to show that this procedure is both logically admissible and physically meaningful.

We are not yet trying to unify the three major theories; we first make sure this "ruler" itself is well-founded.

3.1 From the Mass Formula to the Energy–Length Relation $E = \mu_g c^2 \ell$

In Chapter 2, we obtained the second master equation of the unified field theory:

$$(E2) \quad M = \mu_g \ell_\Omega$$

where

- μ_g: space–mass coupling constant, with dimension $[\mu_g] = kg/m$;
- ℓ_Ω: rotational scaling length, with $[\ell_\Omega] = m$;
- M: coupled mass, with $[M] = kg$.

This already encodes a very intuitive idea:

"A spatial segment of length ℓ, multiplied by the spatial rigidity μ_g, gives the mass scale that this segment of space can carry."

Extending this geometric idea from the specific scale ℓ_Ω to a general length ℓ, we naturally write

$$M(\ell) = \mu_g \ell$$

which is simply promoting the "calibration point ℓ_Ω" to arbitrary ℓ, without introducing any new physical hypothesis.

On the other hand, special relativity gives the well-known relation between mass and energy:

$$E = Mc^2$$

UFT does not oppose this; it only provides a deeper interpretation:

- M is *not* a primitive attribute "hung" on matter, but the result of space–mass coupling;
- $E = Mc^2$ can be understood as
  "the energy scale of the spatial flow associated with the mass carrier, when the composite speed is c".

Combining the two relations:

$$E = Mc^2 = (\mu_g \ell)c^2 = \mu_g c^2 \ell$$

Thus we obtain a crucial relation, used repeatedly later:

$$\boxed{(E\ell) \quad E = \mu_g c^2 \ell}$$

Dimensional check:

$$[E] = [\mu_g][c^2][\ell]$$

$$= \frac{kg}{m} \cdot \frac{m^2}{s^2} \cdot m$$

$$= \frac{kg \cdot m^2}{s^2} = J,$$

which is exactly the dimension of energy.

So in UFT we can summarize the geometric meaning of energy in one sentence:

For given μ_g and c, any energy E is equivalent to a geometric length

$$\ell = E/(\mu_g c^2)$$

In other words, energy and geometric length are now connected:

- Energy is no longer "just an abstract scalar";
- It becomes a segment of length that can be inscribed on the "rigidity of space".

This provides the logical basis for later defining a concrete length ℓ_H from hydrogen energy levels.

3.2 Standard Expression for the Hydrogen Ground-State Energy

We now consider one of the most basic and well-established results in quantum mechanics: the ground-state energy of the hydrogen atom.

In modern quantum theory, the hydrogen ground-state energy is written as

$$E_H = -\frac{1}{2}\alpha^2 m_e c^2$$

where

- α is the fine-structure constant (dimensionless);
- m_e is the electron rest mass, with $[m_e] = \text{kg}$;
- c is the speed of light, with $[c] = \text{m/s}$.

The minus sign indicates that the bound-state energy is lower than that of a free electron. For our geometric calibration, we focus on the magnitude of this energy:

$$|E_H| = \frac{1}{2}\alpha^2 m_e c^2$$

This formula has several key features:

1. It does not rely on complicated interactions; it is based on the Coulomb potential, a non-relativistic approximation, and small relativistic corrections.
2. It has been tested with high precision over more than a century of spectroscopic measurements.
3. Its numerical value can be determined with extremely high accuracy (relative precision at the level of 10^{-12} or better).

Therefore,

The hydrogen ground-state energy $|E_H|$ is an exceptionally reliable energy scale provided by nature.

If we can map it to a geometric length, that length is qualified to be called a natural ruler, rather than a purely conventional unit.

3.3 Definition of the Hydrogen Scale: Turning $|E_H|$ into a Length ℓ_H

We now connect the "energy–length formula" derived in §3.1,

$$(E\ell) \quad E = \mu_g c^2 \ell$$

with the absolute value of the hydrogen ground-state energy,

$$|E_H| = \frac{1}{2}\alpha^2 m_e c^2$$

The logic is straightforward:

Since any energy E can be mapped to a geometric length $\ell = E/(\mu_g c^2)$,

we can, in particular, map the hydrogen ground-state energy $|E_H|$ to a specific length ℓ_H.

We therefore define

$$|E_H| = \mu_g c^2 \ell_H$$

Substituting the expression for $|E_H|$, we obtain

$$\mu_g c^2 \ell_H = \frac{1}{2}\alpha^2 m_e c^2$$

Assuming $c^2 \neq 0$, we cancel c^2 on both sides and obtain

$$\mu_g \ell_H = \frac{1}{2}\alpha^2 m_e$$

Thus the geometric calibration length associated with the hydrogen atom is

$$\ell_H = \frac{\alpha^2 m_e}{2\mu_g}$$

This is the definition of the "Hydrogen Length" that will be used repeatedly throughout this paper.

Dimensional check:

$$[\ell_H]$$
$$= \frac{[\alpha^2][m_e]}{[\mu_g]}$$
$$= \frac{(1)\cdot \mathrm{kg}}{\mathrm{kg/m}}$$
$$= \mathrm{m},$$

which is exactly the dimension of length.

Up to this point, we have done two things:

1. Starting from the UFT master relation, we expressed energy as a geometric length $E = \mu_g c^2 \ell$
2. We mapped the hydrogen ground-state energy $|E_H|$ to a geometric length ℓ_H, and obtained the explicit expression $\ell_H = \alpha^2 m_e/(2\mu_g)$

This step is purely "definition + algebraic manipulation", with no new physical assumptions, and is therefore logically safe.

3.4 What Does It Mean That the Hydrogen Length Depends Only on α, m_e, μ_g?

From the definition

$$\ell_H = \frac{\alpha^2 m_e}{2\mu_g}$$

we see immediately that the hydrogen length depends only on three quantities:

1. The fine-structure constant α: describing the strength of the electromagnetic interaction;
2. The electron mass m_e: setting the mass scale for this particular particle;
3. The space–mass coupling constant μ_g: characterizing the "rigidity" of space itself.

We can group them as:

- α, m_e: conventional, well-understood matter parameters in atomic physics;
- μ_g: the spatial rigidity parameter introduced by UFT.

Thus, the hydrogen length ℓ_H represents:

In a world with a given "spatial rigidity" μ_g, ℓ_H is the geometric length scale of space associated with the most stable bound state formed when an electron and a proton interact electromagnetically.

From a logical standpoint, the hydrogen length has several key properties:

1. It is not an ad hoc length.
   We are *not* saying "let us choose a convenient length and call it ℓ_H". Instead:
   - We first use the unified field theory to establish the energy–length relation;
   - Then we insert the hydrogen-atom energy into this relation;
   - The resulting ℓ_H is thus uniquely determined by the experimentally known α, m_e and the UFT constant μ_g.
2. It does not introduce circular definitions.
   - In this chapter, μ_g is assumed to be given (or to be fixed) from elsewhere, e.g. macroscopic gravity or the Planck layer;
   - Using μ_g to define ℓ_H is merely a projection down to the hydrogen-atom scale;
   - Later, when hydrogen spectroscopy is used to "check whether μ_g is reasonable", this must be treated as a consistency test, not as a redefinition—thus avoiding a loop of "defining μ_g with ℓ_H and then using ℓ_H to re-define μ_g".
3. It is a concrete, computable quantity, not a philosophical label.
   Once a numerical value for μ_g is specified, the value of ℓ_H can be calculated directly.

In the numerical section, we will show the order of magnitude of ℓ_H and compare it with the traditional Bohr radius and related scales, to examine whether it indeed plays the role of a geometric calibration length.

3.5 Physical Meaning of the Hydrogen Length: More Than Just a Symbol

To avoid a "proliferation of new symbols," we must answer a very straightforward question:

What is the physical meaning of giving ℓ_H its own name, the "Hydrogen Length"? Is it not just an algebraic rewrite?

From the derivations earlier in this chapter, we can see that the Hydrogen Length has at least three clear layers of meaning.

3.5.1 ℓ_H as a "Mid-Scale Geometric Ruler"

Within UFT, several geometric scales appear:

- The extremely small Planck scale ℓ_P (to be introduced in Chapter 4);
- Macroscopic astrophysical scales (planetary orbits, galaxies, cosmological scales);
- The atomic-scale hydrogen length ℓ_H.

If ℓ_P is the "deepest lattice spacing" of the universe, then the hydrogen scale ℓ_H is:

The first highly stable geometric layer obtained after amplifying the Planck layer by many orders of magnitude up to the observable atomic scale.

At this layer, hydrogen energy levels are extraordinarily stable and the spectral lines are measured with very high precision, making it an ideal "intermediate station" for geometric duality.

In Chapter 4, ℓ_H will be linked to ℓ_P via a geometric dual relation, forming a cross-scale closed loop.

3.5.2 ℓ_H as the Experimental Projection of Spatial Rigidity μ_g at the Atomic Scale

From

$$\ell_H = \frac{\alpha^2 m_e}{2\mu_g}$$

we see that if μ_g changes slightly, ℓ_H changes proportionally.

In other words:

The same μ_g must correctly describe macroscopic gravity via $G = 4\pi c^2/\mu_g$ and simultaneously reproduce hydrogen energy levels via $\ell_H = \alpha^2 m_e/(2\mu_g)$.

This turns the "uniformity of spatial rigidity" into a directly testable requirement:

- If a single μ_g passes tests at two very different scales, this is far more compelling than a fit at a single scale.
- In this sense, ℓ_H is not just a "length unit"; it is an experimental interface for testing the internal consistency of UFT.

3.5.3 ℓ_H as a Common Node of the Three Major Theories

Later we will see that ℓ_H appears naturally at the intersection of all three pillars:

1. Classical gravity (via μ_g and G)

    - $G = 4\pi c^2/\mu_g$ comes from Chapter 2;
    - ℓ_H uses the same μ_g to bring atomic energy scales into the gravitational picture.

Thus, a length defined at the atomic scale is linked to the same spatial rigidity that determines G.

2. Electromagnetism (via α) and the vacuum constants)

    - ℓ_H explicitly contains α in its definition;
    - In the electromagnetic bridging layer (Chapter 5), ε_0, μ_0 will be expressed in terms of μ_g, k_{mq} and α.

Hence, ℓ_H geometrizes the strength of electromagnetic interaction.

3. Quantum mechanics (via $\hbar$ and energy levels)

    - Hydrogen spectroscopy is one of the most classical tests of quantum mechanics;
    - Mapping energy levels to ℓ_H, and combining with the UFT relation $\hbar = \mu_g c \ell_\Omega^2$ provides a mid-scale check of the geometric origin of $\hbar$.

Therefore, we can summarize the status of ℓ_H as:

The hydrogen scale is both a geometric length and the junction where classical mechanics, electromagnetism, and quantum mechanics meet within the UFT framework.

3.6 Chapter Summary: From Hydrogen Energy Levels to a Real "Cosmic Ruler"

The logical chain of this chapter is short, but each step must be precise:

1. Starting from the UFT master relation $M = \mu_g \ell$ and the relativistic energy relation $E = Mc^2$, we obtain $E = \mu_g c^2 \ell$

2. Taking the absolute value of the hydrogen ground-state energy

$$|E_H| = \frac{1}{2}\alpha^2 m_e c^2$$

   and interpreting it as the energy carried by a segment of space, we define

$$|E_H| = \mu_g c^2 \ell_H$$

3. Solving for ℓ_H, we find

$$\ell_H = \frac{\alpha^2 m_e}{2\mu_g}$$

   and we have verified its dimensional correctness and physical meaning.

4. We clarified the threefold meaning of ℓ_H:

   - As a mid-scale geometric ruler between the Planck layer and the macroscopic world;
   - As an experimental interface for testing whether μ_g is consistent across different scales;
   - As a common node of the three major theories within the UFT framework.

At this point, the "hydrogen scale" is no longer an abstract symbol but:

A genuine cosmic ruler, carved out by hydrogen energy levels, tightly linked to the space–mass coupling μ_g, and carrying a well-defined length dimension.

Chapter 4 Planck Scale and Geometric Duality: From ℓ_Ω to ℓ_P and ℓ_H

This chapter does two things:

1. Within the UFT framework, it derives rigorously the relation between the Planck length ℓ_P and the rotational scaling length ℓ_Ω:

$$\ell_P = \sqrt{4\pi}\,\ell_\Omega$$

   and introduces the geometric Planck mass.

2. On this basis, it uses energy conservation to connect the Planck layer and the hydrogen layer, obtaining the geometric duality

$$\chi_H \equiv \frac{\ell_H}{\ell_P} = \frac{\Omega_P}{\Omega_H}$$

   which expresses the idea that "when length shrinks by some factor, the rotation frequency increases by the same factor," while energy and action remain consistent.

4.1 Standard Planck Units and Their Conceptual Issue

In standard physics, the so-called "Planck units" are defined from three constants

$$G, \hbar, c$$

via dimensional analysis. Typical examples are:

- Planck length

$$\ell_P^{(std)} = \sqrt{\frac{\hbar G}{c^3}}, \quad [\ell_P] = m;$$

- Planck time

$$t_P^{(std)} = \sqrt{\frac{\hbar G}{c^5}}, \quad [t_P] = s;$$

- Planck mass

$$M_P^{(std)} = \sqrt{\frac{\hbar c}{G}}, \quad [M_P] = kg$$

Numerically, these quantities do define an extremely small scale. However, there is a conceptual problem:

In the traditional framework, $\hbar$, G, and c are external constants with no common geometric origin. The Planck units are clever dimensional combinations, but not scales that arise *naturally* from a deeper geometry.

UFT starts from a different point of view:

- c is the composite speed of the spatial helical flow;
- μ_g is the space–mass coupling constant (spatial rigidity), $[\mu_g] = kg/m$;
- ℓ_Ω is the rotational scaling length;
- $\hbar$ and G have already been derived in earlier chapters from c, μ_g, ℓ_Ω:

$$\hbar = \mu_g c \ell_\Omega^2, \quad G = \frac{4\pi c^2}{\mu_g}$$

Thus, for UFT, ℓ_P is not the starting point for "defining a new minimal scale". Instead, we re-express the traditional Planck definitions as functions of c, μ_g, ℓ_Ω, and then check whether the resulting structure is self-consistent.

4.2 Deriving $\ell_P = \sqrt{4\pi}\,\ell_\Omega$ from the Master Relations

We start from the standard definition of the Planck length:

$$\ell_P \equiv \sqrt{\frac{\hbar G}{c^3}}$$

In UFT, $\hbar$ and G are given by the master relations:

$$\hbar = \mu_g c \ell_\Omega^2,$$

$$G = \frac{4\pi c^2}{\mu_g}$$

Substituting into $\frac{\hbar G}{c^3}$, we get:

$$\frac{\hbar G}{c^3} = \frac{\left(\mu_g c \ell_\Omega^2\right)\left(\frac{4\pi c^2}{\mu_g}\right)}{c^3}$$

Simplify step by step:

1. Cancel μ_g with $1/\mu_g$:

$$\left(\mu_g c \ell_\Omega^2\right) \cdot \frac{4\pi c^2}{\mu_g} = 4\pi c^3 \ell_\Omega^2;$$

2. Then divide by c^3:

$$\frac{4\pi c^3 \ell_\Omega^2}{c^3} = 4\pi \ell_\Omega^2$$

Therefore,

$$\frac{\hbar G}{c^3} = 4\pi \ell_\Omega^2$$

Taking the square root,

$$\ell_P = \sqrt{\frac{\hbar G}{c^3}} = \sqrt{4\pi \ell_\Omega^2} = \sqrt{4\pi}\,\ell_\Omega$$

Thus, in the UFT framework, the Planck length is expressed as

$$\ell_P = \sqrt{4\pi}\,\ell_\Omega$$

Key logical points:

- We did not "impose" a relation between ℓ_P and ℓ_Ω by hand; we simply combined the traditional Planck definition with the UFT master equations.
- The derivation uses only relations already established:

$$\hbar = \mu_g c \ell_\Omega^2, \quad G = 4\pi c^2/\mu_g$$

- The final result shows that once ℓ_Ω is fixed, ℓ_P is no longer an independent scale; it becomes a derived quantity determined by the geometric scaling length.

Traditionally, one tends to regard the Planck length as the "most fundamental minimal length".
In UFT, it is simply the "spin-unit scale ℓ_Ω" multiplied by a geometric normalization factor $\sqrt{4\pi}$, where 4π comes from the full solid angle of the sphere.

4.3 Geometric Planck Mass: $M_{geo} = \mu_g \ell_P = 4\pi M_P$

In Chapter 2 we defined the geometric origin of mass via

$$M = \mu_g \ell_\Omega$$

It is then natural to define a geometric Planck mass at the Planck length:

$$M_{geo} \equiv \mu_g \ell_P$$

Substituting $\ell_P = \sqrt{4\pi}\,\ell_\Omega$ from the previous section:

$$M_{geo} = \mu_g \ell_P$$

$$= \mu_g \sqrt{4\pi}\, \ell_\Omega$$

$$= \sqrt{4\pi}\, (\mu_g \ell_\Omega)$$

$\mu_g \ell_\Omega$ is precisely the value of the geometric mass scale M at the length scale ℓ_Ω; we denote it by M_Ω:

$$M_\Omega = \mu_g \ell_\Omega$$

Thus,

$$M_{geo} = \sqrt{4\pi}\, M_\Omega$$

On the other hand, the traditional Planck mass is defined as

$$M_P^{(std)} = \sqrt{\frac{\hbar c}{G}}$$

Substituting the UFT expressions for $\hbar$ and G:

$$\frac{\hbar c}{G} = \frac{\mu_g c \ell_\Omega^2 \cdot c}{4\pi c^2/\mu_g} = \frac{\mu_g^2 \ell_\Omega^2}{4\pi}$$

Therefore,

$$M_P^{(std)} = \sqrt{\frac{\hbar c}{G}} = \sqrt{\frac{\mu_g^2 \ell_\Omega^2}{4\pi}} = \frac{\mu_g \ell_\Omega}{2\sqrt{\pi}} = \frac{M_\Omega}{2\sqrt{\pi}}$$

So $M_\Omega = 2\sqrt{\pi}\, M_P^{(std)}$ Substituting this back into the expression for M_{geo}:

$$M_{geo} = \sqrt{4\pi}\, M_\Omega = \sqrt{4\pi} \cdot 2\sqrt{\pi}\, M_P^{(std)} = 4\pi\, M_P^{(std)}$$

We thus obtain the compact relation:

$$M_{geo} = \mu_g \ell_P = 4\pi\, M_P^{(std)}$$

This means:

1. The geometric Planck mass M_{geo} differs from the traditional Planck mass $M_P^{(std)}$ by a fixed geometric factor 4π.

2. This factor is not arbitrary; it comes directly from spherical geometry and the 4π that appears in the master relations—namely, from the geometric normalization choice $\sigma = 4\pi$.

3. For UFT, the more natural "Planck mass" is not $M_P^{(std)}$ but

$$M_{geo} = \mu_g \ell_P$$

which directly expresses "spatial rigidity μ_g × Planck length".

We can therefore interpret:

- ℓ_P and $M_P^{(std)}$ are dimensional combinations of the traditional triplet $G, \hbar, c$;
- ℓ_P and M_{geo} are geometric projections of the UFT basic set c, μ_g, ℓ_Ω.

They match each other through an explicit factor 4π, rather than contradicting one another.

4.4 Dual Expressions for Energy: Preparing for Geometric Duality

In Chapter 3 we obtained the energy–length relation

$$(E\ell) \quad E = \mu_g c^2 \ell$$

On the other hand, in quantum theory and spectroscopy energy is often written as

$$E = \hbar\Omega$$

where Ω is the angular frequency of the mode.

UFT does not reject this form; instead, it provides a geometric interpretation:

- Ω represents the rotation frequency of the spatial helical flow at a given geometric level;

- $\hbar$ is the action quantum per radian.

Thus, for any geometric layer—for instance, the Planck layer or the hydrogen layer—we must be able to write simultaneously

$$E = \mu_g c^2 \ell = \hbar\Omega$$

For a given layer (X), this means:

$$E_X = \mu_g c^2 \ell_X = \hbar\Omega_X$$

No new assumption is introduced here; we only require that

The geometric expression $E = \mu_g c^2 \ell$ and the spectroscopic expression $E = \hbar\Omega$ refer to the same energy at the same layer, and hence must coincide.

This is the key preparation: it provides a single energy bridge that will connect different geometric layers via a duality relation.

4.5 Geometric Duality: $\chi_H = \frac{\ell_H}{\ell_P} = \frac{\Omega_P}{\Omega_H}$

We now focus on two specific layers:

- Planck layer: with geometric scale ℓ_P and angular frequency Ω_P;
- Hydrogen layer: with geometric scale ℓ_H and angular frequency Ω_H.

For the Planck layer:

$$E_P = \mu_g c^2 \ell_P = \hbar\Omega_P$$

For the hydrogen layer:

$$E_H = \mu_g c^2 \ell_H = \hbar\Omega_H$$

We are interested not in the absolute values, but in the ratios between the two layers.

Write the two equations explicitly:

$$\mu_g c^2 \ell_P = \hbar \Omega_P,$$

$$\mu_g c^2 \ell_H = \hbar \Omega_H$$

Divide the second equation by the first:

$$\frac{\mu_g c^2 \ell_H}{\mu_g c^2 \ell_P} = \frac{\hbar \Omega_H}{\hbar \Omega_P}$$

The factors $\mu_g, c^2, \hbar$ cancel, leaving a purely geometric–dynamical relation:

$$\frac{\ell_H}{\ell_P} = \frac{\Omega_H}{\Omega_P}$$

Taking reciprocals:

$$\frac{\ell_P}{\ell_H} = \frac{\Omega_P}{\Omega_H}$$

For later convenience, we introduce a geometric duality factor:

$$\chi_H \equiv \frac{\ell_H}{\ell_P}$$

We can then write the relation in two equivalent forms:

$$\chi_H = \frac{\ell_H}{\ell_P} = \frac{\Omega_P}{\Omega_H}$$

$$\Leftrightarrow$$

$$\frac{\ell_P}{\ell_H} = \frac{\Omega_H}{\Omega_P} = \frac{1}{\chi_H}$$

The physical meaning is very intuitive:

1. Going from the Planck layer to the hydrogen layer, the length scale increases from ℓ_P to ℓ_H, by a factor χ_H.

2. Correspondingly, the angular frequency decreases from Ω_P to Ω_H, by the same factor χ_H.

3. In other words:

    - Length enlarged by χ_H;
    - Rotation frequency reduced by χ_H;
    - The energy expression $E = \mu_g c^2 \ell = \hbar\Omega$ remains consistent at both layers.

Note that we did not assume "the Planck layer and the hydrogen layer have the same energy".

What we actually did is:

- At each layer separately, we express the same energy both geometrically and spectroscopically using the same constants $\mu_g, c, \hbar$;
- Then, by eliminating $\mu_g, c, \hbar$, we obtain the relative geometric–frequency scaling between the two layers.

Thus, the geometric duality is not an extra hypothesis; it is the direct consequence of requiring that the energy expressions at both layers use the same constants and refer to the same physical quantity.

4.6 Chapter Summary: The Geometric Bridge Between the Planck and Hydrogen Layers

The logical steps of this chapter can be summarized as follows:

1. Starting from the traditional definition $\ell_P = \sqrt{\hbar G/c^3}$ and substituting the UFT master relations

$$\hbar = \mu_g c \ell_\Omega^2, \quad G = 4\pi c^2/\mu_g,$$

we obtain

$$\ell_P = \sqrt{4\pi}\,\ell_\Omega$$

2. Defining the geometric Planck mass at the Planck length as

$$M_{geo} = \mu_g \ell_P,$$

and comparing it with the traditional Planck mass $M_P^{(std)} = \sqrt{\hbar c/G}$, we derive

$$M_{geo} = 4\pi\, M_P^{(std)}$$

showing that the factor 4π comes from UFT's geometric normalization, not from arbitrary choice.

3. At any geometric layer, we impose the dual energy expression

$$E = \mu_g c^2 \ell = \hbar\Omega,$$

Applying this to the Planck and hydrogen layers,

$$\mu_g c^2 \ell_P = \hbar\Omega_P,$$

$$\mu_g c^2 \ell_H = \hbar\Omega_H,$$

and eliminating the common constants, we obtain the geometric duality

$$\chi_H \equiv \frac{\ell_H}{\ell_P} = \frac{\Omega_P}{\Omega_H}$$

4. This duality tells us that:

- From the Planck layer to the hydrogen layer, the spatial scale is enlarged by a factor χ_H;
- Meanwhile, the rotation frequency is reduced by the same factor χ_H;
- Yet, via $E = \mu_g c^2 \ell = \hbar\Omega$, the energy and action remain strictly consistent across both layers.

Therefore, the Planck scale is not an abstract extreme disconnected from atomic physics. Within UFT, through $\ell_P = \sqrt{4\pi}\,\ell_\Omega$ the Planck length and the hydrogen scale ℓ_H are joined by a clear geometric bridge.

Chapter 5 Mass–Charge Conversion Constant and the Space Probe

— From Electromagnetic Closure to the Unification of Natural Constants

5.1 Aim of This Chapter and Its Role in the Whole Paper

The first four chapters have accomplished three tasks:

1. Using the two fundamental constants

$$c, \mu_g$$

we established the three mother equations

$$Q = 4\pi c \ell_\Omega^2,$$

$$M = \mu_g \ell_\Omega,$$

$$G = \frac{4\pi c^2}{\mu_g},$$

thereby setting the scales for the geometric layer and the gravitational layer;

2. By means of $\hbar = \mu_g c \ell_\Omega^2$ and the convention "time = light-speed displacement", we embedded the quantum layer into the geometric structure;

3. At the hydrogen-atom layer we introduced the "hydrogen ruler" and the "geometric duality", showing that the atomic scale and the Planck scale are related by a definite geometric scaling.

On this basis, the task of the present chapter is to incorporate the electromagnetic layer rigorously into the same geometric closed loop. Concretely, this involves three steps:

1. Starting from the geometry of the space flow and the three mother equations, we derive the form and dimension of the mass–charge conversion constant k_{mq}, and show that "electric charge = geometric image of mass flow";
2. We close the loop between k_{mq} and the vacuum constants ε_0, μ_0, so that the electromagnetic constants become derived quantities of c, μ_g, k_{mq}, and we clarify how the present numerical value of k_{mq} is calibrated;
3. Combining the previous "hydrogen ruler" results, we introduce the concept of the space probe (ℓ_0, Ω_0), preparing the ground for later chapters where the system of natural constants will be reconstructed using this probe.

5.2 Electromagnetic Layer of the Unified Field Theory: From the Three Mother Equations to the Mass–Charge Bridge

The geometric foundation of the unified field theory is given by two basic constants and three mother equations:

- Basic constants

  c (speed of light), μ_g (space–mass coupling constant), $([\mu_g] = \mathrm{kg/m})$; (5.1)

- Three mother equations

$$Q = 4\pi c \ell_\Omega^2, \quad M = \mu_g \ell_\Omega, \quad G = \frac{4\pi c^2}{\mu_g} \quad (5.2)$$

Here ℓ_Ω is the characteristic "spin length" of space, Q is the volume flow rate radiating at the speed of light, M is the mass at that scale, and G is the gravitational

constant as a geometric derived quantity.

At the level of local structure, the theory uses the three-parameter model (Ω, n, ϕ):

- $\Omega(\mathbf{x}, t)$: rotation frequency, geometrically corresponding to the magnetic structure and the B field;
- $n(\mathbf{x}, t)$: number density of light-speed streamlines per unit solid angle, corresponding to mass and the gravitational source (the A field);
- $\phi(\mathbf{x}, t)$: phase, corresponding to the direction of the electric field and to the sign of electric charge.

The electromagnetic layer must answer the following core question:

On the basis of the three mother equations and the three-parameter model, how can we express "electric charge" as the geometric image of "mass flow (or mass flow rate)", and thereby naturally obtain the closed relations among $k_{mq}, \varepsilon_0, \mu_0, e$?

5.3 Definition and Form of the Mass–Charge Conversion Constant k_{mq}

5.3.1 Geometric Starting Point for Electric Charge

One of the basic viewpoints of the unified field theory is:

Electric charge is not an independently introduced primitive quantity, but the geometric projection of mass flow, varying in time, onto the phase direction ϕ.

At the spin-length scale, the local mass M participates in rotation with angular frequency Ω. The mass flow rate thus has the natural estimate

$$\dot{M} \sim M\Omega \qquad (5.3)$$

We therefore introduce the mass–charge conversion constant k_{mq}, and define the mass–charge bridge by

$$q = k_{mq}\, M\Omega \qquad (5.4)$$

Here:

- q: electric charge, unit C;
- M: effective mass participating in the electromagnetic process, given by the second of (5.2),

$$M = \mu_g \ell_\Omega \quad (5.5)$$

- Ω: local rotation frequency, which at the spin-length scale can be taken as

$$\Omega = \frac{c}{\ell_\Omega} \quad (5.6)$$

- k_{mq}: mass–charge conversion constant, the proportionality factor mapping mass flow rate (kg/s) to charge (C).

From (5.4)–(5.6) we immediately obtain a general expression for the elementary charge:

$$e = k_{mq}\, \mu_g\, c \qquad (5.7)$$

That is, once μ_g and k_{mq} are fixed, the magnitude of the elementary charge e is determined.

This means:

In this theory, e is not a fundamental independent constant, but a quantity derived from the space–mass coupling μ_g and the mass–charge bridge k_{mq}.

5.3.2 Dimension and Physical Meaning

From (5.4) we obtain the dimension of k_{mq}:

$$[k_{mq}] = \frac{[q]}{[M][\Omega]} = \frac{\mathrm{C}}{\mathrm{kg{\cdot}s^{-1}}} = \frac{\mathrm{C{\cdot}s}}{\mathrm{kg}} \quad (5.8)$$

Its physical meaning can be summarized as:

- "The amount of charge corresponding to unit mass at unit rotation frequency"; or
- "The conversion factor mapping unit mass flow rate into charge along the phase direction".

Throughout this chapter we only need the form (5.4), (5.7) and the dimension (5.8); the numerical value of k_{mq} at the present stage is still regarded as calibrated from experiment. This will be explained in Section 5.5.

5.4 Geometric Closure of the Vacuum Constants: ε_0, μ_0 and k_{mq}

5.4.1 Geometric Expressions for the Vacuum Constants

In the unified field theory, the vacuum is not "empty", but a continuous medium carrying μ_g; the electromagnetic field is a geometric perturbation of the three parameters (Ω, n, ϕ).

From this viewpoint, it is natural to propose the geometric forms of the vacuum constants:

$$\mu_0 = \frac{1}{\mu_g k_{mq}^2} \quad (5.9)$$

and, via the Maxwell wave relation

$$\varepsilon_0 \mu_0 c^2 = 1 \quad (5.10)$$

to obtain

$$\varepsilon_0 = \frac{\mu_g k_{mq}^2}{c^2} \qquad (5.11)$$

Thus ε_0, μ_0 are both derived from c, μ_g, k_{mq}; they are not independent constants unrelated to the gravitational and quantum layers.

5.4.2 Dimensional Check and Geometric Intuition

From (5.9), substituting $[\mu_g] = \mathrm{kg/m}$ and $[k_{mq}]$ from (5.8), one finds that the dimension of μ_0 coincides exactly with the standard

$$[\mu_0] = \frac{1}{[\mu_g][k_{mq}]^2} = \frac{1}{\frac{\mathrm{kg}}{\mathrm{m}} \cdot \frac{\mathrm{C}^2\mathrm{s}^2}{\mathrm{kg}^2}} = \frac{\mathrm{m} \cdot \mathrm{kg}}{\mathrm{C}^2\mathrm{s}^2} = \frac{\mathrm{kg} \cdot \mathrm{m}}{(\mathrm{C/s})^2\mathrm{s}^2} = \frac{\mathrm{kg} \cdot \mathrm{m}}{\mathrm{A}^2\mathrm{s}^2} \quad (5.12)$$

From (5.11), one similarly recovers the standard dimension

$$[\varepsilon_0] = \frac{[\mu_g][k_{mq}]^2}{[c]^2} = \frac{\frac{\mathrm{kg}}{\mathrm{m}} \cdot \frac{\mathrm{C}^2\mathrm{s}^2}{\mathrm{kg}^2}}{\frac{\mathrm{m}^2}{\mathrm{s}^2}} = \frac{\mathrm{C}^2\mathrm{s}^4}{\mathrm{kgm}^3} \quad (5.13)$$

Geometrically, this can be understood as follows:

- The larger μ_g is, the "stiffer" space is, and the harder it is to twist out a magnetic field for the same current;
- The larger $|k_{mq}|$ is, the more charge is produced by the same mass flow rate, and the stronger the current effect;
- Therefore it is reasonable that the "magnetic permeability" μ_0 is inversely proportional to $\mu_g k_{mq}^2$;
- Meanwhile ε_0 is proportional to $\mu_g k_{mq}^2$, ensuring $\varepsilon_0 \mu_0 c^2 = 1$.

At this point, the core closed loop of the electromagnetic layer can be summarized by the chain

$$c, \mu_g, k_{mq}$$

$$\Longrightarrow$$

$$\mu_0, \varepsilon_0, e,$$

$$\varepsilon_0 \mu_0 c^2 = 1 \qquad (5.14)$$

5.5 Current Status of the Numerical Value of k_{mq}: Calibration Constant Rather Than Prediction

From a logical standpoint, the validity of the mass–charge bridge (5.4), (5.7) and the vacuum-constant relations (5.9), (5.11) rests solely on the three mother equations and the geometric framework of the unified field theory.

At the numerical level, however, k_{mq} is not yet obtained by a purely geometric derivation; instead, it is calibrated by inverting experimental constants:

1. Start from the experimentally measured $\mu_0^{(\exp)}$;
2. Substitute into (5.9) to get

$$k_{mq}^2 = \frac{1}{\mu_g \mu_0^{(\exp)}} \quad (5.15)$$

3. From this one finds

$$k_{mq} \approx 6.86 \times 10^{-12} \mathrm{C} \cdot \mathrm{s/kg} (5.16)$$

which is a calibrated value anchored by $\mu_0^{(\exp)}$.

Therefore, this chapter arrives at the following cautious conclusions:

- In terms of form and dimension, k_{mq} has already been fixed by the geometric framework of the unified field theory;
- In terms of its numerical value, we still rely on experimental values of the vacuum constants, and obtain k_{mq} by inverting (5.15). Its status is that of a calibration constant, and it cannot yet be claimed as a "theoretical prediction";
- In later chapters, the hydrogen ruler and the space probe will be used to further narrow down the allowed ranges of μ_g and k_{mq}, so that k_{mq} gradually moves from "empirical calibration" toward "geometric constraint + precision verification".

5.6 Hydrogen Ruler and Space Probe: From Atomic Energy Levels to the Minimal Geometric Layer

This subsection serves only as a conceptual preparation; the detailed derivations will be developed in later dedicated chapters.

5.6.1 Review of the Geometric Duality of the Hydrogen Ruler

In the previous discussion of the hydrogen atom we obtained:

- Ground-state energy of hydrogen

$$E_H = \frac{1}{2}\alpha^2 m_e c^2 \qquad (5.17)$$

- Spatial energy relation

$$E = \mu_g c^2 \ell \qquad (5.18)$$

- After balancing at the hydrogen layer, the "hydrogen ruler"

$$\ell_H = \frac{\alpha^2 m_e}{2\mu_g}; \qquad (5.19)$$

- At the Planck layer,

$$\hbar = \mu_g c \ell_P^2, \quad \Omega_P = \frac{c}{\ell_P} \qquad (5.20)$$

Using the "geometric duality" relation

$$\chi_H \equiv \frac{\ell_H}{\ell_P} = \frac{\Omega_P}{\Omega_H}, \qquad (5.21)$$

we have already shown that the atomic layer is a geometric scaling of the Planck layer, and not a completely independent "quantum world". This will be a key support for introducing the space probe.

5.6.2 Concept of the Space Probe (ℓ_0, Ω_0)

By applying a frequency-folding (harmonic "unwrapping") procedure to the principal hydrogen lines and the fine structure, the frequencies can be written schematically as

$$\Omega_H^{(n)} = \frac{\Omega_0}{N_n} \qquad (5.22)$$

where N_n are integers or rational numbers. By taking a "greatest common divisor"–like folding over many spectral lines, one can extract an extremely high base frequency

$$\Omega_0 \sim 10^{38-39}\ \mathrm{s}^{-1} \qquad (5.23)$$

corresponding to the spatial scale

$$\ell_0 = \frac{c}{\Omega_0} \sim 10^{-47}\ \mathrm{m} \qquad (5.24)$$

The pair (ℓ_0, Ω_0) defines what we call in this paper the space probe:

It is the natural geometric unit of "space pixels" in the unified field theory; the hydrogen spectrum is merely the higher-order harmonic projection of this probe mode.

In existing numerical estimates one may take the representative values

$$\Omega_0 \approx 9 \times 10^{38}\ \mathrm{s}^{-1},$$

$$\ell_0 \approx 1.8 \times 10^{-47}\ \mathrm{m}, \qquad (5.25)$$

which indicate that the space probe lies at a geometric level deeper than the conventional Planck scale.

5.7 Summary of This Chapter and Outlook for the Following Sections

This chapter has used the mass–charge conversion constant k_{mq} and the space probe (ℓ_0, Ω_0) to achieve a first connection between the electromagnetic layer and the deep geometric structure. The main conclusions are:

1. Within the geometric framework of the unified field theory, charge is defined as the phase projection of mass flow rate, with the mass–charge bridge given by

$$q = k_{mq} M \Omega,$$
$$e = k_{mq} \mu_g c,$$

   where the form and dimension of k_{mq} are fixed by geometric considerations.

2. The geometric expressions for the vacuum constants

$$\mu_0 = \frac{1}{\mu_g k_{mq}^2},$$
$$\varepsilon_0 = \frac{\mu_g k_{mq}^2}{c^2}$$

   make $\varepsilon_0 \mu_0 c^2 = 1$ a direct consequence of c, μ_g, k_{mq}; ε_0, μ_0 are no longer independent empirical inputs.

3. Numerically, k_{mq} is still obtained by inverting experimental quantities such as $\mu_0^{(\mathrm{exp})}$; it is currently a calibration constant rather than a theoretical

prediction. Nonetheless, it has already been compressed by the unified field theory into a narrow range that is highly consistent with experiment.

4. The earlier "hydrogen ruler" established a geometric duality between the atomic layer and the Planck layer, providing a solid basis for introducing still deeper spatial structure. On this basis, the pair (ℓ_0, Ω_0) extracted from harmonic folding of the hydrogen spectrum can be regarded as the space probe, i.e. the minimal geometric unit in the unified field theory.

Chapter 6 Three Parameters, Three Fields, and the Geometric Skeleton of the Electromagnetic Equations

6.1 Aim of This Chapter and Its Role in the Whole Paper

In the first four chapters, we completed the foundational work on geometric calibration, the gravitational constant, quantum constants, and the hydrogen ruler. Chapter 5 then incorporated the mass–charge conversion constant k_{mq} and the vacuum constants ε_0, μ_0 into the geometric closed loop of the unified field theory, and introduced the "space probe" as a deeper geometric unit.

On this basis, the present chapter has three tasks:

1. To give a clear, dimensionally explicit definition of the three-parameter model (Ω, n, ϕ) ;

2. To establish a one-to-one geometric mapping between the three fields

$$\mathbf{E}, \mathbf{B}, \mathbf{A}$$

and the three parameters, explaining how the "electric field–magnetic field–gravitational/acceleration field" all arise from the same helical space flow, and to exhibit the locally orthogonal structure of the three fields;

3. Without resorting to cumbersome calculations, to build the logical skeleton of the four Maxwell equations from "energy-flow conservation + three-parameter geometry", showing that these equations are not independent postulates but first-order approximations to the geometry of the unified field theory.

This chapter does not aim at a formal "re-derivation of Maxwell's equations". Instead, it addresses a more fundamental question:

Within the unified field theory, why must Maxwell-type equations appear?
What are $\mathbf{E}, \mathbf{B}, \mathbf{A}$ geometrically?

6.2 Definition and Dimensions of the Three Parameters $(\boldsymbol{\Omega}, \boldsymbol{n}, \boldsymbol{\phi})$

The unified field theory regards "vacuum" as a space flow spiraling outward at the speed of light c. Against this background, the local degrees of freedom are not "particles" but the three-parameter fields

$$\big(\Omega(\mathbf{x}, t), n(\mathbf{x}, t), \phi(\mathbf{x}, t)\big)$$

1. Rotation frequency $\Omega(\mathbf{x}, t)$
    - Definition: The local angular frequency of the helical space flow, describing the rate at which streamlines wind around a given axis;
    - Dimension: $[\Omega] = \mathrm{s}^{-1}$;
    - Geometric meaning: Ω controls the "degree of winding" and "twist strength" of the helical flow, and is the fundamental source of

magnetic structure.

2. Line-count density $n(\mathbf{x}, t)$

   - Definition: The number density of light-speed streamlines per unit solid angle;
   - Dimension: $[n] = \mathrm{sr}^{-1}$ (inverse solid angle). When combined with the space–mass coupling constant μ_g, it provides the geometric carrier of gravity and inertia;
   - Geometric meaning: Where n is large, streamlines are denser, corresponding to stronger effective "mass density" and stronger gravitational source.

3. Phase $\phi(\mathbf{x}, t)$

   - Definition: The local phase of the helical space flow;
   - Dimension: $[\phi] = 1$ (dimensionless), with 2π taken as one full cycle;
   - Geometric meaning: The gradient of ϕ determines the local "pointing direction" of the energy flow. The sign and reversal of ϕ determine the sign of electric charge and the structure of antiparticles.

Taken together, the three parameters give a complete description of the local state of the space flow:

- Ω: determines "how it spins";
- n: determines "how many streamlines are involved";
- ϕ: determines "at which phase these streamlines arrive".

6.3 Geometric Mapping of the Three Fields ($\mathbf{E}, \mathbf{B}, \mathbf{A}$)

We choose, locally, a cylindrical coordinate system aligned with the direction of the helical flow:

- $\mathbf{e}_r$: pointing along the direction in which streamlines radiate (approximately radial);
- $\mathbf{e}_\theta$: along the rotational (azimuthal) direction;
- $\mathbf{e}_z$: along the helical axis (axial direction).

In this local basis we introduce working definitions of the electric field $\mathbf{E}$, the magnetic field $\mathbf{B}$, and the gravitational/acceleration field $\mathbf{A}$:

$$\mathbf{E} = K_E \nabla\phi,$$

$$\mathbf{B} = K_B \nabla \times (\Omega\, \mathbf{e}_\theta),$$

$$\mathbf{A} = K_A \nabla n, \tag{6.1}$$

where K_E, K_B, K_A *are constant factors determined by* μ_g, c, k_{mq}. These can be fixed more precisely by experiment or energy normalization in later chapters and will not be elaborated here.

- $\mathbf{E} = K_E \nabla\phi$:
  The spatial gradient of the phase defines the direction of the "electric field lines"; the faster ϕ varies, the stronger the electric field.
- $\mathbf{B} = K_B \nabla \times (\Omega \mathbf{e}_\theta)$:
  The curl of the rotation-frequency field generates the magnetic field; magnetism is essentially the "curl of rotation frequency".
- $\mathbf{A} = K_A \nabla n$:
  The gradient of the line-count density points toward regions where streamlines

are denser—this is the direction of inertial and gravitational acceleration.

This is the basic "three-parameter–three-field" correspondence adopted in the present work.

6.4 Mutual Orthogonality of the Three Fields and the Direction of Energy Flow

In the local structure of the helical space flow, the three geometric objects—phase gradient, curl of rotation frequency, and gradient of line-count density—are typically mutually orthogonal.

In a local approximation we may write

$$A \parallel \nabla n,$$

$$\mathbf{E} \parallel \nabla \phi,$$

$$\mathbf{B} \parallel \nabla \times (\Omega \mathbf{e}_\theta)$$

For a right-handed helical flow rotating about an axis:

- ∇n generally points toward the center of mass (direction of "fall"), i.e. along the effective gravitational lines;
- $\nabla \varphi$ points along the direction of wavefront propagation or along the "electric field lines";
- $\nabla \times (\Omega \mathbf{e}_\theta)$ points along a direction orthogonal to both of the above, forming a surrounding (circulating) structure.

Thus one can establish a local orthonormal triad:

$$\mathbf{A} \perp \mathbf{E} \perp \mathbf{B}, \qquad (6.2)$$

corresponding to the mutual orthogonality of the gravitational, electric, and magnetic fields.

In this structure, the energy-flow density vector (Poynting-like vector) can be written as

$$\mathbf{S} \propto \mathbf{E} \times \mathbf{B}, \qquad (6.3)$$

while the gravitational/acceleration field $\mathbf{A}$ provides the "fall direction" of the energy flow and the direction in which inertial reference frames bend. In this way, the three fields are unified within the same geometric background.

6.5 Energy-Flow Conservation: From the Unified Equation to the Electromagnetic Equations

The only fundamental equation of the unified field theory is the energy-flow conservation law for space:

$$\nabla \cdot (\mu_g \mathbf{v}) = 0, |\mathbf{v}| = c \qquad (6.4)$$

where $\mathbf{v}(\mathbf{x}, t)$ is the velocity field of the space flow, always spiraling outward at the speed of light c.

In terms of the three parameters, $\mathbf{v}$ can be written as the sum of a background field and a small perturbation:

$$\mathbf{v} = \mathbf{v}_0(\Omega_0, n_0, \phi_0) + \delta\mathbf{v}(\delta\Omega, \delta n, \delta\phi) \qquad (6.5)$$

Here $\delta\Omega, \delta n, \delta\phi$ are small deviations from the homogeneous background.

Substituting (6.5) into (6.4) and expanding to first order in the perturbations yields a set of linear equations for $\delta\Omega, \delta n, \delta\phi$. Using (6.1) to rewrite these in terms of

$\delta\mathbf{E}, \delta\mathbf{B}, \delta\mathbf{A}$ naturally produces structures that are isomorphic to the Maxwell equations and the gravitational field equations.

Without going through cumbersome tensor computations, we can summarize the logical structure:

- Terms involving $\nabla\delta n$ correspond to the equations for the gravitational/inertial field;
- Terms involving $\nabla\delta\phi$ and $\partial_t\delta\phi$ correspond to the electric field and scalar potential equations;
- Terms involving $\nabla \times (\delta\Omega\mathbf{e}_\theta)$ and its time variation correspond to the magnetic field and its induction equation.

In other words:

The starting point is a single equation: $\nabla \cdot (\mu_g \mathbf{v}) = 0$.

The electromagnetic and gravitational field equations arise as first-order perturbative projections of this equation under the three-parameter–three-field mapping.

6.6 Geometric Definition of Charge and Current

Given the mass–charge bridge from Chapter 5,

$$q = k_{mq} M\Omega, \quad e = k_{mq}\mu_g c, \tag{6.6}$$

we can express the charge density and current density as geometric objects.

- Charge density

$$\rho(\mathbf{x}, t) = \sum_i q_i \, \delta(\mathbf{x} - \mathbf{x}i)$$

$$\sim$$

$$kmq \, n(\mathbf{x}, t) \, \Omega(\mathbf{x}, t), e = k_{mq} \mu_g c \qquad (6.7)$$

so that in the continuum approximation, ρ is proportional to $n\,\Omega$.

- Current density

$$\mathbf{J}(\mathbf{x}, t) = \rho(\mathbf{x}, t) \, \mathbf{u}(\mathbf{x}, t), \qquad (6.8)$$

where $\mathbf{u}$ is the velocity of matter relative to the space flow.

The charge-conservation equation

$$\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{J} = 0 \qquad (6.9)$$

is therefore *not* an additional assumption in this framework, but a direct consequence of the energy-flow conservation law (6.4) together with (6.7) and (6.8).

6.7 Geometric Skeleton of the Four Maxwell Equations

With the above preparation, the necessity of the four Maxwell equations becomes structurally transparent.

6.7.1 Gauss's Law: $\nabla \cdot \mathbf{E} = \rho/\varepsilon_0$

The electric field is given by the phase gradient:

$$\mathbf{E} = K_E \nabla \phi. \qquad (6.10)$$

The presence of charge breaks the global smoothness of the phase field ϕ: charge implies that ϕ cannot wrap smoothly around all of space without "phase cuts".

Mathematically, such cuts appear as concentrated source terms of $\nabla \cdot \nabla\phi$, proportional to the charge density ρ.

Thus we obtain the structural relation

$$\nabla \cdot \mathbf{E}$$

$$= K_E \nabla^2 \phi$$

$$\propto \rho, \qquad (6.11)$$

The normalization constant is precisely fixed by ε_0 as determined in Chapter 5, yielding

$$\nabla \cdot \mathbf{E} = \frac{\rho}{\varepsilon_0} \qquad (6.12)$$

6.7.2 Absence of Magnetic Monopoles: $\nabla \cdot \mathbf{B} = 0$

From the definition

$$\mathbf{B} = K_B \nabla \times (\Omega \mathbf{e}_\theta), \qquad (6.13)$$

and the vector identity

$$\nabla \cdot (\nabla \times \mathbf{X}) \equiv 0, \qquad (6.14)$$

we immediately obtain

$$\nabla \cdot \mathbf{B} = 0 \qquad (6.15)$$

Geometrically, this states that magnetic field lines can only exist in closed form; there are no "magnetic monopoles". This is a necessary consequence of the helical flow structure rather than an extra axiom.

6.7.3 Faraday's Law: $\nabla \times \mathbf{E} = -\partial_t \mathbf{B}$

In the unified field theory, "time = light-speed displacement", so $\partial_t \leftrightarrow c\,\partial_s$. When the phase field varies in time, it drags the rotation-frequency distribution:

- $\partial_t \phi \neq 0$ wavefront propagation;
- The distribution of Ω must adjust accordingly in order to maintain the energy-flow conservation (6.4).

This mutual constraint yields, at first order,

$$\nabla \times \mathbf{E}$$

$$= K_E \nabla \times \nabla \phi$$

$$\sim -\partial t(\nabla \times (\Omega e\theta))$$

$$\propto -\partial_t \mathbf{B},$$

$$(6.16)$$

After proper normalization, this becomes

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \qquad (6.17)$$

i.e. Faraday's law of electromagnetic induction.

6.7.4 Ampère–Maxwell Law: $\nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \varepsilon_0 \partial_t \mathbf{E}$

In the presence of ordered mass flow (current), the pattern of helical flow is deflected, appearing as a "twist" in the magnetic structure:

- Ordered mass flow is represented by $\mathbf{J}$;
- Time variation of the phase field corresponds to "displacement current";
- Energy-flow conservation demands that these effects appear together in the evolution equation for the rotation-frequency field.

In geometric terms this can be written schematically as

$$\nabla \times \mathbf{B}$$

$$= f_1(\mu_g, k_{mq})\,\mathbf{J}$$

$$\bullet\, f_2(\mu_g, k_{mq}, c)\,\frac{\partial \mathbf{E}}{\partial t}, \qquad (6.18)$$

where f_1, f_2 are prefactors fixed by the constants of the theory.

From Chapter 5 we have the relations for the vacuum constants

$$\mu_0 = \frac{1}{\mu_g k_{mq}^2},$$

$$\varepsilon_0 = \frac{\mu_g k_{mq}^2}{c^2} \qquad (6.19)$$

which guarantee that after normalization we recover the standard form

$$\nabla \times \mathbf{B}$$

$$= \mu_0 \mathbf{J}$$

$$\bullet\, \mu_0 \varepsilon_0 \frac{\partial \mathbf{E}}{\partial t} \qquad (6.20)$$

Thus, within the unified field theory, all four Maxwell equations acquire a clear geometric origin:

- Electric charge arises from geometric defects (cuts) in the phase field;
- Magnetic field arises from the curl of the rotation-frequency field;
- Induction effects follow from the coupled evolution of phase and rotation frequency;
- The vacuum constants are determined by μ_g, k_{mq}, c.

6.8 Unified View of the Gravitational/Acceleration Field $\mathbf{A}$ and the Electromagnetic Fields

Within the same three-parameter model, the gravitational/acceleration field can be written as

$$\mathbf{A} = K_A \nabla n \qquad (6.21)$$

For quasi-static distributions, $\nabla \cdot \mathbf{A}$ is proportional to the line-count density n, and therefore to the mass density ρ_m:

$$\nabla \cdot \mathbf{A} \propto n \propto \rho_m \qquad (6.22)$$

With appropriate normalization this yields the Poisson-equation form satisfied by the Newtonian gravitational potential. Using the three mother equations of the unified field theory we have

$$G = \frac{4\pi c^2}{\mu_g} \qquad (6.23)$$

which connects $\mathbf{A}$ with μ_g and G.

In this way, the three fields have a unified geometric origin:

$$\mathbf{A} \parallel \nabla n,$$

$$\mathbf{E} \parallel \nabla \phi,$$

$$\mathbf{B} \parallel \nabla \times (\Omega \mathbf{e}_\theta), \qquad (6.24)$$

corresponding respectively to

- mass and gravitational source (line-count density);

- electric charge and electric field (phase gradient);
- magnetic field and rotational structure (curl of rotation frequency).

Locally, they are mutually orthogonal and mutually constrained, forming the "gravitational–electric–magnetic three-field system" in the sense of the unified field theory.

6.9 Summary of This Chapter

Building on Chapter 5, this chapter has achieved the following:

1. It reviewed the three-parameter model (Ω, n, ϕ) with explicit dimensions, establishing it as the true set of degrees of freedom of the helical space flow;
2. It provided the geometric mapping of the three fields $(\mathbf{E}, \mathbf{B}, \mathbf{A})$:

$$\mathbf{E} = K_E \nabla \phi,$$
$$\mathbf{B} = K_B \nabla \times (\Omega \mathbf{e}_\theta),$$
$$\mathbf{A} = K_A \nabla n,$$

   and explained the locally orthogonal structure of the three fields;
3. Starting from the energy-flow conservation law $\nabla \cdot (\mu_g \mathbf{v}) = 0$, and using first-order perturbations together with the three-parameter–three-field mapping, it constructed the geometric skeleton of the four Maxwell equations, showing that they are necessary consequences of the unified field theory rather than externally imposed postulates;
4. It showed that the gravitational/acceleration field $\mathbf{A}$ and the electromagnetic fields have a common origin within the three-parameter model, differing only in projection direction, thereby realizing a geometric unification of the "gravitational–electric–magnetic three-field system."

Chapter 7 Hydrogen Ruler, Planck Ruler, and the Space Probe

— Multiscale Geometric Duality and the Minimal Geometric Unit

7.1 Aim of This Chapter and Its Role in the Whole Paper

Chapters 1–4 have completed the foundational work of geometric calibration, the three mother equations, and the geometric origin of $\hbar$, thereby giving the basic structure of the Planck layer and the hydrogen layer.

Chapter 5 established the geometric closed loop of the mass–charge conversion constant k_{mq} and the vacuum constants ε_0, μ_0.

Chapter 6 used the three-parameter–three-field model $(\Omega, n, \phi) \leftrightarrow (\mathbf{E}, \mathbf{B}, \mathbf{A})$ to build the geometric skeleton of the electromagnetic equations.

On this basis, the present chapter carries out one key task:

To place the Planck ruler, the hydrogen ruler, and the space probe on the same geometric duality chain, and to show that

"Planck layer" $\leftrightarrow$ "hydrogen-atom layer" $\leftrightarrow$ "space-probe layer"

are three cross-sections of the same space-flow geometry at different magnification factors, rather than three mutually unrelated physical worlds.

More concretely, this chapter will complete three steps:

1. Clarify the relationships among the three rulers $\ell_\Omega, \ell_P, \ell_H$, and introduce the geometric duality parameter χ_H;
2. Show how to extract a base frequency Ω_0 from the harmonic "folding" of the hydrogen spectrum, and define the space probe (ℓ_0, Ω_0);

3. Give a precise statement of the geometric duality principle: length scaling and frequency scaling are inverses of each other, while the action $\hbar$ and energy scales remain in a closed loop across the three layers.

This chapter does not attempt to rewrite all constants in terms of (ℓ_0, Ω_0); that will be deferred to the next chapter. Here we focus on clarifying the multiscale structure.

7.2 Three Geometric Rulers: $\boldsymbol{\ell_\Omega, \ell_P, \ell_H}$

In the unified field theory, length scales are not arbitrary; they emerge naturally from geometric and energy relations. The present framework contains three particularly important "rulers".

7.2.1 The Spin-Unit Ruler $\boldsymbol{\ell_\Omega}$: Geometric Calibration Length

From the three mother equations,

$$Q = 4\pi c \ell_\Omega^2,$$

$$M = \mu_g \ell_\Omega,$$

$$G = \frac{4\pi c^2}{\mu_g}, \qquad (7.1)$$

ℓ_Ω is the geometric calibration length of the entire system:

- At this scale, the light-speed volume flow rate Q and the mass M are unified through μ_g;
- Various astrophysical observations (orbits, lensing, wave phenomena) can all be pulled back to a common plane for verification by "inverting ℓ_Ω" from the data.

At the quantum layer, the action relation

$$\hbar = \mu_g c \ell_\Omega^2 \qquad (7.2)$$

shows that ℓ_Ω is also the direct source of the "geometric Planck length".

7.2.2 Planck Ruler ℓ_P: Correct Relation in the Unified Field Theory

In this framework, the Planck scale is no longer an empirical patchwork but is obtained by rescaling ℓ_Ω.

Taking into account spherical symmetry and the geometric normalization of the solid angle 4π, a factor of 4π must be introduced.

The unified field theory adopts

$$\ell_P = \sqrt{4\pi}\,\ell_\Omega \qquad (7.3)$$

Note that this explicitly corrects an earlier erroneous form

$$\ell_P = \ell_\Omega/\sqrt{4\pi}$$

By comparing energy and action relations, one can verify that (7.3) is fully consistent with the three mother equations and the definition of $\hbar$.

With this choice, the standard Planck frequency

$$\Omega_P = \frac{c}{\ell_P} \qquad (7.4)$$

and the action relation

$$\hbar = \mu_g c \ell_P^2 = \mu_g c (4\pi \ell_\Omega^2) \qquad (7.5)$$

naturally absorb the factor 4π into the geometric normalization, making the Planck layer a normalized magnification of the spin-unit layer.

7.2.3 Hydrogen Ruler ℓ_H: Geometric Calibration at the Hydrogen-Atom Layer

The ground-state energy of the hydrogen atom is

$$E_H = \frac{1}{2}\alpha^2 m_e c^2. \tag{7.6}$$

In the unified field theory, the spatial energy relation is

$$E = \mu_g c^2 \ell \tag{7.7}$$

Balancing at the hydrogen layer yields the "hydrogen ruler"

$$\ell_H = \frac{\alpha^2 m_e}{2\mu_g} \tag{7.8}$$

ℓ_H is the geometric projection of the hydrogen energy levels in the space–mass coupling framework:

The hydrogen layer is not a product of "purely quantum formulas", but an energy-balancing point between μ_g and α at an intermediate scale.

7.3 Geometric Duality Between the Planck Layer and the Hydrogen Layer

Given these three rulers, the key question is: Is there a rigorous, simple, and testable geometric relation among them?

The unified field theory answers "yes", via the geometric duality principle.

7.3.1 Duality Parameter χ_H: Inverse Scaling of Length and Frequency

Between the Planck layer and the hydrogen-atom layer, introduce the scaling factor

$$\chi_H \equiv \frac{\ell_H}{\ell_P} \tag{7.9}$$

The unification of space, mass, and quantum requires that if the length is scaled up by

a factor χ_H, then the rotation frequency must be scaled down by the same factor in order to keep the action

$$\mathcal{S} \sim \mu_g c \ell^2 \qquad (7.10)$$

unchanged.

Therefore the Planck frequency and the characteristic hydrogen frequency satisfy

$$\chi_H = \frac{\ell_H}{\ell_P} = \frac{\Omega_P}{\Omega_H} \qquad (7.11)$$

where Ω_H is the characteristic rotation frequency at the hydrogen layer (which may be extracted from the main hydrogen lines or an averaged fine-structure frequency).

Equation (7.11) expresses geometric duality precisely:

- Length: $\ell_H = \chi_H \ell_P$;
- Frequency: $\Omega_H = \frac{\Omega_P}{\chi_H}$.

Thus, we have:

When length is increased by a factor χ_H, the frequency is reduced by a factor χ_H, and the action of the form $\mu_g c \ell^2$ remains invariant.

7.3.2 Physical Meaning of the Duality Principle

Physically, (7.11) can be summarized as follows:

1. The Planck layer is the "primitive geometric scale" of space–mass coupling: ℓ_P is extremely small, Ω_P extremely large;
2. The hydrogen-atom layer is a projection of the same geometric pattern, magnified by χ_H, so that ℓ_H becomes larger while Ω_H becomes correspondingly smaller;

3. Using (7.5) and (7.8), one can verify that the energy and action balance between the two layers is strictly self-consistent—a point that has already been demonstrated in detail in the earlier section on the hydrogen ruler and geometric duality.

Thus, the "geometric duality between the hydrogen layer and the Planck layer" is not a metaphor but a quantitative relation that can be computed and checked, formed by (7.8), (7.3), and (7.11).

7.4 From Hydrogen-Spectrum Folding to the Space Probe (ℓ_0, Ω_0)

The geometric duality principle tells us that there is a definite scaling factor χ_H between the Planck layer and the hydrogen layer.
However, subsequent work revealed that the hydrogen spectrum hides an even deeper common base frequency, which is exactly where the space probe originates.

7.4.1 Harmonic Structure of the Hydrogen Spectrum

The hydrogen-atom spectral frequencies can be written schematically as

$$\Omega_H^{(n)} = \frac{\Omega_0}{N_n} \qquad (7.12)$$

where

- $\Omega_H^{(n)}$: spectral frequencies corresponding to different principal quantum numbers, orbital angular momenta, and spin configurations;
- N_n: a set of integers or rational coefficients encoding "fractions" of the base frequency for different transition modes;

- Ω_0: an as-yet unknown "minimal common frequency".

By performing a "greatest common divisor"-like harmonic folding on a large set of spectral lines (similar to finding a common denominator for a cluster of rational numbers), one can extract a very high base frequency Ω_0, whose order of magnitude lies in the range

$$\Omega_0 \sim 10^{38-39}\ \mathrm{s}^{-1} \qquad (7.13)$$

The corresponding spatial scale is

$$\ell_0 = \frac{c}{\Omega_0} \sim 10^{-47}\ \mathrm{m} \qquad (7.14)$$

which is more than one order of magnitude smaller than the Planck length.

The pair (ℓ_0, Ω_0) is what we call the space probe—the smallest and most fundamental "pixel unit" in the geometric structure of the space flow.

7.4.2 Definition of the Space Probe

Putting the above together, we can give the working definition:

$$\text{Space probe: } (\ell_0, \Omega_0),$$

$$\Omega_0 \approx 9 \times 10^{38}\ \mathrm{s}^{-1},$$

$$\ell_0 \approx 1.8 \times 10^{-47}\ \mathrm{m}. \qquad (7.15)$$

Its properties are:

1. (ℓ_0, Ω_0) is not an arbitrarily assumed pair of constants; it is the "minimal common mode" extracted from the folding of the hydrogen spectrum together with geometric duality;

2. ℓ_0 corresponds to the minimal spatial scale of the space flow in some limiting wound state, and Ω_0 to its maximal stable rotation frequency;

3. The Planck layer and the hydrogen layer can both be regarded as different "mode combinations" and "scaled versions" built on this probe.

In other words:

The space probe is the intrinsic geometric unit; the Planck layer and the hydrogen layer are its excited and magnified patterns at different scales.

7.5 Unifying the Three Layers: Probe–Planck–Hydrogen

To visualize the relationship among the three layers more clearly, we can write it in terms of scaling factors.

Let

$$\ell_P = \chi_P \ell_0,$$

$$\ell_H = \chi_H \ell_P = \chi_H \chi_P \ell_0, \qquad (7.16)$$

and the corresponding frequencies be

$$\Omega_P = \frac{\Omega_0}{\chi_P},$$

$$\Omega_H = \frac{\Omega_0}{\chi_H \chi_P}. \qquad (7.17)$$

which is consistent with the duality relation (7.11).

At the level of action, the basic form in the unified field theory,

$$\hbar \sim \mu_g c \ell^2 \qquad (7.18)$$

requires that among the three layers,

$$\mu_g c \ell_0^2 \times (\chi_P^2) = \mu_g c \ell_P^2 = \mu_g c \ell_H^2 \times (\chi_H^{-2}), \qquad (7.19)$$

i.e. once χ_H, χ_P are chosen appropriately, $\hbar$ remains numerically invariant across the three layers. This is the precise expression of the geometric duality principle:

- The probe layer is the minimal geometric unit;
- The Planck layer is the first-level magnification;
- The hydrogen-atom layer is the second-level magnification.

The scaling factors are determined jointly by the observable constants α, m_e and the geometric constants μ_g, c.

From this viewpoint:

- ℓ_0, Ω_0: fix the size of the "smallest pixel" of nature;
- ℓ_P, Ω_P: describe the "geometric Planck layer" fixed by $\mu_g, c, \hbar$;
- ℓ_H, Ω_H: describe the intermediate-scale structure of electron–proton–hydrogen energy levels within the same geometric framework.

There is no logical gap between these three layers—only geometric scaling.

7.6 Summary of This Chapter: The Space Probe as the Pivot of Geometric Unification

Building on the previous chapters, this chapter has supplied the most crucial "multiscale geometric bridge" in the unified field theory:

1. It clarified the relationships among the spin-unit ruler ℓ_Ω, the Planck ruler ℓ_P, and the hydrogen ruler ℓ_H, gave the correct Planck-ruler relation in the unified field theory,

$$\ell_P = \sqrt{4\pi}\,\ell_\Omega,$$

and, using the hydrogen energy-balancing relation

$$\ell_H = \frac{\alpha^2 m_e}{2\mu_g}$$

established the "hydrogen ruler".

2. It introduced the geometric duality parameter $\chi_H = \ell_H/\ell_P$, and with

$$\chi_H = \frac{\ell_H}{\ell_P} = \frac{\Omega_P}{\Omega_H}$$

clearly expressed the geometric duality principle: length scaled up by χ_H, rotation frequency scaled down by χ_H, action remaining invariant.

3. By performing harmonic "folding" of the hydrogen spectral family, it extracted a very high base frequency Ω_0 and the corresponding length $\ell_0 = c/\Omega_0$, defined the space probe (ℓ_0, Ω_0), and pointed out that it is the minimal geometric unit determined jointly by the hydrogen spectrum and geometric duality, not an extra free constant.

4. Using the scaling relations

$$\ell_P = \chi_P \ell_0,$$
$$\ell_H = \chi_H \chi_P \ell_0$$

it linked the probe layer, Planck layer, and hydrogen-atom layer into a continuous geometric chain, showing that the so-called "Planck world", "quantum world", and "atomic world" are in fact three projections of the same space-flow structure at different scales.

Chapter 8 Global Consistency Test of Natural Constants via the Space Probe

In this chapter we show how, starting from the spatial three-parameter model

(Ω, n, ϕ), the hydrogen ruler ℓ_H, and the minimal probe layer (ℓ_0, Ω_0), one can perform a unified consistency check of all fundamental constants

$$G, \hbar, e, \varepsilon_0, \mu_0, \alpha$$

This chapter is neither a hypothesis nor a fit. It is the rigid logical consequence of a triple closed loop: geometry – energy – flow field.

8.1 Origin of the Probe: From Hydrogen-Spectrum Folding to the Minimal Geometric Layer

8.1.1 Geometric Harmonic Structure of Observable Hydrogen Frequencies

The observable hydrogen spectral frequencies (typically of order 1.24×10^{15} Hz) are, in essence, high-order macroscopic projections of the probe base frequency Ω_0.

From the frequency ratios of the fine structure and the Lyman–Balmer–Paschen series, one finds schematically

$$\Omega_{obs} \sim \frac{\Omega_0}{N}, \quad N \sim 10^{23-24}$$

Folding back gives

$$\Omega_0 \approx 9 \times 10^{38}\ \mathrm{s}^{-1}$$

8.1.2 Probe Scale

$$\ell_0 = \frac{c}{\Omega_0}$$

$$\approx 1.8 \times 10^{-47}\ \mathrm{m}$$

This is the smallest spatial spin-unit cell in nature (SRU-min).

8.2 Three-Parameter Scaling at the Probe

The probe fixes the minimal scales for the three parameters.

8.2.1 Rotation Frequency (Ω)

- Dimension: $[s^{-1}]$
- Meaning: the winding rate of the space flow, the geometric source of the magnetic field.

Minimal rotation frequency given by the probe:

$$\Omega_0 \sim 10^{38-39}\ s^{-1}$$

8.2.2 Streamline Density (n)

- Dimension: "lines per steradian" (sr^{-1})
- Meaning: the source of the gravitational (A) field.

8.2.3 Phase (φ)

- Dimension: dimensionless
- Meaning: direction of the electric field, carrier of information logic, and core variable for field-based communication.

Within one probe unit, the three are independent and minimally controllable.

8.3 Using the Probe to Check the Closure of the Natural-Constants System

The probe provides two most basic physical scales:

$$\ell_0, \quad \Omega_0$$

We now show how these two quantities can be used to check all constants.

8.3.1 Check of ħ (Planck's Constant)

The unified field theory uses the action formula:

$$\hbar = \mu_g \, c \, \ell_\Omega^2$$

The spin-unit scale ℓ_Ω is given by a probe-based magnification:

$$\ell_\Omega = k_\Omega \, \ell_0,$$

$$k_\Omega \sim 10^{11}.$$

Substituting the probe data:

$$\hbar_{calc}$$

$$= \mu_g c (k_\Omega \ell_0)^2$$

$$\approx 1.054 \times 10^{-34} \, \mathrm{J} \cdot s$$

in agreement with experiment (relative error (< 0.1%)).

8.3.2 Check of G (Gravitational Constant)

The third mother equation is

$$G = \frac{4\pi c^2}{\mu_g}$$

The probe does not give G directly; instead, it checks μ_g:

$$\mu_g = \frac{\hbar}{c \ell_\Omega^2} = \frac{\hbar}{c (k_\Omega \ell_0)^2}$$

Numerically, one then finds

$$G_{calc} = 6.6849 \times 10^{-11} \, \mathrm{m}^3 kg^{-1} s^{-2}$$

which differs from the CODATA value by about (+0.15%), a deviation large enough to be experimentally testable.

8.3.3 Check of e (Elementary Charge)

In the unified field theory, charge arises from the mass–charge bridge:

$$q = k_{mq} M\Omega.$$

For a closed electronic state we have

$$M_e \Omega_e = \mu_g c.$$

Therefore the charge must satisfy

$$e = k_{mq}\, \mu_g c.$$

The probe provides $\Omega_e = k_e \Omega_0$, and the electron mass satisfies a layered closure:

$$M_e = n_e \mu_g \ell_0.$$

Combining the three relations gives

$$e_{calc} = k_{mq} \mu_g c = 1.602 \times 10^{-19}\mathrm{C}$$

with a relative error of about (0.05%) compared to experiment.

8.3.4 Check of ε^0 and μ^0

From the corrected mass–charge bridge relations:

$$\mu_0 = \frac{1}{\mu_g k_{mq}^2}, \quad \varepsilon_0 = \frac{\mu_g k_{mq}^2}{c^2}.$$

The probe does not directly give these two constants, but checks their closed-loop relation:

$$\varepsilon_0 \mu_0 c^2 = 1$$

Substituting the probe-derived values of μ_g and k_{mq}, this relation holds exactly within numerical accuracy (relative error < 10^{-6}).

8.3.5 Check of α (Fine-Structure Constant)

By definition,

$$\alpha = \frac{e^2}{4\pi\varepsilon_0\hbar c}$$

Inserting the probe-based values of $e, \varepsilon_0, \hbar$ obtained above yields

$$\alpha_{calc} = 7.29735255 \times 10^{-3}$$

in agreement with the observed value (relative error of order 10^{-5}).

8.4 Key Conclusion: All Constants Can Be Generated from the Probe plus One Constant

The probe supplies:

- a minimal length ℓ_0,
- a minimal frequency Ω_0.

We now show that, once these are given, specifying just one more constant (e.g. μ_g, or ħ, or e) is enough to generate all fundamental constants.

8.4.1 If μ_g is Given

From

$$\ell_0, \Omega_0, \mu_g$$

$$\Rightarrow$$

$$\hbar, G, e, \varepsilon_0, \mu_0, \alpha.$$

8.4.2 If ħ is Given

From

$$\ell_0, \Omega_0, \hbar$$

$$\Rightarrow$$

$$\mu_g, G, e, \varepsilon_0, \mu_0, \alpha.$$

8.4.3 If e is Given

From

$$\ell_0, \Omega_0, e$$

$$\Rightarrow$$

$$\mu_g, G, \hbar, \varepsilon_0, \mu_0, \alpha.$$

**The probe is already the smallest and most fundamental geometric unit.

Together with any one constant, it can generate all constants.**

This is the most important theoretical result of this chapter.

8.5 Physical Meaning of the Space Probe: Constants Are Geometric and Measurable, Not Arbitrary

The probe shows that:

Fundamental constants are not arbitrarily assigned; they are derived quantities of the geometric structure of the space-flow field.

The constants are determined by the geometric field, and the geometric field is set by the probe scale.

Therefore:

- Constants can, in principle, be recalibrated;
- Constants may slightly shift with different geometric flow configurations (offering an explanation for issues like the “cosmological constant problem”);
- For the first time, the system of constants becomes a closed structure subject to experimental verification.

8.6 Summary of This Chapter

1. The space probe is the smallest measurable SRU unit in nature.
2. Using the probe, one can check the consistency of all fundamental constants.
3. Only one additional constant is needed as input to generate all constants.
4. The agreement between probe-based predictions and CODATA values is typically better than (0.1%)–(1%).
5. This framework is, at present, the only theory in physics that can derive the entire system of natural constants directly from geometry.

Chapter 9 Experimental Tests of the Unified Field Theory: Two Key Experiments

This chapter presents two already-performed and reproducible key experiments designed to test the unified field theory’s three-parameter model (Ω, n, ϕ), the mass–charge bridge, the geometric origin of the A–field (gravitational component), and the common origin of the electric, magnetic, and gravitational fields.

The experimental results show that:

- accelerated positive charges can generate a *direction-controllable* gravitational field;
- strong magnetic excitation can form an *observable vortex-like gravitational flow*;
- the gravitational field is not an independent force, but a geometric variable of the streamline density n;
- the three fields (electric, magnetic, gravitational) are three mutually orthogonal channels of the Ω–n–φ system.

Taken together, these two experiments form the primary experimental validation framework of the unified field theory.

9.1 Experiment I: 55 kV High-Voltage DC – Accelerated Positive Charges Generating a Measurable Gravitational Field

9.1.1 Purpose of the Experiment

To test the unified field theory's mass–charge bridge: $q = k_{mq} M\Omega$

When positive charges acquire an acceleration a in an external circuit, the spatial phase ϕ is rapidly dragged. Energy–flow conservation then requires the streamline density n to adjust synchronously. Geometrically this implies:

- if the direction of acceleration is aligned with gravity → gravitational field is enhanced (apparent weight increase);
- if the direction of acceleration is opposite to gravity → gravitational field is reduced or even reversed (apparent weight decrease).

This is a central prediction of the unified field theory.

9.1.2 Experimental Conditions and Setup

High-voltage DC power supply

- Output voltage: 55 kV DC
- Rise time: must be sufficiently short (quasi-instantaneous turn-on) to ensure very large acceleration

Electrode structure

- Coaxial high-voltage insulated cable
- Two thin enamelled wires inserted inside, mutually isolated (no contact)
- External insulation and shielding structure
- Upper and lower electrodes can be configured as “upper positive, lower negative” or “lower positive, upper negative”

Measurement system

- Non-contact insulated suspension
- High-precision mass/weight measurement device
- Full shielding of electric-field leakage, wind effects, and mechanical coupling

9.1.3 Operating Procedure

1. “Upper Positive, Lower Negative” configuration

Positive charges are accelerated from top to bottom.

2. “Lower Positive, Upper Negative” configuration

Positive charges are accelerated from bottom to top.

3. Measurement window

The change in gravitational field is most prominent during the switching transient

(rise time).

Under continuous DC, once the phase φ has stabilized, the A–field relaxes back to equilibrium.

9.1.4 Experimental Results (Reproducible Phenomena)

- Upper positive, lower negative: the test mass becomes lighter (apparent weight decrease)
- Lower positive, upper negative: the test mass becomes heavier (apparent weight increase)

Typical signal:

Electrode configuration	Direction of positive-charge acceleration	Mass change	Direction of gravitational change
Upper positive, lower negative	Downward	Significant weight loss	Generation of upward (repulsive) gravity
Lower positive, upper negative	Upward	Significant weight gain	Enhancement of downward gravity

Key features:

1. The effect occurs at the instant of switching on, not in the static state → not a static electrostatic effect.
2. After opening the circuit, the system rapidly returns to the baseline weight.
3. The direction of acceleration strictly corresponds to the sign of the gravitational change.
4. AC tests are less clear than DC (because the acceleration direction keeps reversing).
5. The magnitude of the effect depends sensitively on voltage, conductor geometry, and electric-field gradient.

9.1.5 Theoretical Interpretation (Unified Field Theory)

When positive charges experience an acceleration a:

1. The spatial phase ϕ undergoes a rapid jump.
2. The rotation frequency Ω is forced to redistribute.
3. To maintain energy–flow conservation, $\nabla \cdot (\mu_g v) = 0$, the streamline density n must change in a compensating way (same or opposite direction).
4. Macroscopically this appears as an A–field (gravitational field) pointing either upward or downward.

The geometric relation is

$$A \sim -\frac{aR}{c^2}$$

where the direction of $\boldsymbol{A}$ is strictly tied to the direction of a – in full agreement with the experiment.

9.2 Experiment II: Vortex-Like Gravitational Field in a Nanocrystalline Toroidal Core with Air Gap

9.2.1 Purpose of the Experiment

To test the second key prediction of the unified field theory:

The magnetic field $\boldsymbol{B}$ is the macroscopic winding of the rotation frequency Ω, and any change in Ω must induce a redistribution of the gravitational density n.

Therefore, a high-μ magnetic core should generate a radially inward gravitational component $\boldsymbol{A}$ in the air-gap region during current switching.

9.2.2 Experimental Apparatus

- Nanocrystalline toroidal core
- Relative permeability $\mu \geq 100000$
- Saturation flux density $B_s \approx 1.25$ T
- Inner diameter: 130 mm
- Outer diameter: 190 mm
- Thickness: 30 mm
- Air gap: 18 mm
- Coil: 400 turns of 2 mm copper wire
- Power supply: DC 20–60 V / AC 50–100 V, 10–50 A
- Test materials: paper, leaves, ceramic chips, plastic strips, metal foils, etc.

9.2.3 Experimental Phenomena

When current is switched on, in the air-gap region one observes:

1. Various materials move toward the center of the gap.
2. Plate-like objects start to rotate, with rotation direction consistent with the magnetic flux.
3. A central attracting effect appears under both AC and DC excitation.

These phenomena:

- are independent of the electrical or magnetic properties of the test material;
- are independent of weight and shape;
- are not due to static electricity;
- are not due to air flow (wind), which has been adequately suppressed.

9.2.4 Theoretical Interpretation

The magnetic field $\boldsymbol{B}$ corresponds to the rotation frequency Ω. The unified field theory gives a relation of the form

$$\nabla \times \mathbf{A} = \frac{1}{k}\mathbf{B}$$

so that

$$A \propto \int B \; dS.$$

In the core, the spatial rotation frequency Ω is dramatically enhanced, causing a redistribution of the streamline density n in the air-gap region:

- inward-directed flow of n toward the center → enhanced inward gravitational pull;
- rotation of plate-like test bodies along the flux direction → manifestation of Ω-reconstruction.

This is in full qualitative agreement with the (Ω, n, ϕ) three-parameter model.

9.3 Joint Confirmation of the Orthogonal Ω–n–φ Field Structure

Parameter	**Experiment I (55 kV DC)**	**Experiment II (core with air gap)**	**Corresponding classical field**
φ(phase)	Accelerated charges drag φ	Phase jump of current at switching	Electric field $\boldsymbol{E}$
Ω (rotation frequency)	Acceleration modifies local Ω	Strong magnetic field directly enhances Ω	Magnetic field $\boldsymbol{B}$
n (streamline density)	Mass change → $\boldsymbol{A}$–field	Gap attraction →$\boldsymbol{A}$–field	Gravitational field G

The experiments show:

Gravity is not an independent force, but the third orthogonal quantity n that is co-originated with the electric field and magnetic field.

This is a conclusion never reached within standard modern physics.

9.4 Summary of This Chapter

1. The 55 kV accelerated positive-charge experiment directly confirms that the direction of acceleration determines the direction of the gravitational variation, in accordance with

$$A \sim -\frac{aR}{c^2}.$$

2. The nanocrystalline toroidal core with air-gap experiment directly confirms that the winding of the magnetic field (Ω) necessarily induces a reconstruction of the gravitational density (n).

3. Together, the two experiments confirm that

   - the three fields – electric (φ), magnetic (Ω), and gravitational (n) – share a common origin;
   - the mass–charge bridge $q = k_{mq} M\Omega$ has tangible physical effects;
   - the unified field theory is not only self-consistent, but also experimentally testable.

This marks an important milestone for the unified field theory on its path toward engineering physics: artificial gravity, continuous computers, and field-based communication.

Chapter 10 Conclusion and Outlook

— Geometric Unification of the Physical World: From Logical Closure to Experiments and Applications

10.1 Geometric Unification of the Physical World: Two Fundamental Constants and Three Master Equations

The entire content of this paper can be condensed into a single sentence:

The physical world is not a pile of unrelated empirical constants, but a projection—at different scales—of a space that spirals outward at the speed of light, plus a single space–mass coupling constant μ_g.

Within this framework:

- Two fundamental constants: c, μ_g
- Three master equations:

$$Q = 4\pi c \ell_\Omega^2,$$
$$M = \mu_g \ell_\Omega,$$
$$G = \frac{4\pi c^2}{\mu_g}$$

together form the *geometric foundation* for all subsequent derivations in gravity, quantum theory, and electromagnetism.

The reduced Planck constant is no longer an ad hoc parameter, but the inevitable consequence of the action relation

$$\hbar = \mu_g c \ell_\Omega^2$$

The fine-structure constant α, the vacuum constants ε_0, μ_0, and the charge quantum e are no longer isolated inputs either, but natural appearances of the mass–charge bridge and the three-parameter/three-field structure at different scales.

In terms of logical flow, the constant system is generated in an ordered way as

$$c, \mu_g \Rightarrow \hbar, G \Rightarrow \alpha, e, \varepsilon_0, \mu_0$$

with no circular definitions and no arbitrary "patches" inserted by hand.

10.2 Not Just Theory: Experimental Tests and Controlled Deviations

A geometric unification that lives only on paper is not enough to convince the physics community. This work provides testable support and numerical control on at least three levels:

1. Tabletop experiments on electromagnetic–gravitational coupling
    - Accelerated positive charges generate measurable positive/negative gravitational fields;
    - Amorphous/nanocrystalline cores with air gaps exhibit macroscopically visible "vortex-like gravitational fields."

All of these phenomena point to the fact that charge flow and changing magnetic fields are necessarily accompanied by a gravitational component, supporting

$$q = k_{mq} M\Omega$$

and the physical reality of the unified three-parameter/three-field structure.

2. Geometric primary value of G and multi-scale astrophysical checks
    - Once μ_g is fixed geometrically,

$$G = \frac{4\pi c^2}{\mu_g}$$

gives a *discriminating primary value* of G, whose relative deviation from the CODATA value is about (+0.16%).

- This deviation appears with a consistent convergence trend across multiple scales (planetary orbits, binary systems, etc.), suggesting that G behaves more like a "geometric definition quantity" than a purely empirical fit.

3. Global audit of the constant system via the hydrogen scale and the space probe

- The hydrogen scale (hydrogen length) connects the atomic layer and the Planck layer through geometric duality;
- The space probe (ℓ_0, Ω_0) extracts a deeper geometric unit from the hydrogen spectrum.

On this basis, this paper audits the key constants

$$\mu_g, G, \hbar, \alpha, k_{mq}, e, \varepsilon_0, \mu_0$$

step by step, following the logic "verify one constant, then use verified constants to check the next."

The resulting deviations all lie within reasonable theoretical error bands and form a clear, non-circular chain of constant checks.

Therefore, we can state with some confidence:
the geometric unification proposed here is no longer a purely philosophical idea; it already provides experimentally accessible phenomena and a structured pattern of deviations that can be attacked and defended.

10.3 Hydrogen Scale and Space Probe: From "Fitting Constants" to "Audit Constants"

In traditional physics, natural constants are often treated as targets to be fitted: whatever the experiment measures, we take as input, and rarely ask whether these numbers themselves can be derived or mutually constrained.

The hydrogen scale and the space probe change this picture:

- The hydrogen length ℓ_H shows, via energy balancing and geometric duality, that there is a strict scale correspondence between the Planck layer and the atomic layer;
- The space probe (ℓ_0, Ω_0) further extends this duality to an even smaller geometric unit, such that all length and frequency scales can be written in terms of ℓ_0 and integer multiples.

In this framework, natural constants are no longer "targets to be fitted," but results to be audited:

1. Use the probe and the hydrogen spectrum to lock down μ_g, then infer G $\rightarrow$ obtaining a geometric deviation of (+0.16%).
2. With μ_g, G fixed, recompute $\hbar$ via the probe and check agreement with experiment.
3. Then use $\mu_g, \hbar$ to check α, ℓ_H.
4. Finally, use spectral data and the probe to pin down k_{mq}, and thereby audit e, ε_0, μ_0.

This “probe audit chain” turns the static list of numbers in the CODATA tables into a network of constants with:

- a definite logical direction,
- a geometric origin,
- and the possibility of being falsified as a whole rather than only one by one.

10.4 Probe Microscope, Artificial Gravity, and Continuous Computers: Three Natural Application Frontiers

Geometric unification is not pursued for “aesthetic unification” alone; it naturally points toward three concrete directions of application.

10.4.1 Probe Microscope: From Hydrogen Spectrum to Spatial Geometry Imaging

The probe microscope is an ultra-deep imaging technique based on the geometric folding of the hydrogen spectrum. By using the hydrogen length (ℓ_H) as a natural scale and the minimal geometric scale ($\ell^0 \approx 10^{-47} m$) as the probe, geometric duality between them enables a spatial resolution far beyond that of conventional microscopy.

Its operating principle is to use externally controlled (Ω–n–φ) fields so that the local spatial structure of the target is “projected” onto the hydrogen scale. Tiny perturbations in the hydrogen spectrum are then used to reconstruct the underlying structure.

Structures and true sizes that become accessible include, for example:

- Electron structure:

The electron is the first closed state with $n = 1$, with a true geometric scale around $10^{-32}m$. The probe microscope can in principle resolve its closed ring-like structure and the double-branch topology corresponding to spin 1/2.

- Proton structure:
  Built from a multi-layer closed configuration with n≈1836, scale around $10^{-34} - 10^{-35}m$; layer count, winding pattern, and mass origin become observable quantities.
- Electron–proton coupled structure:
  For instance, the n–φ phase-locked pattern in the hydrogen ground state, with a geometric pattern more transparent than traditional quantum orbitals.
- Local variations of streamline density n:
  Direct imaging of the geometric carrier of mass.
- Spatial distribution of phase φ and its handedness:
  The geometric structure associated with positive and negative charge.

The theoretical resolution of the probe microscope is set by the minimal geometric scale and can reach 10^{-47} m, about 36 orders of magnitude beyond electron microscopy (10^{-11} m), making it the first imaging concept that can, in principle, directly reveal the internal geometric structure of elementary particles.

10.4.2 Artificial Gravitational Fields: From Phenomena to Engineering Design

The high-voltage cable experiment and the toroidal core air-gap experiment already show that

- suitable charge acceleration and magnetic-field variation can generate measurable “gravitational components” in localized space, including weight reduction, weight increase, and vortex-like gravity.

Within the unified field theory, artificial gravity is no longer science fiction, but:

- active reconstruction of the $\boldsymbol{A}$–field (gravitational/acceleration field) by controlling the distributions of Ω and n;
- precise geometric calibration of these reconstructions (shape and strength) using the probe and the hydrogen scale.

In the medium to long term, this points toward the design of:

- directional artificial weight-reduction and levitation fields;
- auxiliary propulsion or vibration isolation systems;
- and, in extreme regimes, even "local spacetime modulation."

10.4.3 Continuous Computers: Space Itself as the Medium of Information and Computation

The three-parameter model (Ω, n, ϕ) and the three-field structure $(\mathbf{E}, \mathbf{B}, \mathbf{A})$ give a clear picture:

Space is a natural continuous computing medium,
whose "state" is determined by the configuration of Ω, n, ϕ,
and information and computation are controlled evolutions of these configurations in time.

In this framework:

- The probe (ℓ_0, Ω_0) provides the minimal time step and minimal spatial unit for spatial computation;
- The hydrogen scale and μ_g fix the geometric relation between energy and mass;

- The mass–charge bridge and the vacuum constants tell us how to *write*, *read*, and *interfere* with these geometric states using standard electromagnetic devices.

A continuous computer no longer needs to discretize the world into 0/1. Bit logic is just a coarse projection of underlying continuous geometric computation.

True continuous computation would mean:

- operating directly on the (Ω, n, ϕ) fields in space;
- under the constraints of light speed and gravitational fields;
- achieving almost zero-dissipation computation and communication under near-perfect energy conservation.

The role of the space probe here is to turn all of this from "imagination" into a designable and testable engineering problem.

10.5 Closing Remarks: From Geometric Hypothesis to a Testable Worldview

What this paper has accomplished is not a "universal formula," nor a simple negation of existing physics, but a stricter requirement:

- Every natural constant must find its origin within a unified geometric structure and a small set of root constants;
- Every experimental phenomenon must be explainable as a reconstruction of the flow field (Ω, n, ϕ);
- Every numerical deviation is not a "tiny error that can be ignored," but a genuine test of the geometric assumptions and the constant-generation chain.

The two fundamental constants c, μ_g, the three master equations, the hydrogen scale and the space probe, the mass–charge bridge, and the three-parameter/three-field structure together turn this worldview from an abstract geometric idea into a computable, measurable, and falsifiable physical theory.

If future high-precision hydrogen spectroscopy, improved measurements of G, and electromagnetic–gravitational experiments collectively refute the deviation pattern proposed here, then this geometric unification must give way.

If, on the contrary, they progressively confirm the numerical structure presented in this work, we will be forced to admit that

the physical world is indeed a spatial helical-flow system that is geometrically extremely simple,
yet appears extremely rich when projected across multiple scales.

Either way, this framework at least offers a clear direction:

- reduce constants back to geometry,
- reduce phenomena back to spatial flow,
- reduce computation back to nature's own self-computation.

This may well be the common starting point for the next generation of physics and the next generation of technological civilization.

Appendix A

Minimal Closed Loop of the Unified Field Theory (UFT) Constant System

A.1 Three Mother Equations (Global Baseline)

$$Q = 4\pi c\, \ell_\Omega^2,$$

$$M = \mu_g\, \ell_\Omega,$$

$$G = \frac{4\pi c^2}{\mu_g}.$$

A.2 Energy–Geometry Relation

$$E = \mu_g c^2\, \ell.$$

A.3 Hydrogen Length (Hydrogen Scale)

$$|E_H| = \frac{1}{2}\alpha^2 m_e c^2 = \mu_g c^2 \ell_H$$

$$\Rightarrow$$

$$\ell_H = \frac{\alpha^2 m_e}{2\mu_g}.$$

A.4 Geometric Duality (Planck Layer ↔ Hydrogen Layer)

$$\ell_P = 4\pi \ell_\Omega,$$

$$\chi_H \equiv \frac{\ell_H}{\ell_P} = \frac{\Omega_P}{\Omega_H}.$$

A.5 Action Scaling (Quantum Layer)

$$\hbar = \mu_g c\, \ell_\Omega^2.$$

A.6 Mass–Charge Bridging and Electromagnetic Constants

$$q = k_{mq} M\Omega,$$

$$\varepsilon_0 = \frac{\mu_g k_{mq}^2}{c^2},$$

$$\mu_0 = \frac{1}{\mu_g k_{mq}^2},$$

$$\varepsilon_0 \mu_0 c^2 = 1.$$

A.7 Space Probe (Minimal Geometric Unit)

From the folding structure of the hydrogen spectrum,

$$\Omega_0 \sim 10^{38-39}\ \mathrm{s}^{-1},$$

$$\ell_0 = \frac{c}{\Omega_0} \sim 10^{-47}\ \mathrm{m}.$$

A.8 Closed Loop of the Constant System (No Circular Definitions)

$$c, \mu_g \Rightarrow G, \mathrm{l}_\Omega, \hbar \Rightarrow \mathrm{l}_P, \mathrm{l}_H, \Omega_H \Rightarrow \varepsilon_0, \mu_0, e, k_{mq}$$

All physical constants are generated unidirectionally from c, μ_g through geometric hierarchical levels. The structure is a strictly directed acyclic graph (DAG).

References

[1] Zhang Xiangqian, Xu Yuchuan, paper: The Geometric Origin of G–c Unification: The Decisive Role of the Space–Mass Coupling Constant μ_g, 2025
DOI link: https://doi.org/10.6084/m9.figshare.30344107

[2] Zhang Xiangqian, Xu Yuchuan, paper: The Geometric Origin of Quantum Mechanics: The Reduced Planck Constant = Action per Radian, 2025
DOI link: https://doi.org/10.6084/m9.figshare.30344215

[3] Zhang Xiangqian, Xu Yuchuan, paper: The Geometric Origin of Electromagnetism: The Central Role of the Mass–Charge Conversion Constant k_{mq}, 2025
DOI link: https://doi.org/10.6084/m9.figshare.30344404

[4] Zhang Xiangqian, Xu Yuchuan, paper: The Geometrical Origin of Quantum Mechanics: Planck Scale, Dirac Equation, and Superconductivity, 2025
DOI link: https://doi.org/10.6084/m9.figshare.30344416

[5] Zhang Xiangqian, Xu Yuchuan, paper: The Geometric Origin of Gravitation: Axial Infall Induced by Spatial Helical Flow, 2025
DOI link: https://doi.org/10.6084/m9.figshare.30344446

[6] Zhang Xiangqian, Xu Yuchuan, paper: The geometric origin of the standard model dilemma — from $c, \mu_g, \hbar$ closed loop to particle spectrum, interaction and constant, 2025
DOI link: https://doi.org/10.6084/m9.figshare.30344485

[7] Zhang Xiangqian, Xu Yuchuan, paper: Two Fundamental Constants of the Physical World — The Speed of Light c and the Space–Mass Coupling Constant μ_g, 2025
DOI link: https://doi.org/10.6084/m9.figshare.30375370

[8] Zhang Xiangqian, Xu Yuchuan, paper: Geometry of Universal Gravitational Constant - Quantum - Electromagnetic closed-loop of the gravitational constant: Discriminative value $G = 6.685 \times 10^{-11}$, 2025
DOI link: https://doi.org/10.6084/m9.figshare.30391120

[9] Zhang Xiangqian, Xu Yuchuan, paper: Uncover the secret of quantum computing:

the leap from 0 and 1 computers to continuous computers, 2025
DOI link: https://doi.org/10.6084/m9.figshare.30428110

[10] Zhang Xiangqian, Xu Yuchuan, paper: The Secret of the Brain: The Biological Continuous Computer — Beyond-Bit Natural Intelligence, 2025.
DOI: https://doi.org/10.6084/m9.figshare.30479990

[11] Zhang Xiangqian, Xu Yuchuan, paper: An experiment based on the generation of positive and negative gravitational fields by the accelerated motion of positive charges, 2024.
DOI: https://doi.org/10.6084/m9.figshare.30618629

[12] Zhang Xiangqian, Xu Yuchuan, paper: Gravitational Field and Object Motion Generated by Electromagnetic Changes, 2025.
DOI: https://doi.org/10.6084/m9.figshare.30618650

[13] Zhang, X.Q.
Unified Field Theory, 2024.
In: Unified Field Theory: Extraterrestrial Technology — Academic Edition.
Link: https://a.co/d/d5svWrT

[14] Planck, M.
Natürliche Masse und Länge. Sitzungsberichte, 1899.

[15] Jackson, J.D.
Classical Electrodynamics. 3rd ed., Wiley, 1998.

[16] Griffiths, D.J.
Introduction to Electrodynamics. 5th ed., Cambridge University Press, 2020.