Copyright Notice

These slides are distributed under the Creative Commons License.

DeepLearning.ai makes these slides available for educational purposes. You may not use or distribute these slides for commercial purposes. You may make copies of these slides and use or distribute them for educational purposes as long as you cite DeepLearning.AI as the source of the slides.

For the rest of the details of the license, see https://creativecommons.org/licenses/by-sa/2.0/legalcode

DeepLearning.AI

Math for Machine Learning

Linear algebra - Week 4

W4 Lesson 1

DeepLearning.AI

Determinants and Eigenvectors

Machine learning motivation

Principal Component Analysis



Principal Component Analysis





Principal Component Analysis



Principal Component Analysis

- Reduce dimensions (columns) of dataset
- Preserve as much information as possible



species	culmen_length_mm	culmen_depth_mm	flipper_length_mm	body_mass_g
Adelie	40.6	17.2	187.0	3475.0
Adelie	38.9	17.8	181.0	3625.0
Adelie	35.7	16.9	185.0	3150.0
Gentoo	50.0	15.3	220.0	5550.0
Adelie	34.5	18.1	187.0	2900.0

PC1	PC2	species
1.353843	-0.422253	Adelie
1.760446	-0.350965	Adelie
2.005766	-1.113797	Adelie
-2.585758	0.061768	Gentoo
2.438111	-0.786227	Adelie

What to expect?



What to expect?

Linear transformation

1. Singular / Non-singular
2.
3.

Characterize your transformation

Non-singular

3	1
1	2

Singular

1	1
2	2



What to expect?

Linear transformation

1

2 Determinant as area

3

Characterize your transformation



What to expect?



$$\begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 \\ -2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ -3 & 3 \end{bmatrix}$$

Det = 5 Det = 3 Det = 15

$= 5 \cdot 3$

$$\begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix}^{-1} = \begin{bmatrix} 0.4 & -0.2 \\ -0.2 & 0.6 \end{bmatrix}$$

Det = 5 Det = 0.2 = $\frac{1}{5}$

What to expect?



What to expect?



What to expect?





What to expect?





1 element
Dimensions: 1



2 elements
Dimensions: 2

What to expect?

Linear transformation

1. Singular / Non-singular
2. Determinant as area
3. Properties of determinant

Characterize your transformation

1. Eigenvalues and Eigenvectors

PCA



2	1	1	=	2
0	3	0		0

(1,0) → (2,0)

$$A v_1 = \lambda_1 v_1$$

What to expect?

Linear transformation

1. Singular / Non-singular
2. Determinant as area
3. Properties of determinant

Characterize your transformation

1. Basis
2. Span
3. Eigenvalues and Eigenvectors

PCA

DeepLearning.AI

Determinants and Eigenvectors

Singularity and rank of linear transformations

Non-singular transformation



3	1
1	2



Singular transformation



1	1	1	=	2
2	2	1		4

(0,0) → (0,0)
(1,0) → (1,2)
(0,1) → (1,2)
(1,1) → (2,4)



Singular transformation



0	0	a	=	0
0	0	b		0



Singular and non-singular transformations

Non-singular

3	1
1	2

Singular

1	1
2	2

Singular

0	0
0	0



Rank of linear transformations



DeepLearning.AI

Determinants and Eigenvectors

Determinant as an area

Determinant as an area



3	1
1	2

$$\text{Det} = 3 \cdot 2 - 1 \cdot 1$$

$$\text{Det} = 5$$



Determinant as an area



1	1
2	2

$$\text{Det} = 1 \cdot 2 - 1 \cdot 2$$

$$\text{Det} = 0$$



Determinant as an area



0	0
0	0

$$\text{Det} = 0 \cdot 0 - 0 \cdot 0$$

$$\text{Det} = 0$$



Determinant as an area

Non-singular

3	1
1	2

Determinant = 5



Area = 5

Singular

1	1
2	2

Determinant = 0



Area = 0

Singular

0	0
0	0

Determinant = 0



Area = 0

Negative determinants?



Det = $3 \cdot 2 - 1 \cdot 1$

Det = 5



Det = $1 \cdot 1 - 3 \cdot 2$

Det = -5

Determinant as an area



1	3
2	1

$$\text{Det} = 1 \cdot 1 - 3 \cdot 2$$

$$\text{Det} = -5$$



Determinant as an area



1	3
2	1

$$\text{Det} = 1 \cdot 1 - 3 \cdot 2$$

$$\text{Det} = -5$$

Negative



DeepLearning.AI

Determinants and Eigenvectors

Determinant of a product

Determinant of a product



Determinant of a product

$$\det(AB) = \det(A)\det(B)$$

Determinant of a product



3	1
1	2

Det = 5



Area blows up by 5

Determinant of a product



1	1
-2	1

Det = 3



Area blows up by 3

Determinant of a product



Quiz

- The product of a singular and a non-singular matrix (in any order) is:
  - Singular
  - Non-singular
  - Could be either one

Solution

- If A is non-singular and B is singular, then det(AB) = det(A) x det(B) = 0, since det(B) = 0. Therefore det(AB) = 0, so AB is **singular**.

When one factor is zero

$$5 \cdot 0 = 0$$

When one factor is singular…



If one factor is singular…



3	1
1	2

Det = 5



1	2
1	2

Det = 0



Area blows up by 0

4	8
3	6

Det = 0

DeepLearning.AI

Determinants and Eigenvectors

Determinant of inverse

Quiz

- Find the determinants of the following matrices

0.4	-0.2
-0.2	0.6

0.25	-0.25
-0.125	0.625

Solution

Det $\begin{vmatrix} 0.4 & -0.2 \\ -0.2 & 0.6 \end{vmatrix}$ = (0.4)(0.6) - (-0.2)(-0.2) = 0.2

Det $\begin{vmatrix} 0.25 & -0.25 \\ -0.125 & 0.625 \end{vmatrix}$ = (0.25)(0.625) - (-0.125)(-0.25) = 0.125

Determinant of an inverse



det = 5 det = 0.2

$$5^{-1} = 0.2$$



det = 8 det = 0.125

$$8^{-1} = 0.125$$



det = 0 det = ???

$$0^{-1} = ???$$

Determinant of an inverse

$$\det(A^{-1}) = \frac{1}{\det(A)}$$

Why?

Why is this?

$$\det(A^{-1}) = \frac{1}{\det(A)}$$

$$\det(AB) = \det(A)\det(B)$$

$$\det(AA^{-1}) = \det(A)\det(A^{-1})$$

$$\det(I) = \det(A)\det(A^{-1})$$

1

$\frac{1}{\det(A)}$

Determinant of the identity matrix

$$\det \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = 1 \cdot 1 - 0 \cdot 0 = 1$$

$$\det(I) = 1$$

W4 Lesson 2

DeepLearning.AI

Determinants and Eigenvectors

Bases

Bases



3	1
1	2



Bases



3	1
1	2



Bases





Bases





Bases





Bases





What is not a basis?





DeepLearning.AI

Determinants and Eigenvectors

Span

Span



Span



Span



Span



Span



Is this a basis?



No

Is this a basis for something?





A basis is a minimal spanning set



Basis



Not a basis

A basis is a minimal spanning set



Basis



Not a basis

Number of elements in the basis is the dimension



Dimensions: 1
1 element in the basis



Dimensions: 2
2 elements in the basis

Linearly independent and linearly dependent vectors



Linearly independent

Linearly independent and linearly dependent vectors



Linearly independent



Linearly independent

Linearly independent and linearly dependent vectors



Linearly independent



Linearly independent

Linearly independent and linearly dependent vectors



Linearly independent



Linearly independent



Linearly dependent

Linearly independent and linearly dependent vectors



Linearly independent



Linearly independent

Linearly independent and linearly dependent vectors



Linearly independent



Linearly independent



Linearly dependent

Linearly independent and linearly dependent vectors



Linearly independent



Linearly independent



Linearly dependent

Linearly independent and linearly dependent vectors



Linearly independent



Linearly independent



Linearly dependent

Linearly independent and linearly dependent vectors



Linearly independent



Linearly independent



Linearly dependent

Let's see how to check for linear dependence



Linearly dependent

$\alpha \quad +\beta \quad =$

Let's see how to check for linear dependence



Linearly dependent

$$\alpha v_1 + \beta v_2 = v_3$$

Let's see how to check for linear dependence



Linearly dependent

$$\alpha v_1 + \beta v_2 = v_3$$

$$\alpha \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \beta \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} -5 \\ 3 \end{bmatrix}$$

1. $-\alpha + 2\beta = -5$
2. $\alpha + \beta = 3$

Let's see how to check for linear dependence



Linearly dependent

$$\alpha v_1 + \beta v_2 = v_3$$

$$\alpha \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \beta \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} -5 \\ 3 \end{bmatrix}$$

v_3 is a linear combination of v_1 and v_2

1 $-\alpha + 2\beta = -5$

2 $\alpha + \beta = 3$

1 + 2

$3\beta = -2 \rightarrow \beta = -\frac{2}{3}$

2

$\alpha - \frac{2}{3} = 3 \rightarrow \alpha = \frac{11}{3}$

Quiz

Are these vectors linearly independent?

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \quad \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \quad \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

Solution

Are these vectors linearly independent?

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \quad \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \quad \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

$$1 \qquad -1 \qquad =$$

Linearly dependent

Solution

Are these vectors linearly independent?

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \quad \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \quad \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

$$1 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \quad -1 \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

Linearly dependent

Solution

Are these vectors linearly independent?

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \quad \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \quad \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

Not a basis!

Linearly independent

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$$

Linearly independent

$$\begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

Linearly independent

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

Basis: a formal definition

A basis is a set of vectors that:

- Spans a vector space
- Is linearly independent

Not all sets of N vectors are a basis
for N-dimensional space



Spans a line
Linearly independent
Is a basis



Spans the plane
Linearly independent
Is a basis



Spans a line
Linearly dependent
Not a basis



Spans the plane
Linearly dependent
Not a basis

DeepLearning.AI

Determinants and Eigenvectors

Eigenbasis

Basis



2	1	0	=	1
0	3	1		3

(1,0) → (2,0)

(0,1) → (1,3)



Eigenbasis



2	1	1	=	3
0	3	1		3





Eigenbasis



2	1	3	=	8
0	3	2		6

(3,2) → (8,6)



DeepLearning.AI

Determinants and Eigenvectors

Eigenvalues and Eigenvectors

Eigenvalues and eigenvectors



$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 3 \\ 3 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 6 \end{bmatrix} \neq \lambda \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$



Eigenvalues and eigenvectors



Eigenvalues and eigenvectors



$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

form a basis

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 6 \end{bmatrix}$$



Eigenvalues and eigenvectors



$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ form a basis

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 6 \end{bmatrix}$$



Eigenvalues and eigenvectors



$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ form a basis

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} -3 & +2 \end{bmatrix}$$



Eigenvalues and eigenvectors



Eigenvalues and eigenvectors



$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

1. Calculate the inverse of the eigenbasis
2. Multiply it by your vector

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

form a basis

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = -3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 6 \end{bmatrix}$$



Eigenvalues and eigenvectors

- $Av = \lambda v$ for each eigenvector / eigenvalue
- Eigenvectors: direction of stretch
- Eigenvalues: how much stretch
- Eigenbasis: the set of a matrix's eigenvectors, can be arranged as a matrix with one eigenvector in each column
- Save work and characterize a transformation

DeepLearning.AI

Determinants and Eigenvectors

Calculating eigenvalues and eigenvectors

Finding eigenvalues





Finding eigenvalues



Finding eigenvalues





Finding eigenvalues



Finding eigenvalues

If λ is an eigenvalue:

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

For infinitely many (x,y)

$$\begin{bmatrix} 2-\lambda & 1 \\ 0 & 3-\lambda \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Has infinitely many solutions

$$\det \begin{bmatrix} 2-\lambda & 1 \\ 0 & 3-\lambda \end{bmatrix} = 0$$

Characteristic polynomial

$$(2-\lambda)(3-\lambda) - 1 \cdot 0 = 0$$

$$\lambda = 2$$
$$\lambda = 3$$

Finding eigenvectors

Eigenvalues: $\lambda = 2$, $\lambda = 3$

Solve the equations

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 2 \begin{bmatrix} x \\ y \end{bmatrix}$$

$$2x + y = 2x \qquad x = 1$$

$$0x + 3y = 2y \qquad y = 0$$

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 3 \begin{bmatrix} x \\ y \end{bmatrix}$$

$$2x + y = 3x \qquad x = 1$$

$$0x + 3y = 3y \qquad y = 1$$

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Quiz

- Find the eigenvalues and eigenvectors of this matrix:

9	4
4	3

Solution

- Eigenvalues: 11, 1
- Eigenvectors: (2,1), (-1,2)

9	4
4	3

- The characteristic polynomial is $\det \begin{vmatrix} 9-\lambda & 4 \\ 4 & 3-\lambda \end{vmatrix} = (9-\lambda)(3-\lambda) - 4 \cdot 4 = 0$
- Which factors as $\lambda^2 - 12\lambda + 11 = (\lambda - 11)(\lambda - 1)$
- The solutions are $\lambda = 11$, $\lambda = 1$

Finding eigenvalues

$A =$

2	1	-1
1	0	-3
-1	-3	0

$\lambda I =$

λ	0	0
0	λ	0
0	0	λ

Characteristic polynomial: $\det(A - \lambda I) = 0$

$\det$

2 - λ	1	-1
1	-λ	-3
-1	-3	-λ

$= 0$

$$(2-\lambda)\lambda^2 \; +3 \; +3 \; -9(2-\lambda) \; +\lambda \; +\lambda \quad = -\lambda^3 + 2\lambda^2 + 11\lambda - 12 = 0$$

$$-(\lambda + 3)(\lambda - 1)(\lambda - 4) = 0$$

Eigenvalues: $-3, 1, 4$

Finding eigenvalues

$$A = \begin{bmatrix} 2 & 1 & -1 \\ 1 & 0 & -3 \\ -1 & -3 & 0 \end{bmatrix}$$

Eigenvalues: $-3, 1, 4$

$$Av = \lambda v$$

$$\underbrace{\begin{bmatrix} 2 & 1 & -1 \\ 1 & 0 & -3 \\ -1 & -3 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_{\begin{bmatrix} 2x_1 + x_2 - x_3 \\ x_1 - 3x_3 \\ -x_1 - 3x_2 \end{bmatrix}} = 4 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 4x_1 \\ 4x_2 \\ 4x_3 \end{bmatrix}$$

Finding eigenvalues

$$A = \begin{bmatrix} 2 & 1 & -1 \\ 1 & 0 & -3 \\ -1 & -3 & 0 \end{bmatrix}$$

Eigenvalues: $-3, 1, 4$

$$Av = \lambda v$$

$$\begin{bmatrix} 2 & 1 & -1 \\ 1 & 0 & -3 \\ -1 & -3 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 4 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$\begin{bmatrix} 2x_1 + x_2 - x_3 \\ x_1 - 3x_3 \\ -x_1 - 3x_2 \end{bmatrix} = \begin{bmatrix} 4x_1 \\ 4x_2 \\ 4x_3 \end{bmatrix}$$

$$\begin{aligned} 2x_1 + x_2 - x_3 &= 4x_1 \\ x_1 - 3x_3 &= 4x_2 \\ -x_1 - 3x_2 &= 4x_3 \end{aligned}$$

$$\begin{aligned} R_1 \quad & -2x_1 + x_2 - x_3 = 0 \\ R_2 \quad & x_1 - 4x_2 - 3x_3 = 0 \\ R_3 \quad & -x_1 - 3x_2 - 4x_3 = 0 \end{aligned}$$

$$\begin{aligned} R_2 + R_3 \\ -7x_2 - 7x_3 = 0 \\ x_2 = -x_3 \end{aligned} \qquad \begin{aligned} 3R_1 + R_3 \\ -7x_1 - 7x_3 = 0 \\ x_1 = -x_3 \end{aligned}$$

$$\begin{aligned} x_1 &= k \\ x_2 &= k \\ x_3 &= -k \end{aligned}$$

infinite solutions of this form

$$\begin{aligned} x_1 &= 1 \\ x_2 &= 1 \\ x_3 &= -1 \end{aligned}$$

this works!

$$\begin{aligned} x_1 &= 2 \\ x_2 &= 2 \\ x_3 &= -2 \end{aligned}$$

so does this!

Eigenvector: $\begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$

Finding eigenvalues

$A =$

2	1	-1
1	0	-3
-1	-3	0

Eigenvalues $\lambda_1 = 4$ $\lambda_2 = 1$ $\lambda_3 = -3$

Eigenvectors

1	0	2
1	1	-1
-1	1	1

Note on dimensions

Eigenvalues → Determinant → Square Matrix

9	4
4	3

9	4	5
4	3	-2

DeepLearning.AI

Determinants and Eigenvectors

On the number of eigenvectors

Number of eigenvectors

$A =$

2	1	-1
1	0	-3
-1	-3	0

3 by 3 matrix → ? → 3 distinct eigenvalues → ? → 3 distinct eigenvectors

Eigenvalues

$\lambda_1 = 4 \quad \lambda_2 = 1 \quad \lambda_3 = -3$

Eigenvectors

1	0	2
1	1	−1
−1	1	1

Repeated eigenvalues - Example 1

$A =$

2	0	0
-1	4	-0.5
0	0	2

Characteristic polynomial = $\det(A - \lambda I)$ = det

2 - λ	0	0
1	4 - λ	0.5
0	0	2 - λ

$$(2-\lambda)^2(4-\lambda)+0+0-0-0-0=0$$

Eigenvalues: $4, \boxed{2,\ 2}$ Repeated eigenvalue

Repeated eigenvalues - Example 1

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{bmatrix}$$

Eigenvalue: 4

$$Av = 4v$$

$$\underbrace{\begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_{\begin{bmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 2x_3 \end{bmatrix}} = 4 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 4x_1 \\ 4x_2 \\ 4x_3 \end{bmatrix}$$



Repeated eigenvalues - Example 1

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{bmatrix}$$

Eigenvalue: 4

$$Av = 4v$$

$$\begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 4 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 4x_1 \\ 4x_2 \\ 4x_3 \end{bmatrix}$$

$$\begin{bmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 2x_3 \end{bmatrix}$$

$$\begin{aligned} 2x_1 &= 4x_1 \\ -x_1 + 4x_2 - 0.5x_3 &= 4x_2 \\ 2x_3 &= 4x_3 \end{aligned}$$

$$\begin{aligned} -2x_1 &= 0 \\ -x_1 - 0.5x_3 &= 0 \\ -2x_3 &= 0 \end{aligned}$$

$x_1 = 0$

$x_2 =$ any number

$x_3 = 0$

Eigenvector

$$\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

Repeated eigenvalues - Example 1

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{bmatrix}$$

Eigenvalue: 2

$$Av = 2v$$

$$\underbrace{\begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_{\begin{bmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 2x_3 \end{bmatrix}} = 2 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2x_1 \\ 2x_2 \\ 2x_3 \end{bmatrix}$$

Repeated eigenvalues - Example 1

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{bmatrix}$$

Eigenvalue: 2

$$Av = 2v$$

$$\begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 2 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2x_1 \\ 2x_2 \\ 2x_3 \end{bmatrix}$$

$$\begin{bmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 2x_3 \end{bmatrix}$$

$$\begin{aligned} 2x_1 &= 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 &= 2x_2 \\ 2x_3 &= 2x_3 \end{aligned}$$

$$\begin{aligned} 0 &= 0 \\ -x_1 + 2x_2 - 0.5x_3 &= 0 \\ 0 &= 0 \end{aligned}$$

$$x_1 = 2x_2 - 0.5x_3$$

$$\begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix} \quad \begin{aligned} x_1 &= 2 \\ x_2 &= 1 \\ x_3 &= 0 \end{aligned}$$

$$\begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \quad \begin{aligned} x_1 &= 1 \\ x_2 &= 1 \\ x_3 &= 2 \end{aligned}$$

Point in different directions

Different eigenvectors

Repeated eigenvalues - Example 1

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 0 & 0 & 2 \end{bmatrix}$$

Eigenvalues $\lambda_1 = 4$ $\lambda_2 = 2$ $\lambda_3 = 2$

Eigenvectors $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ $\begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$ $\begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$

Repeated eigenvalues - Example 2

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix}$$

Repeated eigenvalues - Example 2

$A =$

2	0	0
-1	4	-0.5
4	0	2

Characteristic polynomial = $\det(A - \lambda I)$ = det

2-λ	0	0
1	4-λ	0.5
-4	0	2-λ

$$(2-\lambda)^2(4-\lambda) + 0 \;+0\; -0\; -0 -0$$

Eigenvalues: $4, \boxed{2,\ 2}$ Repeated eigenvalue

Repeated eigenvalues - Example 2

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix}$$

Eigenvalue: 4

$$Av = 4v$$

$$\underbrace{\begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_{\begin{bmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 4x_1 + 2x_3 \end{bmatrix}} = 4 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 4x_1 \\ 4x_2 \\ 4x_3 \end{bmatrix}$$

Repeated eigenvalues - Example 2

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix}$$

Eigenvalue: 4

$$\begin{aligned} 2x_1 &= 4x_1 \\ -x_1 + 4x_2 - 0.5x_3 &= 4x_2 \\ 4x_1 + 2x_3 &= 4x_3 \end{aligned}$$

$$Av = 4v$$

$$\underbrace{\begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_{\begin{bmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 4x_1 + 2x_3 \end{bmatrix}} = 4 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 4x_1 \\ 4x_2 \\ 4x_3 \end{bmatrix}$$

$$\begin{aligned} -2x_1 &= 0 \\ -x_1 - 0.5x_3 &= 0 \\ 4x_1 - 2x_3 &= 0 \end{aligned}$$

$x_1 = 0$ $\quad x_3 = 0$ $\quad x_2 =$ any number

$$\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad \begin{aligned} x_1 &= 0 \\ x_2 &= 1 \\ x_3 &= 0 \end{aligned}$$

Same as before!

Repeated eigenvalues - Example 2

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix}$$

Eigenvalue: 2

$$Av = 2v$$

$$\underbrace{\begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_{\begin{bmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 4x_1 + 2x_3 \end{bmatrix}} = 2 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2x_1 \\ 2x_2 \\ 2x_3 \end{bmatrix}$$

Repeated eigenvalues - Example 2

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix}$$

Eigenvalue: 2

$$Av = 2v$$

$$\begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 2 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2x_1 \\ 2x_2 \\ 2x_3 \end{bmatrix}$$

$$\begin{bmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 4x_1 + 2x_3 \end{bmatrix}$$

$$\begin{aligned} 2x_1 &= 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 &= 2x_2 \\ 4x_1 + 2x_3 &= 2x_3 \end{aligned}$$

$$\begin{aligned} 0 &= 0 \\ -x_1 + 2x_2 - 0.5x_3 &= 0 \\ 4x_1 &= 0 \end{aligned}$$

$$\begin{bmatrix} 0 \\ k \\ 4k \end{bmatrix}$$

$x_1 = 0 \qquad x_3 = 4x_2$

$$\begin{bmatrix} 0 \\ 1 \\ 4 \end{bmatrix} \quad x_1 = 0,\ x_2 = 1,\ x_3 = 4 \qquad \begin{bmatrix} 0 \\ 0.5 \\ 2 \end{bmatrix} \quad x_1 = 0,\ x_2 = 0.5,\ x_3 = 2$$

On the same line

Same eigenvector

Repeated eigenvalues - Example 2

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix}$$

Eigenvalue: 2

$$\begin{aligned} 2x_1 &= 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 &= 2x_2 \\ 4x_1 + 2x_3 &= 2x_3 \end{aligned}$$

$$Av = 2v$$

$$\begin{aligned} 0 &= 0 \\ -x_1 + 2x_2 - 0.5x_3 &= 0 \\ 4x_1 &= 0 \end{aligned}$$

$$\begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 2 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2x_1 \\ 2x_2 \\ 2x_3 \end{bmatrix}$$

$$\begin{bmatrix} 2x_1 \\ -x_1 + 4x_2 - 0.5x_3 \\ 4x_1 + 2x_3 \end{bmatrix}$$

$x_1 = 0$ $\qquad$ $x_3 = 4x_2$

$$\begin{bmatrix} 0 \\ k \\ 4k \end{bmatrix}$$

Repeated eigenvalues - Example 2

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 4 & -0.5 \\ 4 & 0 & 2 \end{bmatrix}$$

Eigenvalues $\lambda_1 = 4$ $\lambda_2 = 2$ $\lambda_3 = 2$

Eigenvectors $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ $\begin{bmatrix} 0 \\ 1 \\ 4 \end{bmatrix}$

Can’t create an eigenbasis from this matrix

Summary

a	b
c	d

Eigenvalues λ_1, λ_2

If $\lambda_1 \neq \lambda_2$ → 2 eigenvectors (2 different directions)

If $\lambda_1 = \lambda_2$ → 1 eigenvector (1 direction)

If $\lambda_1 = \lambda_2$ → 2 eigenvectors (2 different directions)

a	b	c
d	e	f
g	h	i

$\lambda_1, \lambda_2, \lambda_3$

If $\lambda_1 \neq \lambda_2 \neq \lambda_3$ → 3 eigenvectors (3 different directions)

If $\lambda_1 = \lambda_2 \neq \lambda_3$ → 2 eigenvectors (2 different directions)

If $\lambda_1 = \lambda_2 \neq \lambda_3$ → 3 eigenvectors (3 different directions)

If $\lambda_1 = \lambda_2 = \lambda_3$ → 1 eigenvector (1 direction)

If $\lambda_1 = \lambda_2 = \lambda_3$ → 2 eigenvectors (2 different directions)

If $\lambda_1 = \lambda_2 = \lambda_3$ → 3 eigenvectors (3 different directions)

DeepLearning.AI

Determinants and Eigenvectors

Dimensionality reduction annd projection

Dimensionality Reduction

- Reduce dimensions (# of columns) of dataset
- Preserve as much information as possible



Dimensionality Reduction

- Leads to smaller datasets
- Easier to visualize



Customer Age	Account Age	Days Since Login	Total Purchases	Total $ Spent
23	1 month	10 days	1	$100
71	45 months	2 days	3	$150
54	30 months	15 days	2	$70
36	22 months	12 days	4	$210

Easy approach - just delete columns

Loses valuable information

Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

1

1

=

$(1 + 1)$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

1

1

=

$(1 + 1)$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix}1\\1\end{matrix} \frac{1}{\sqrt{2}} = (1+1)/\sqrt{2}$$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

Norm of 1

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{1}{\sqrt{2}} = \frac{1}{\left\| \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\|_2}$$

$$(1+1)/\sqrt{2}$$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{1}{\sqrt{2}} = \begin{bmatrix} (1+1)/\sqrt{2} \\ \end{bmatrix}$$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{1}{\sqrt{2}} = \begin{bmatrix} (1+1)/\sqrt{2} \\ (1.2+1.6) \end{bmatrix}$$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{1}{\sqrt{2}} = \begin{bmatrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \end{bmatrix}$$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{1}{\sqrt{2}} = \begin{bmatrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \\ \end{bmatrix}$$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{1}{\sqrt{2}} = \begin{bmatrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \\ (-0.5+0.2) \end{bmatrix}$$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{1}{\sqrt{2}} = \begin{bmatrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \\ (-0.5+0.2)/\sqrt{2} \end{bmatrix}$$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{1}{\sqrt{2}} = \begin{bmatrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \\ (-0.5+0.2)/\sqrt{2} \\ \end{bmatrix}$$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{1}{\sqrt{2}} = \begin{bmatrix} (1+1)/\sqrt{2} \\ (1.2+1.6)/\sqrt{2} \\ (-0.5+0.2)/\sqrt{2} \\ (-1.3,-0.6) \end{bmatrix}$$



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{matrix}1\\1\end{matrix}\ \frac{1}{\sqrt{2}} = \begin{matrix}(1+1)/\sqrt{2}\\(1.2+1.6)/\sqrt{2}\\(-0.5+0.2)/\sqrt{2}\\(-1.3,-0.6)/\sqrt{2}\end{matrix}$$

Final coordinates



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{1}{\sqrt{2}} =$$

Final coordinates

1.4142
1.9799
-0.2121
-1.344



Projections

x	y
1.0	1.0
1.2	1.6
-0.5	0.2
-1.3	-0.6

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{1}{\sqrt{2}} =$$

Final coordinates

1.4142
1.9799
-0.2121
-1.344



Projections

To project a matrix A onto a vector v

$$A_P = A \frac{v}{\|v\|_2}$$

$r \times 1 \quad r \times c \quad c \times 1$

Projections

To project a matrix A onto vectors v_1 and v_2

$$A_P = A \overbrace{\left[\frac{v_1}{\|v_1\|_2} \quad \frac{v_2}{\|v_2\|_2} \right]}^{V}$$

$r \times 2 \qquad r \times c \qquad c \times 2$

Projections

To project a matrix A onto vectors v_1 and v_2

$$A_P = A \overbrace{\left[\frac{v_1}{\|v_1\|_2} \quad \frac{v_2}{\|v_2\|_2} \right]}^{V}$$

$r \times 2 \qquad r \times c \qquad c \times 2$

Projections

To project a matrix A onto vectors v_1 and v_2

$$A_P = AV$$

$r \times 2 \qquad r \times c \;\; c \times 2$

DeepLearning.AI

Determinants and Eigenvectors

Motivating PCA

Dimensionality Reduction



Principal Component Analysis (PCA)



Principal Component Analysis (PCA)



Principal Component Analysis (PCA)



Principal Component Analysis (PCA)



Principal Component Analysis (PCA)



Principal Component Analysis (PCA)



Principal Component Analysis (PCA)





Principal Component Analysis (PCA)





Principal Component Analysis (PCA)





Principal Component Analysis (PCA)





Principal Component Analysis (PCA)





Principal Component Analysis (PCA)





Principal Component Analysis (PCA)





Principal Component Analysis (PCA)



Principal Component Analysis (PCA)



Principal Component Analysis (PCA)





Principal Component Analysis (PCA)





Principal Component Analysis (PCA)



Benefits of Dimensionality Reduction

- Easier dataset to manage
- PCA reduces dimensions while minimizing information loss
- Simpler visualization



DeepLearning.AI

Determinants and Eigenvectors

Variance and covariance

Mean

"The average of the data"



$$Mean(x) = \frac{1}{n}\sum_{i=1}^{n} x_i$$

$$Mean(y) = \frac{1}{n}\sum_{i=1}^{n} y_i$$

Variance



Variance



Variance



Variance



Variance

$$Variance(x) = \frac{1}{n-1} \sum_{i=1}^{n} (x_i - Mean(x))^2 \quad = 16$$

	x_i	x_i - $Mean(x)$	$(x_i - Mean(x))^2$
1	10	1	1
2	4	-5	25
3	11	2	4
4	14	5	25
5	6	-3	9

→ 64

$$Mean(x) = 9$$

Variance

$$Variance(x) = \frac{1}{n-1}\sum_{i=1}^{n}(x_i - Mean(x))^2$$

$Var(x)$ μ_x

“The average squared distance from the mean”

Variance

$$Var(y) = \frac{1}{n-1}\sum_{i=1}^{n}(y_i - \mu_Y)^2$$



$$Var(x) = \frac{1}{n-1}\sum_{i=1}^{n}(x_i - \mu_x)^2$$

Problem



Solution: Covariance



Negative covariance

Positive covariance

Covariance

"Take the average"

$$Cov(x, y) = \frac{1}{n-1} \sum_{i=1}^{n} (x_i - \mu_x)(y_i - \mu_y)$$

$$Var(x) = \frac{1}{n-1} \sum_{i=1}^{n} (x_i - \mu_x)^2 (x_i - \mu_x)$$



negative covariance



covariance zero (or very small)



positive covariance

Covariance

$$Cov(x, y) = \frac{1}{n-1} \sum_{i=1}^{n} (x_i - \mu_x)(y_i - \mu_y)$$

“The direction of the relationship between two variables”



negative
covariance



covariance zero
(or very small)



positive
covariance

DeepLearning.AI

Determinants and Eigenvectors

The covariance matrix

Covariance matrix



Covariance matrix







$$\begin{bmatrix} 3 & \\ & 1 \end{bmatrix}$$

$$\begin{bmatrix} 3 & \\ & 1 \end{bmatrix}$$

$$\begin{bmatrix} 3 & \\ & 1 \end{bmatrix}$$

Covariance matrix



$$\begin{bmatrix} 3 & -2 \\ -2 & 1 \end{bmatrix}$$



$$\begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix}$$



$$\begin{bmatrix} 3 & 2 \\ 2 & 1 \end{bmatrix}$$

Covariance matrix



$$C = \begin{matrix} & \begin{matrix} x & \quad y \end{matrix} \\ \begin{matrix} x \\ y \end{matrix} & \begin{bmatrix} Var(x) & Cov(x,y) \\ Cov(y,x) & Var(y) \end{bmatrix} \end{matrix}$$

$$Cov(x, x) = Var(x)$$

Covariance matrix

$$\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix}$$

Covariance matrix

$$A = \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} \qquad \mu = \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}$$

$$C = \frac{1}{n-1}(\quad - \quad)^T(\quad - \quad)$$

Covariance matrix

$$A = \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} \qquad \mu = \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}$$

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu)$$

Covariance matrix

$$A = \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} \qquad \mu = \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}$$

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu) = \frac{1}{n-1}\left(\quad - \quad \right)^T \left(\quad - \quad \right)$$

$$= \frac{1}{n-1} \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}^T \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ x_2 - \mu_x & y_2 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix}$$

Covariance matrix

$$A = \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} \qquad \mu = \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}$$

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu) = \frac{1}{n-1}\left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)^T \left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}^T \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

Covariance matrix

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu) = \frac{1}{n-1}\left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)^T \left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}^T \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

$$= \frac{1}{n-1}\underbrace{\begin{bmatrix} x_1-\mu_x & x_2-\mu_x & \dots & x_n-\mu_n \\ y_1-\mu_y & y_2-\mu_y & \dots & y_n-\mu_y \end{bmatrix}}_{2 \times n} \underbrace{\begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}}_{n \times 2}$$

Covariance matrix

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu) = \frac{1}{n-1}\left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)^T \left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}^T \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & x_2-\mu_x & \dots & x_n-\mu_n \\ y_1-\mu_y & y_2-\mu_y & \dots & y_n-\mu_y \end{bmatrix} \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

$$(x_1-\mu_x)(x_1-\mu_x) + (x_2-\mu_x)(x_2-\mu_x) + \dots + (x_n-\mu_x)(x_n-\mu_x)$$

Covariance matrix

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu) = \frac{1}{n-1}\left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)^T \left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}^T \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & x_2-\mu_x & \dots & x_n-\mu_n \\ y_1-\mu_y & y_2-\mu_y & \dots & y_n-\mu_y \end{bmatrix} \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

$$\sum_{i=1}^{n}(x_i-\mu_x)^2 = Var(x)$$

Covariance matrix

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu) = \frac{1}{n-1}\left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)^T \left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}^T \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & x_2-\mu_x & \dots & x_n-\mu_n \\ y_1-\mu_y & y_2-\mu_y & \dots & y_n-\mu_y \end{bmatrix} \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix} = \begin{bmatrix} Var(x) & \\ & \end{bmatrix}$$

$$\frac{1}{n-1}\sum_{i=1}^{n}(x_i-\mu_x)^2 = Var(x)$$

Covariance matrix

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu) = \frac{1}{n-1}\left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)^T \left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}^T \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & x_2-\mu_x & \dots & x_n-\mu_n \\ y_1-\mu_y & y_2-\mu_y & \dots & y_n-\mu_y \end{bmatrix} \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix} = \begin{bmatrix} Var(x) & \\ & \end{bmatrix}$$

$$(x_1-\mu_x)(y_1-\mu_y) + (x_2-\mu_x)(y_2-\mu_y) + \dots + (x_n-\mu_x)(y_n-\mu_y)$$

Covariance matrix

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu) = \frac{1}{n-1}\left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)^T \left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}^T \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & x_2-\mu_x & \dots & x_n-\mu_n \\ y_1-\mu_y & y_2-\mu_y & \dots & y_n-\mu_y \end{bmatrix} \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix} = \begin{bmatrix} Var(x) & \\ & \end{bmatrix}$$

$$\sum_{i=1}^{n}(x_i-\mu_x)(y_i-\mu_y) = Cov(x,y)$$

Covariance matrix

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu) = \frac{1}{n-1}\left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)^T \left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}^T \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & x_2-\mu_x & \dots & x_n-\mu_n \\ y_1-\mu_y & y_2-\mu_y & \dots & y_n-\mu_y \end{bmatrix} \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix} = \begin{bmatrix} Var(x) & Cov(x,y) \\ & \end{bmatrix}$$

$$= \frac{1}{n-1}\sum_{i=1}^{n}(x_i-\mu_x)(y_i-\mu_y) = Cov(x,y)$$

Covariance matrix

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu) = \frac{1}{n-1}\left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)^T \left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}^T \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & x_2-\mu_x & \dots & x_n-\mu_n \\ y_1-\mu_y & y_2-\mu_y & \dots & y_n-\mu_y \end{bmatrix} \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix} = \begin{bmatrix} Var(x) & Cov(x,y) \\ Cov(y,x) & \end{bmatrix}$$

Covariance matrix

$$C = \frac{1}{n-1}(A-\mu)^T(A-\mu) = \frac{1}{n-1}\left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)^T \left(\begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \end{bmatrix} - \begin{bmatrix} \mu_x & \mu_y \\ \mu_x & \mu_y \\ \vdots & \vdots \\ \mu_x & \mu_y \end{bmatrix}\right)$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}^T \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix}$$

$$= \frac{1}{n-1}\begin{bmatrix} x_1-\mu_x & x_2-\mu_x & \dots & x_n-\mu_n \\ y_1-\mu_y & y_2-\mu_y & \dots & y_n-\mu_y \end{bmatrix} \begin{bmatrix} x_1-\mu_x & y_1-\mu_y \\ x_2-\mu_x & y_2-\mu_y \\ \vdots & \vdots \\ x_n-\mu_x & y_n-\mu_y \end{bmatrix} = \begin{bmatrix} Var(x) & Cov(x,y) \\ Cov(y,x) & Var(y) \end{bmatrix}$$

Matrix formula

$$A - \mu = \begin{bmatrix} x_1 - \mu_x & y_1 - \mu_y \\ \vdots & \vdots \\ x_n - \mu_x & y_n - \mu_y \end{bmatrix} \qquad C = \frac{1}{n-1}(A - \mu)^T(A - \mu)$$



Similar X and Y variances
Negative covariance

A

x	y
10	5
12	3
6	9
6	4
5	11
14	2
8	1
3	13

$\mu_x = 8 \quad \mu_y = 6$

$A - \mu$

$x - \mu_x$	$y - \mu_y$
2	-1
4	-3
-2	3
-2	-2
-3	5
6	-4
0	-5
-5	8

$\frac{1}{7}$

$(A - \mu)^T$

$x - \mu_x$	2	4	-2	-2	-3	6	0	-5
$y - \mu_y$	-1	-3	3	-2	5	-4	-5	8

$A - \mu$

$x - \mu_x$	$y - \mu_y$
2	-1
4	-3
-2	3
-2	-2
-3	5
6	-4
0	-5
-5	8

$$C = \begin{bmatrix} 14 & -11.86 \\ -11.86 & 19.71 \end{bmatrix}$$

Matrix formula

$$A = \begin{bmatrix} x_1 & y_1 & z_1 \\ \vdots & \vdots & \vdots \\ x_n & y_n & z_n \end{bmatrix} \qquad C = \frac{1}{n-1}(A-\mu)^T(A-\mu)$$

1. Arrange data with a different feature in each column
2. Calculate column averages
3. Subtract each average from their respective column to generate $A - \mu$
4. $\frac{1}{n-1}(A-\mu)^T(A-\mu)$ gives the covariance matrix C

DeepLearning.AI

Determinants and Eigenvectors

PCA - Overview

Principal Component Analysis (PCA)



Principal Component Analysis (PCA)



Principal Component Analysis (PCA)



Principal Component Analysis (PCA)



Best line = most variance

Principal Component Analysis (PCA)



Best line = most variance

Projections

Eigenvalues / Eigenvectors

Covariance Matrix

PCA

Principal Component Analysis (PCA)



Principal Component Analysis (PCA)



$$C = \begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

$$\begin{bmatrix} 2 \\ 1 \end{bmatrix} \quad \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

Eigenvectors (direction)

11 1

Eigenvalues (magnitude)

Principal Component Analysis (PCA)



$$C = \begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

$$\begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Eigenvectors
(direction)

11

Eigenvalues
(magnitude)

Principal Component Analysis (PCA)



$$C = \begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

$$\begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Eigenvectors
(direction)

11

Eigenvalues
(magnitude)

Principal Component Analysis (PCA)



$$C = \begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

$$\begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Eigenvectors (direction)

11

Eigenvalues (magnitude)

Principal Component Analysis (PCA)



Principal Component Analysis (PCA)



PCA: Principal Component Analysis



PCA: Principal Component Analysis



PCA: Principal Component Analysis



PCA: Principal Component Analysis



DeepLearning.AI

Determinants and Eigenvectors

PCA - Why it works

PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"





PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"





PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"





PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"





PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"





PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"





PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"





PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"



PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"



PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"





PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"





PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"





PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

“How spread out is the data?”



PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"





PCA: Why It Works



Covariance matrix

$$C$$

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"



PCA: Why It Works



Covariance matrix

C

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"

Eigenvalues and Eigenvectors

$$\begin{matrix} 11 \\ \begin{bmatrix} 2 \\ 1 \end{bmatrix} \end{matrix} \quad \begin{matrix} 1 \\ \begin{bmatrix} -1 \\ 2 \end{bmatrix} \end{matrix}$$

"Everything is just stretches"







PCA: Why It Works



Covariance matrix

C

$$\begin{bmatrix} 9 & 4 \\ 4 & 3 \end{bmatrix}$$

"How spread out is the data?"

Eigenvalues and Eigenvectors

$$\begin{matrix} 11 \\ \begin{bmatrix} 2 \\ 1 \end{bmatrix} \end{matrix} \quad \begin{matrix} 1 \\ \begin{bmatrix} -1 \\ 2 \end{bmatrix} \end{matrix}$$

"Everything is just stretches"



Norm Equation

$$\sqrt{x^2 + y^2}$$





DeepLearning.AI

Determinants and Eigenvectors

PCA - Mathematical formulation

PCA Mathematical formulation

You have n observations of 5 variables $(x_1, x_2, x_3, x_4, x_5)$

Goal: Reduce to 2 variables

1 Create matrix

5 variables

$$X = \begin{bmatrix} x_{11} & x_{12} & \cdots & x_{15} \\ x_{21} & x_{22} & \cdots & x_{25} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{n5} \end{bmatrix}$$

n Observations

2 Center the data

$$X - \mu = \begin{bmatrix} x_{11} - \mu_1 & x_{12} - \mu_2 & \cdots & x_{15} - \mu_5 \\ x_{21} - \mu_1 & x_{22} - \mu_2 & \cdots & x_{25} - \mu_5 \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} - \mu_1 & x_{n2} - \mu_2 & \cdots & x_{n5} - \mu_5 \end{bmatrix}$$

PCA Mathematical formulation

You have n observations of 5 variables $(x_1, x_2, x_3, x_4, x_5)$

Goal: Reduce to 2 variables

3 Calculate Covariance Matrix

$$C = \frac{1}{n-1}(X-\mu)^T(X-\mu) = \begin{bmatrix} Var(X_1) & Cov(X_1,X_2) & Cov(X_1,X_3) & Cov(X_1,X_4) & Cov(X_1,X_5) \\ Cov(X_1,X_2) & Var(X_2) & Cov(X_2,X_3) & Cov(X_2,X_4) & Cov(X_2,X_5) \\ Cov(X_1,X_3) & Cov(X_2,X_3) & Var(X_3) & Cov(X_3,X_4) & Cov(X_3,X_5) \\ Cov(X_1,X_4) & Cov(X_2,X_4) & Cov(X_3,X_4) & Var(X_4) & Cov(X_4,X_5) \\ Cov(X_1,X_5) & Cov(X_2,X_5) & Cov(X_3,X_5) & Cov(X_4,X_5) & Var(X_5) \end{bmatrix}$$

PCA Mathematical formulation

You have n observations of 5 variables $(x_1, x_2, x_3, x_4, x_5)$

Goal: Reduce to 2 variables

4 Calculate Eigenvectors and Eigenvalues

5 Create Projection Matrix

6 Project Centered Data



$$V = \left[\frac{}{\|v_1\|_2} \quad \frac{}{\|v_2\|_2} \right]$$

$$X_{PCA} = (X - \mu)V$$

DeepLearning.AI

Determinants and Eigenvectors

Conclusion