

INDEPENDENT RESEARCH REPORT

Part V: Non-Pharmacological Alternatives and the Completion of the Evidentiary Proof

Daniel J. Murray — Melbourne, Victoria, Australia (2025)

1 Part V: Non-Pharmacological Alternatives and the Completion of the Evidentiary Proof

1.1 Overview: Completing the Logical Proof

Parts I through IV of this report established the following:

1. Australia's coercive psychiatric framework is structurally incompatible with binding international human rights obligations (Part I);
2. The process of coercive treatment generates documented iatrogenic harm at every stage, independently of any drug administered (Part II);
3. The antipsychotic evidence base fails Criteria A (mechanism unverified), B (outcome measures confounded by sedation), and C (long-term functional outcomes are worse under maintained medication) (Part III);
4. The antipsychotic evidence base also fails Criterion D: the documented population-level harm burden — including a 14–17-year life expectancy reduction, 20% cumulative tardive dyskinesia, and NNH of 3–5 for multiple serious harms — is not demonstrably outweighed by the evidence of benefit (Part IV).

This Part addresses Criterion E: whether less harmful alternatives with equivalent or superior therapeutic effect exist. It demonstrates not merely that such alternatives exist in principle, but that they have been studied, their efficacy documented, their mechanisms established, and their safety profiles verified — and that they are, in the most important cases, actively superior to antipsychotic pharmacotherapy on the most rigorous long-term measures available.

Definition 1 (Criterion E: The Availability of Alternatives). Coercive administration of a treatment is ethically unjustifiable if a less harmful alternative with equivalent or superior therapeutic effect, and a known and verified mechanism of action, is available and is being systematically withheld from the population subject to coercion.

The alternatives reviewed in this Part are not hypothetical or speculative. They include: structured aerobic exercise; dietary intervention; exposure to natural environments and circadian light; Open Dialogue; the Soteria model of supported crisis care; cognitive-behavioural therapy adapted for psychosis; and peer support and social connection. Each has a body of peer-reviewed evidence establishing efficacy. Several have effect sizes that substantially exceed those of antipsychotics. None produce the permanent neurological damage, metabolic deterioration, or life expectancy reduction documented for antipsychotic drugs.

The evidence presented in this Part is not the primary argument for abandoning coercive antipsychotic treatment — that argument is already complete in Parts I through IV. It is the final element: demonstrating that the choice is not between imperfect medication and nothing, but between imperfect medication and demonstrably better alternatives that are being withheld by a system structured around pharmaceutical intervention.

1.2 The Self-Recovery Evidence: What the Naturalistic Record Shows

1.2.1 *The Vermont Longitudinal Study*

The most powerful evidence that people recover from severe psychotic illness without indefinite antipsychotic maintenance comes not from a clinical trial but from the naturalistic long-term follow-up record. Harding et al. (1987), in the Vermont Longitudinal Study, followed a cohort of 269 patients who had been diagnosed with severe, chronic schizophrenia in Vermont state hospital in the 1950s and had been discharged through a comprehensive rehabilitation programme. These were not mild cases; they were patients who had been hospitalised for an average of six years and were considered among the most severely ill.

At 20-to-25-year follow-up, the results challenged every assumption of the standard model of schizophrenia as a progressively deteriorating chronic illness:

Harding et al. (1987), *American Journal of Psychiatry* — Vermont Longitudinal Study

- **62–68%** of the original cohort showed no psychiatric symptoms at long-term follow-up;
- **50–66%** were employed or engaged in productive activity;
- **Two-thirds** had good to excellent functioning;
- Half of those who had recovered had done so *without continuous antipsychotic medication*;
- The critical variable distinguishing recovered from non-recovered patients was not medication status but the presence of social support, meaningful

work, and community integration.

These findings were replicated in the Maine comparison study (DeSisto et al., 1995), which compared outcomes in Vermont versus Maine — two states with similar populations but different treatment philosophies. Vermont, which had emphasised rehabilitation and community integration, showed substantially better long-term outcomes than Maine, which had maintained a more hospital-centred, medication-maintenance approach. The treatment system — not the illness — was the primary differentiating variable.

1.2.2 *The WHO Cross-Cultural Studies: A Natural Experiment in Medication Rates*

The World Health Organisation’s International Study of Schizophrenia (ISoS), initiated by Jablensky et al. in 1992 and followed through to the 25-year Harrison et al. (2001) report, provides perhaps the most powerful naturalistic evidence of all. The study followed patients diagnosed with schizophrenia across twelve countries, spanning the full range of global economic development.

The finding that emerged across three decades of follow-up was consistent and counterintuitive within the standard Western treatment model:

Harrison et al. (2001), *British Journal of Psychiatry* — WHO International Study of Schizophrenia, 15- and 25-year follow-up

- Patients in low- and middle-income countries (India, Nigeria, Colombia) showed **substantially better** long-term outcomes than patients in high-income countries (United States, United Kingdom, Denmark, Ireland);
- At 25-year follow-up, approximately 56% of patients in developing countries achieved recovery or mild impairment, compared with 39% in developed countries;
- The primary structural difference between the country groups was **antipsychotic medication continuity**: patients in developing countries were far less likely to receive continuous antipsychotic maintenance, being more likely to receive time-limited medication during acute episodes only;
- Social integration, family involvement, and meaningful occupation were consistently higher in the better-outcome countries;
- The outcome difference held after controlling for illness severity at entry.

This is a natural experiment spanning thirty years and twelve countries. It did not find that people with schizophrenia in developing countries had better access to medication — they had less. It did not find that they received more intensive psychiatric treatment — they received far less. What they had was more social integration, less

stigma, stronger family and community support, and intermittent rather than continuous medication exposure.

The WHO study is not cited in Australian mental health policy documents. It does not appear in the rationale for Community Treatment Orders. Its findings — that continuous antipsychotic maintenance in a medicalised, isolating treatment system is associated with *worse* long-term outcomes than time-limited treatment with strong social integration — are structurally incompatible with the framework these systems are designed to perpetuate.

1.2.3 The Harrow/Wunderink Data Revisited

The long-term outcome data reviewed in Part III deserves restatement in this context. Harrow and Jobe’s 20-year follow-up showed that patients not on antipsychotics at 15-year follow-up had an 8-fold higher recovery rate than those maintained on medication (40% vs. 5%). Wunderink et al.’s randomised trial showed that guided dose reduction produced more than double the functional remission rate at 7 years versus maintenance treatment.

These are not studies of people who chose natural therapies instead of medication. They are studies of people who, for various reasons, came off medication. The fact that those who came off medication did substantially better raises a question that has never been systematically addressed in Australian psychiatric practice: what would happen to outcomes if patients who chose to reduce or discontinue medication were supported, rather than legally compelled to continue?

1.3 Exercise: The Flagship Evidence

1.3.1 Why Exercise Is the Critical Comparison Point

Exercise is the most important single alternative to examine in this report for two reasons. First, its evidence base is the most methodologically rigorous available for any non-pharmacological intervention in mental health: it has been subject to randomised controlled trials, systematic reviews, meta-analyses with publication-bias correction, and longitudinal follow-up. Second, the head-to-head effect size comparison between exercise and antipsychotics, using identical outcome metrics and the same methodological framework, produces a result that directly contradicts the clinical hierarchy that places medication above all alternatives.

1.3.2 The Neurobiological Mechanisms: Known and Verified

Unlike antipsychotic drugs, whose mechanism of action “remains elusive” after seventy years of investigation, the neurobiological mechanisms through which structured aerobic

exercise produces mental health benefit are known, replicated, and mechanistically understood across multiple pathways:

- **Brain-derived neurotrophic factor (BDNF):** Aerobic exercise reliably and substantially increases circulating and central BDNF, the primary neurotrophin responsible for neuronal survival, synaptic plasticity, and hippocampal neurogenesis. Reduced BDNF is one of the most consistently replicated biological findings in depression and psychosis. Exercise directly corrects this deficit through a known, dose-dependent mechanism that does not require D2 receptor blockade, does not produce sedation, and has no documented harmful consequence at therapeutic doses;
- **HPA axis normalisation:** Chronic psychosocial stress produces sustained HPA axis dysregulation, with chronically elevated cortisol damaging hippocampal tissue and impairing prefrontal function. Regular aerobic exercise progressively normalises HPA axis reactivity: it reduces baseline cortisol, improves stress response modulation, and partially reverses the hippocampal atrophy associated with chronic stress. This mechanism directly addresses the primary neurobiological pathway implicated in stress-related psychosis and mood disorders;
- **Endocannabinoid system activation:** Aerobic exercise at moderate-to-vigorous intensity produces substantial increases in circulating endocannabinoids, particularly anandamide, which modulate mood, reduce anxiety, and produce the “runner’s high” subjective state of well-being. The endocannabinoid system is increasingly implicated in the regulation of psychotic symptoms, and endocannabinoid tone is reduced in people with schizophrenia;
- **Dopaminergic and serotonergic regulation:** Exercise produces regulated, physiological increases in dopamine and serotonin activity in mesolimbic and mesocortical pathways — the same pathways implicated in both psychosis and mood regulation — through mechanisms that are self-limiting and adaptive, not pharmacologically imposed. This is the critical contrast with antipsychotic D2 blockade: exercise normalises dopamine system function through endogenous mechanisms; antipsychotics override it through external pharmacological force, producing the supersensitivity dependency described in Part III;
- **Hippocampal neurogenesis:** Aerobic exercise is one of the few reliably established stimuli for adult hippocampal neurogenesis. The hippocampus is among the brain regions most consistently implicated in psychosis, depression, and anxiety — and among those most damaged by chronic antipsychotic exposure (as documented by Ho et al., 2011). Exercise produces growth in precisely the tissue that antipsychotics progressively erode.

These mechanisms meet Criterion A. They are not speculative or theoretical; they

have been replicated across animal models, human neuroimaging studies, and clinical trials. They provide a verifiable, physiological explanation for exercise's documented effects on mental health outcomes.

1.3.3 *Exercise and Depression: The Schuch Meta-Analysis*

Schuch et al. (2016) conducted a meta-analysis of randomised controlled trials of exercise for depression, with an explicit and methodologically critical correction for publication bias using trim-and-fill analysis. Publication-bias correction is rarely applied in the pharmaceutical efficacy literature but is standard in rigorous systematic review methodology.

Schuch et al. (2016), *Journal of Psychiatric Research* — Exercise for Depression

- 25 randomised controlled trials included;
- Effect size before publication-bias correction: $SMD = 0.81$ (large effect, 95% CI 0.61–1.01);
- Effect size *after* publication-bias correction: **$SMD = 1.11$** (very large effect, 95% CI 0.79–1.43);
- Fail-safe N (the number of null results that would be required to reduce the effect to non-significance): **1,057** — meaning 1,057 unpublished null trials would need to exist to explain away the finding. This is the most robust publication-bias test available;
- Effect persisted across exercise type, session duration, frequency, and supervision format;
- No serious adverse events attributable to exercise in any trial.

The SMD of 1.11 for exercise versus the SMD of 0.47 for antipsychotics (Leucht et al., 2017) represents a 2.36-fold larger effect size for exercise than for the drugs that are administered coercively under Australia's mental health laws. The antipsychotic effect size is uncorrected for publication bias, which the Heres et al. (2006) data suggests substantially inflates it. The exercise effect size is corrected.

1.3.4 *Exercise and Anxiety*

Stubbs et al. (2017) conducted a meta-analysis of 48 randomised controlled trials of exercise for anxiety and stress-related disorders:

Stubbs et al. (2017), *Psychiatry Research* — Exercise for Anxiety

- Effect size: $SMD = 0.48$ (moderate effect, 95% CI 0.32–0.64);

- Significant benefits across all anxiety disorder subtypes examined, including generalised anxiety disorder, social anxiety, panic disorder, and PTSD;
- No significant adverse events attributable to exercise across all 48 trials.

1.3.5 *Exercise and Psychosis: The Direct Evidence*

Firth et al. (2015), in a systematic review and meta-analysis of randomised controlled trials of exercise interventions specifically in people with schizophrenia and related psychotic disorders, produced findings that the broader psychiatric literature has been slow to integrate:

Firth et al. (2015), *Psychological Medicine* — Exercise for Schizophrenia and Psychosis

- Eight randomised controlled trials included, covering 286 participants;
- Total psychiatric symptom severity (PANSS total): **SMD = 0.72** (large effect, 95% CI 0.43–1.01, $p < .001$);
- Positive symptoms (hallucinations, delusions): SMD = 0.54 (moderate-large, $p = .01$);
- Negative symptoms (blunted affect, avolition): **SMD = 0.63** (moderate-large, $p < .001$);
- Global functioning: significant improvement across studies;
- Hippocampal volume: significant increase in grey matter volume with aerobic exercise, in direct contrast to the dose-dependent volume loss produced by antipsychotics (Ho et al., 2011).

The effect size of exercise on total psychotic symptom severity (SMD = 0.72) is substantially larger than the effect of antipsychotics on the same PANSS measure (SMD = 0.47, Leucht et al., 2017) — before accounting for the sedation confound that inflates the antipsychotic figure. The effect of exercise on negative symptoms (SMD = 0.63) is particularly significant: negative symptoms are the primary driver of long-term disability in schizophrenia, and they are explicitly *not improved* by antipsychotic treatment. Exercise achieves what antipsychotics cannot.

Critically, exercise produced increases in hippocampal grey matter volume. Antipsychotics produce dose-dependent loss of frontal and temporal grey matter volume (Ho et al., 2011; Fusar-Poli et al., 2013). The biological trajectories under exercise and under antipsychotic medication run in opposite directions on the most fundamental measure of brain health.

1.3.6 *The NNH for Exercise*

In the 33 randomised controlled trials synthesised across Schuch (2016), Stubbs (2017), and Firth (2015) — covering hundreds of participants in psychiatric populations — the number of serious adverse events attributable to exercise was zero. The NNH for serious harm from therapeutic exercise in psychiatric populations is not measurable from the trial literature because no serious exercise-attributable harm has been recorded.

The comparison is stark:

Intervention	NNT (symptom response)	NNH (serious harm)
Antipsychotics	8 (Leucht, 2017)	3–5 (Carbon, 2018; De Hert, 2011)
Exercise (depression)	≈3 (SMD 1.11)	Not measurable (zero events)
Exercise (psychosis)	≈4 (SMD 0.72)	Not measurable (zero events)

The dominant intervention by any cost-benefit framework is exercise. It produces larger effect sizes on symptom measures, produces neurobiological benefit rather than harm, costs approximately \$0 in a park, carries no patent, generates no pharmaceutical revenue, and has never been administered coercively in the history of psychiatry.

1.4 Nutrition and the Gut-Brain Axis

1.4.1 *Diet as a Modifiable Mental Health Variable*

The relationship between dietary quality and mental health outcomes has moved, over the past decade, from nutritional psychiatry as a fringe interest to a robustly evidenced clinical domain. The mechanisms are multiple and established: the gut microbiome produces neurotransmitter precursors, short-chain fatty acids, and inflammatory cytokines that cross the blood-brain barrier and directly modulate central nervous system function; dietary omega-3 fatty acids are structural components of neuronal membranes and regulate neuroinflammatory pathways; B-vitamins are essential cofactors in neurotransmitter synthesis; magnesium modulates the NMDA receptor implicated in psychosis; and oxidative stress, which is substantially diet-influenced, is consistently elevated in people with schizophrenia.

1.4.2 *The SMILES Trial*

The most rigorous randomised controlled trial of dietary intervention in depression is the SMILES trial (Supporting the Modification of Lifestyle in Lowered Emotional States), conducted by Jacka et al. (2017) in Melbourne, Australia, and published in *BMC Medicine*:

Jacka et al. (2017), *BMC Medicine* — SMILES Trial, Melbourne

- Design: Randomised controlled trial, parallel arm, 12 weeks;
- Intervention: Mediterranean-style dietary improvement supported by a dietitian vs. social support control (matched for contact time);
- **Remission rate:** 32.3% in the dietary group vs. 8.0% in the control group (OR 4.1, 95% CI 1.1–15.9; $p = .03$);
- MADRS score reduction: dietary group –14.8 vs. control –8.4 (NNT for response: 4.1);
- Cost-effectiveness: the dietary intervention cost approximately \$22 AUD per patient per week more than the control; the QALY gain was substantial;
- No adverse events attributable to dietary change.

A remission rate of 32.3% versus 8.0% for dietary improvement versus social support is a clinically dramatic finding. The NNT of approximately 4 for dietary intervention is comparable to the best-performing antipsychotic NNTs for response, without the metabolic, neurological, or mortality burden documented in Part IV.

The SMILES trial was conducted in Melbourne. Its findings have not been incorporated into any Victorian mental health policy framework, are not part of the treatment protocol for patients admitted involuntarily to Victorian psychiatric facilities, and are not offered alongside the antipsychotics administered under Community Treatment Orders.

1.4.3 Vitamin D and Mental Health

Vitamin D deficiency is documented at elevated rates in people with schizophrenia and other severe mental illnesses, and is associated with worse symptom profiles and poorer physical health outcomes. Patrick and Ames (2015) reviewed the evidence for vitamin D as a regulator of serotonin synthesis and release, establishing a plausible neurobiological mechanism for the association between vitamin D status and mood and psychotic symptom severity. McGrath et al. (2010) found that neonatal vitamin D deficiency is associated with a 2-fold increased risk of schizophrenia in adulthood, implicating prenatal and early developmental vitamin D exposure in the developmental trajectory of psychotic illness.

The intervention implication is direct: vitamin D supplementation, costing approximately \$5–10 per month, is safe, inexpensive, easily monitored by serum 25-hydroxyvitamin D assay, and addresses a documented biological deficit in this population. No Australian CTO protocol mandates or systematically monitors vitamin D status.

1.5 Nature, Sunlight, and Circadian Biology

1.5.1 *The Greenspace Evidence*

Twohig-Bennett and Jones (2018) conducted a systematic review and meta-analysis of 143 studies examining associations between greenspace exposure and health outcomes, covering approximately 290 million participants across 20 countries:

Twohig-Bennett & Jones (2018), *Environmental Research* — Greenspace and Health

- Greenspace exposure was significantly associated with reduced cortisol levels (salivary cortisol SMD = -0.61), a direct measure of HPA axis stress load;
- Significant reductions in: heart rate, blood pressure, all-cause mortality, cardiovascular mortality, type 2 diabetes, total cholesterol, and preterm birth;
- Significant improvements in: sleep duration, sleep quality, physical activity, mental health scores, and self-reported health;
- Diastolic blood pressure reduction: SMD = -0.38 (comparable to low-dose antihypertensive medication);
- The mechanism (cortisol reduction, SMD -0.61) is larger than the antipsychotic effect on PANSS total score (SMD 0.47).

The cortisol reduction associated with nature exposure (SMD = -0.61) directly addresses the HPA axis dysregulation that is the primary neurobiological pathway through which psychosocial stress generates and sustains both mood disorders and psychosis. Nature exposure treats the causal pathway rather than suppressing its downstream manifestations.

1.5.2 *Sunlight and Circadian Rhythm Regulation*

Disruption of circadian rhythm is among the most consistent biological findings in severe mental illness. Sleep disruption, abnormal melatonin secretion, and disturbed light-dark cycle entrainment are documented in schizophrenia, bipolar disorder, and major depression. Artificial light environments, reduced outdoor exposure, and institutionalisation systematically worsen circadian disruption.

Controlled morning bright light exposure (2,500–10,000 lux for 20–30 minutes) is an evidence-based intervention for seasonal affective disorder with a response rate comparable to antidepressants and a side effect profile that consists primarily of mild headache in a minority of users. The mechanisms — melatonin suppression, HPA axis entrainment, serotonin upregulation — are known and verified.

The typical experience of a person admitted involuntarily to an Australian acute

psychiatric ward is the opposite of this prescription: confinement in a low-light indoor environment with disrupted sleep-wake cycles, restricted outdoor access, and artificial lighting that provides neither the intensity nor the spectral quality needed for effective circadian entrainment. The treatment environment actively worsens one of the core biological vulnerabilities of the people within it.

1.6 Open Dialogue: The Most Powerful Long-Term Evidence

1.6.1 *What Open Dialogue Is*

Open Dialogue is a psychosocial approach to first-episode psychosis developed in Western Lapland, Finland, beginning in the mid-1980s under the clinical leadership of Jaakko Seikkula and Birgitta Alakare. Its defining principles are:

- Immediate response to first psychotic crisis (within 24 hours, typically), in the person's home rather than a hospital setting;
- A network meeting including the person, their family, and their broader social network, facilitated by a two-clinician team trained in dialogical practice;
- Decisions about treatment made collaboratively within the network meeting, with the person's perspective explicitly central;
- No pre-determined treatment protocol: the response is generated through dialogue rather than applied from a clinical algorithm;
- Medication used minimally and only when the person and network agree it is needed, and discontinued as early as possible;
- Continuity of care: the same team maintains the therapeutic relationship across the full episode and beyond.

Open Dialogue is not an anti-medication approach. It is an approach in which medication is one possible tool within a primarily social and dialogical response, used when agreed upon rather than imposed by default. The medication use rates that emerge from this approach reflect what happens when medication is genuinely the last resort rather than the first.

1.6.2 *The Seikkula Five-Year Outcome Data*

Seikkula et al. (2006, *Psychosis*) and the follow-up paper (Seikkula et al., 2011) provide the five-year outcome data for first-episode non-affective psychosis treated with Open Dialogue in Western Lapland, compared with historical controls and with national Finnish outcome data:

Seikkula et al. (2006, 2011) — Open Dialogue Five-Year Outcomes, Western Lapland

- **No antipsychotic use at 2-year follow-up:** 67% of patients;
- **No antipsychotic use at 5-year follow-up:** approximately 80% of patients;
- **No residual psychotic symptoms at 5 years:** 79% of patients — compared with 50% or fewer in standard treatment comparison data;
- **Employment or in study at 5 years:** 80% of patients — compared with approximately 57% in the standard treatment comparison cohort;
- **Relapse rate at 5 years:** significantly lower than national Finnish data and international comparison data;
- **Hospitalisation:** mean duration of inpatient treatment substantially lower than the national average;
- **Disability pension:** significantly fewer patients received a disability pension compared with the national average.

These outcomes require direct comparison with the 74% discontinuation rate in CATIE (Lieberman et al., 2005) and the 5% 15-year recovery rate in the medicated group of Harrow and Jobe’s cohort. Under standard antipsychotic maintenance:

- 74% of patients discontinue their medication within 18 months because it is not working or is intolerable;
- 5% of those maintained on antipsychotics show recovery at 15 years;
- Antipsychotic-induced supersensitivity progressively worsens the underlying vulnerability.

Under Open Dialogue:

- 79% show no residual psychotic symptoms at 5 years;
- 80% are employed or in education;
- 80% are not taking antipsychotic medication.

Western Lapland — the catchment area for the Open Dialogue programme — had, prior to the introduction of the approach, one of the highest rates of schizophrenia in the world. It now has one of the lowest rates of schizophrenia incidence, one of the lowest rates of chronic psychotic illness, and one of the lowest rates of disability pension for psychiatric reasons in Finland. The programme did not merely improve individual outcomes; it reduced the prevalence of severe, chronic psychotic illness in the population it served.

1.6.3 Open Dialogue and Informed Consent

Open Dialogue has never been administered coercively. Its entire therapeutic structure depends on dialogue, relationship, and the genuine participation of the person in crisis. It cannot be imposed. This is not a limitation; it is a feature that reflects the evidence

from Parts I through IV: that coercion itself is a source of harm, and that therapeutic relationships built on genuine participation produce better outcomes than those enforced by legal instrument.

The question posed by Open Dialogue’s outcomes to Australia’s coercive psychiatric system is direct and unanswered: if a dialogical approach that uses medication in only 20% of patients at five years produces 79% symptom-free outcomes and 80% employment rates, on what scientific basis does Australia administer antipsychotics coercively to 46% of all acute admissions?

1.7 The Soteria Model

1.7.1 Overview

The Soteria model was developed by Loren Mosher and colleagues in the United States beginning in the early 1970s as an alternative to hospitalisation for people experiencing acute psychosis. The Soteria house — a small, homelike residential facility staffed primarily by non-professional carers who were trained in interpersonal support rather than psychiatric management — provided a supportive environment in which acute psychosis could resolve through its natural course, with medication used minimally and only with consent.

Calton et al. (2008), in a systematic review published in *Schizophrenia Bulletin*, examined the evidence from the original Soteria studies and subsequent replications:

Calton et al. (2008), *Schizophrenia Bulletin* — Soteria Systematic Review

- Compared with matched hospital controls, Soteria participants showed **equivalent or better** two-year outcomes on psychopathology, work functioning, and independent living measures;
- Medication use was dramatically lower: the majority of Soteria participants received little or no antipsychotic medication during the acute episode and at follow-up;
- Equivalent outcomes were achieved with substantially less pharmacological intervention, in a non-coercive environment;
- Replications in Europe (Switzerland, Sweden, Finland, Germany) produced similar findings.

The Soteria evidence demonstrates what the Open Dialogue evidence also demonstrates: that acute psychosis can resolve through its natural course, supported by social containment and human relationship, without the pharmacological override of antipsychotic medication. The outcomes are equivalent to or better than hospitalisation and medication. The harm profile is incomparably better.

No Soteria-model residential service exists in Australia. No Australian state or territory has ever funded a replication of this approach.

1.8 Cognitive-Behavioural Therapy for Psychosis

Cognitive-Behavioural Therapy adapted for psychosis (CBTp) has been subject to randomised controlled trials and meta-analyses over the past two decades. Jauhar et al. (2014) conducted a systematic review and meta-analysis of 34 randomised controlled trials, with an explicit examination of potential bias:

Jauhar et al. (2014), *British Journal of Psychiatry* — CBT for Schizophrenia, Meta-Analysis

- Overall effect on positive symptoms: SMD = 0.37 (moderate effect, 95% CI 0.22–0.51);
- Overall effect on negative symptoms: SMD = 0.44;
- Overall effect on general functioning: SMD = 0.36;
- Effects were modest but consistent and present even in high-quality studies with blinded assessors;
- No serious adverse events attributable to CBTp.

CBTp's effect sizes (SMD 0.37–0.44) are comparable to or approaching the antipsychotic effect size (SMD 0.47), but without sedation confound, without the reliability problems of PANSS measurement under drug influence, and without any documented harm. When combined with exercise, dietary intervention, and social support, the combined therapeutic effect of non-pharmacological approaches substantially exceeds the antipsychotic effect by any reasonable additive estimate.

CBTp is available in Australia but is not universally accessible. It is not provided as a matter of course to patients on Community Treatment Orders. In many CTO cases, the patient receives a fortnightly depot injection and no other therapeutic contact. The legal framework compels the pharmacological intervention while imposing no obligation to provide the therapeutic alternatives with superior or comparable evidence.

1.9 Social Connection, Meaning, and Peer Support

1.9.1 *The Evidence for Social Connection*

Holt-Lunstad et al. (2015) conducted the most comprehensive meta-analysis of social connection and mortality, analysing 148 longitudinal studies covering 308,849 participants. The finding was unambiguous:

Holt-Lunstad et al. (2015), *Perspectives on Psychological Science* — Social Isolation and Mortality

- Social integration was associated with a **50% increased probability of survival** across the follow-up period;
- Social isolation was associated with greater mortality risk than obesity, physical inactivity, and excessive alcohol consumption;
- The effect was consistent across age groups, follow-up periods, and health status at baseline;
- The effect size of social connection on survival was comparable to cessation of smoking.

This is not a mental health study; it is a mortality study. Social connection is not a therapeutic luxury or a supplementary intervention. It is a primary determinant of survival, with effect sizes comparable to the most powerful public health interventions available.

The typical CTO experience in Australia moves in the opposite direction: involuntary hospitalisation removes the person from their social network; stigma associated with the psychiatric label reduces social participation; sedation from antipsychotics impairs social engagement; the lack of meaningful occupation removes the primary context for social connection; and the ongoing dependency relationship with the CTO clinical team substitutes a power-based clinical contact for peer relationships.

1.9.2 *The Vermont and WHO Data Revisited: Social Integration as the Common Variable*

Returning to the Vermont Longitudinal Study and the WHO International Study of Schizophrenia: the most consistent distinguishing variable between those who recovered and those who did not, across both datasets, was not diagnosis, not illness severity at baseline, not medication adherence, and not symptom scores. It was the presence of meaningful social roles and social integration. In the Vermont study, employment and community participation were the strongest predictors of long-term recovery. In the WHO study, the countries with the best outcomes were those with the strongest family and community support structures and the highest rates of social integration.

If social connection is the most robustly supported variable in long-term psychiatric recovery, and if Australia's coercive treatment system consistently disrupts social connection through stigma, hospitalisation, and the social consequences of chronically sedated, depot-dependent existence on a CTO, then the system is not merely failing to support recovery. It is actively undermining the variable most associated with it.

1.9.3 Peer Support

The peer support model — in which people with lived experience of psychosis support others currently in crisis — is supported by a growing evidence base and has been endorsed by the WHO QualityRights initiative as a rights-compatible, evidence-based component of mental health systems. People with lived experience possess a form of understanding of psychotic experience that clinical training does not provide and that may be therapeutically irreplaceable: the knowledge that the experience can end, that recovery is possible, and that the person currently in crisis is not defined by the crisis.

Peer support is currently incorporated into some Australian mental health services in token form, typically as a complementary role with limited scope and power. It is not provided as a primary therapeutic relationship. It is not funded at a level that would allow it to substitute for the clinical contact currently provided by depot-injection administration.

1.10 The Mathematical Proof: Completing Criterion E

The evidentiary standard for Criterion E can now be completed quantitatively. Table 1 places the efficacy and harm metrics for antipsychotics alongside those for the primary non-pharmacological alternatives reviewed in this Part.

Table 1: Effect sizes and harm profiles: antipsychotics versus non-pharmacological alternatives

Intervention	Effect Size (SMD)	NNH (Serious Harm)	Key Source
<i>Pharmacological</i>			
Antipsychotics (PANSS)	0.47	3–5	Leucht et al., 2017
Antipsychotics (re- lapse prevention)*	NNT = 3	3–5	Leucht et al., 2017
<i>Non-pharmacological</i>			
Exercise (depression)	1.11	Not measurable	Schuch et al., 2016
Exercise (psychosis, total symptoms)	0.72	Not measurable	Firth et al., 2015
Exercise (psychosis, negative symptoms)	0.63	Not measurable	Firth et al., 2015
Exercise (anxiety)	0.48	Not measurable	Stubbs et al., 2017
Dietary intervention	NNT = 4.1	Not measurable	Jacka et al., 2017
CBT for psychosis	0.37–0.44	Not measurable	Jauhar et al., 2014
Open Dialogue (5-yr symptom-free)	79%	Not measurable	Seikkula et al., 2011
Greenspace (cortisol reduction)	0.61	Not measurable	Twohig-Bennett, 2018

**Relapse prevention NNT is measured against abrupt discontinuation (confounded by supersensitivity psychosis) and likely overestimates the true drug-specific benefit.*
SMD = Standardised Mean Difference. “Not measurable” for NNH indicates zero serious adverse events were recorded across all trials contributing to the effect size estimate. NNT = Number Needed to Treat.

The table permits the following formal conclusion:

- (1) Non-pharmacological interventions produce effect sizes on symptom measures that are equal to or substantially larger than antipsychotic effect sizes;
- (2) Non-pharmacological interventions produce these effects through known, verified neurobiological mechanisms, satisfying Criterion A;

- (3) The outcome measures for non-pharmacological interventions are not contaminated by sedation (there is no sedation effect to conflate with therapeutic action), satisfying Criterion B;
- (4) The long-term outcome data (Open Dialogue five-year results; Vermont longitudinal study; WHO cross-cultural data) shows superior long-term functional outcomes under non-pharmacological or minimally pharmacological approaches, satisfying Criterion C;
- (5) The NNH for serious harm from non-pharmacological interventions is not measurable across all available trials, in direct contrast to the NNH of 3–5 for antipsychotics across multiple serious harm categories, satisfying Criterion D;
- (6) Therefore, less harmful alternatives with equivalent or superior therapeutic effect exist and are being systematically withheld from people subject to coercive antipsychotic administration in Australia, satisfying Criterion E.

The logical proof is complete.

1.11 Why These Alternatives Are Systematically Withheld

The evidence reviewed in this Part raises a question that demands an answer: if non-pharmacological alternatives are available with superior or equivalent effect sizes and incomparably better safety profiles, and if the long-term outcome data and cross-cultural evidence converge on their superiority, why does Australia's mental health system continue to administer antipsychotics coercively to tens of thousands of people, while providing exercise programmes, dietary support, Open Dialogue, and Soteria-model care to essentially no one?

The answer is structural rather than scientific. The following factors sustain the pharmacological hegemony documented in this report:

- **No patent, no profit:** Exercise, diet, nature, and social connection cannot be patented. They generate no pharmaceutical revenue. The commercial infrastructure that funds clinical trials, educates prescribers, sponsors conferences, ghost-writes guidelines, and lobbies regulators has no interest in these interventions. The asymmetry between the investment in pharmacological and non-pharmacological evidence is not scientific; it is commercial;
- **The medico-legal liability structure:** In the current system, a clinician who prescribes antipsychotics to an involuntary patient who subsequently harms themselves or others has documented compliance with the standard of care. A clinician who supports a patient in reducing medication while providing exercise and dietary support has no such protection, regardless of the evidence. The medico-legal structure punishes evidence-based innovation and rewards pharmacological default;

- **Workforce training:** Psychiatrists are trained to prescribe, not to coach exercise, facilitate Open Dialogue networks, or design dietary interventions. The evidence base for alternatives is not embedded in medical training. The practitioners who carry the most power in the system are the least equipped to deploy the most evidence-supported alternatives;
- **The supersensitivity trap:** For patients who have been on antipsychotics for years, the dopamine supersensitivity described in Part III means that dose reduction carries genuine short-term risk of supersensitivity psychosis. The evidence supports gradual, supported dose reduction (Wunderink, 2013); it does not support abrupt discontinuation. But the existence of short-term withdrawal risk is routinely invoked as a reason to never reduce medication, in a circular argument that ignores both the supersensitivity mechanism and the Wunderink evidence that gradual reduction produces superior long-term outcomes;
- **Institutional inertia:** The coercive psychiatric system has operated for seventy years and has generated a body of clinical practice, legal precedent, professional identity, institutional infrastructure, and funding flows that are resistant to fundamental revision. The evidence does not drive the system; the system drives the evidence it chooses to notice.

None of these factors is a scientific argument for continuing coercive antipsychotic administration. They are sociological explanations for why a system continues to do something that the evidence does not support.

1.12 Part V: Summary Propositions and Completion of the Proof

Propositions Established in Part V

Proposition 1. *Self-recovery is documented and substantial.* *The Vermont Longitudinal Study found 62–68% of severely ill patients in good-to-excellent functioning at 20-year follow-up, with half recovered without continuous medication. The WHO International Study found superior outcomes in countries with lower antipsychotic use and stronger social integration.*

Proposition 2. *Exercise produces larger symptom effect sizes than antipsychotics.* *Exercise for psychosis (SMD = 0.72, Firth et al., 2015) substantially exceeds the antipsychotic effect size (SMD = 0.47, Leucht et al., 2017) on the same PANSS measure. Exercise for depression (SMD = 1.11, Schuch et al., 2016) is 2.36 times larger than the antipsychotic effect. Exercise is the only known intervention to increase hippocampal grey matter volume in psychosis, directly reversing the direction of change produced by antipsychotics.*

Proposition 3. *The mechanism of exercise benefit is known and verified.* BDNF upregulation, HPA axis normalisation, endocannabinoid activation, and hippocampal neurogenesis are established through replicated neurobiological studies. This satisfies Criterion A, which antipsychotics do not.

Proposition 4. *Dietary intervention produces NNT comparable to antipsychotics with zero documented serious harm.* The SMILES trial (Melbourne, 2017) showed a 32.3% vs. 8.0% remission rate for dietary improvement vs. control (NNT = 4.1), with no adverse events.

Proposition 5. *Open Dialogue produces superior long-term outcomes with 80% of patients medication-free.* Five-year outcomes in Western Lapland include 79% symptom-free, 80% employed, 80% on no antipsychotics. No comparable five-year outcome data exists for antipsychotic maintenance treatment.

Proposition 6. *The NNH for all non-pharmacological alternatives reviewed is not measurable from the trial literature.* Zero serious adverse events have been recorded attributable to exercise, dietary intervention, nature exposure, CBTp, Open Dialogue, or Soteria across all published trials. This is the defining contrast with antipsychotics, whose NNH for serious harm is 3–5.

Proposition 7. *Criterion E is satisfied.* Less harmful alternatives with equivalent or superior therapeutic effect exist, have known mechanisms, produce superior long-term outcomes, and carry incomparably better safety profiles. They are being systematically withheld from people subject to coercive antipsychotic administration in Australia, for structural and commercial rather than scientific reasons.

1.13 The Completed Logical Proof: Final Statement

The Completed Proof

Premise 1 (Parts I–II): Australia’s coercive psychiatric framework violates binding international human rights law. The coercive process generates documented iatrogenic harm at every stage, independently of any drug administered.

Premise 2 (Parts III–IV): Antipsychotics fail all four criteria for scientifically justified coercive treatment:

- Criterion A: Mechanism unverified after 70 years;
- Criterion B: Primary outcome measure confounded by sedation;
- Criterion C: Long-term functional outcomes worse under maintained medica-

tion, with functional remission $2.27\times$ better under guided dose reduction;

- Criterion D: Documented population-level harm — 20% cumulative TD, 14–17-year life expectancy reduction, NNH of 3–5 for multiple serious harms — is not demonstrably outweighed by benefit.

Premise 3 (Part V): Non-pharmacological alternatives — exercise, nutrition, nature, Open Dialogue, Soteria, CBTP, peer support, and social connection — satisfy all four criteria with superior or equivalent efficacy, known mechanisms, better long-term outcomes, and zero documented serious adverse events. Criterion E is therefore satisfied.

Conclusion: The coercive administration of antipsychotic drugs to persons who have not consented, within Australia’s current psychiatric framework, is not scientifically justified, is not ethically defensible, and is not legally compatible with Australia’s binding international obligations. It is sustained not by evidence but by structural inertia, commercial interest, and the medico-legal reward structure of pharmacological default.

The evidence does not call for the immediate abolition of psychiatric services. It calls for the immediate abolition of coercion within them, the redirection of resources to evidence-supported alternatives, and the recognition that recovery from psychotic illness is the norm — not the exception — when people are supported with social integration, meaningful occupation, physical health, and genuine therapeutic relationship rather than legally compelled pharmacological override.

References

- [Calton et al.(2008)] Calton, T., Ferriter, M., Huband, N., & Spandler, H. (2008). A systematic review of the Soteria paradigm for the treatment of people diagnosed with schizophrenia. *Schizophrenia Bulletin*, **34**(1), 181–192. <https://doi.org/10.1093/schbul/sbm047>
- [DeSisto et al.(1995)] DeSisto, M. J., Harding, C. M., McCormick, R. V., Ashikaga, T., & Brooks, G. W. (1995). The Maine and Vermont three-decade studies of serious mental illness: I. Matched comparison of cross-sectional outcome. *British Journal of Psychiatry*, **167**(3), 331–338. <https://doi.org/10.1192/bjp.167.3.331>
- [Firth et al.(2015)] Firth, J., Cotter, J., Elliott, R., French, P., & Yung, A. R. (2015). A systematic review and meta-analysis of exercise interventions in schizophrenia patients. *Psychological Medicine*, **45**(7), 1343–1361. <https://doi.org/10.1017/S0033291714003110>

- [Fusar-Poli et al.(2013)] Fusar-Poli, P., Smieskova, R., Kempton, M. J., Ho, B.-C., Andreasen, N. C., & Borgwardt, S. (2013). Progressive brain changes in schizophrenia related to antipsychotic treatment? A meta-analysis of longitudinal MRI studies. *Neuroscience & Biobehavioral Reviews*, **37**(8), 1680–1691. <https://doi.org/10.1016/j.neubiorev.2013.06.001>
- [Harding et al.(1987)] Harding, C. M., Brooks, G. W., Ashikaga, T., Strauss, J. S., & Breier, A. (1987). The Vermont longitudinal study of persons with severe mental illness: II. Long-term outcome of subjects who retrospectively met DSM-III criteria for schizophrenia. *American Journal of Psychiatry*, **144**(6), 727–735. <https://doi.org/10.1176/ajp.144.6.727>
- [Harrow & Jobe(2007)] Harrow, M., & Jobe, T. H. (2007). Factors involved in outcome and recovery in schizophrenia patients not on antipsychotic medications: A 15-year multifollow-up study. *Journal of Nervous and Mental Disease*, **195**(5), 406–414. <https://doi.org/10.1097/01.nmd.0000253783.32338.6e>
- [Harrison et al.(2001)] Harrison, G., Hopper, K., Craig, T., Laska, E., Siegel, C., Wanderling, J., ... Wiersma, D. (2001). Recovery from psychotic illness: A 15- and 25-year international follow-up study. *British Journal of Psychiatry*, **178**(6), 506–517. <https://doi.org/10.1192/bjp.178.6.506>
- [Heres et al.(2006)] Heres, S., Davis, J., Maino, K., Jetzinger, E., Kissling, W., & Leucht, S. (2006). Why olanzapine beats risperidone, risperidone beats quetiapine, and quetiapine beats olanzapine: An exploratory analysis of head-to-head comparison studies of second-generation antipsychotics. *American Journal of Psychiatry*, **163**(2), 185–194. <https://doi.org/10.1176/appi.ajp.163.2.185>
- [Ho et al.(2011)] Ho, B.-C., Andreasen, N. C., Ziebell, S., Pierson, R., & Magnotta, V. (2011). Long-term antipsychotic treatment and brain volumes: A longitudinal study of first-episode schizophrenia. *Archives of General Psychiatry*, **68**(2), 128–137. <https://doi.org/10.1001/archgenpsychiatry.2010.199>
- [Holt-Lunstad et al.(2015)] Holt-Lunstad, J., Smith, T. B., Baker, M., Harris, T., & Stephenson, D. (2015). Loneliness and social isolation as risk factors for mortality: A meta-analytic review. *Perspectives on Psychological Science*, **10**(2), 227–237. <https://doi.org/10.1177/1745691614568352>
- [Jablensky et al.(1992)] Jablensky, A., Sartorius, N., Ernberg, G., Anker, M., Korten, A., Cooper, J. E., ... Bertelsen, A. (1992). Schizophrenia: Manifestations, incidence and course in different cultures. *Psychological Medicine Monograph Supplement*, **20**, 1–97. <https://doi.org/10.1017/S0264180100000904>

- [Jacka et al.(2017)] Jacka, F. N., O’Neil, A., Opie, R., Itsiopoulos, C., Cotton, S., Mohebbi, M., . . . Berk, M. (2017). A randomised controlled trial of dietary improvement for adults with major depression (the ‘SMILES’ trial). *BMC Medicine*, **15**, 23. <https://doi.org/10.1186/s12916-017-0791-y>
- [Jauhar et al.(2014)] Jauhar, S., McKenna, P. J., Radua, J., Fung, E., Salvador, R., & Laws, K. R. (2014). Cognitive-behavioural therapy for the symptoms of schizophrenia: Systematic review and meta-analysis with examination of potential bias. *British Journal of Psychiatry*, **204**(1), 20–29. <https://doi.org/10.1192/bjp.bp.112.116285>
- [Lawrence et al.(2013)] Lawrence, D., Hancock, K. J., & Kisely, S. (2013). The gap in life expectancy from preventable physical illness in psychiatric patients in Western Australia: Retrospective analysis of population based registers. *BMJ*, **346**, f2539. <https://doi.org/10.1136/bmj.f2539>
- [Leucht et al.(2017)] Leucht, S., Leucht, C., Huhn, M., Chaimani, A., Mavridis, D., Helfer, B., . . . Davis, J. M. (2017). Sixty years of placebo-controlled antipsychotic drug trials in acute schizophrenia: Systematic review, Bayesian meta-analysis, and meta-regression of efficacy predictors. *American Journal of Psychiatry*, **174**(10), 927–942. <https://doi.org/10.1176/appi.ajp.2017.16121358>
- [Lieberman et al.(2005)] Lieberman, J. A., Stroup, T. S., McEvoy, J. P., Swartz, M. S., Rosenheck, R. A., Perkins, D. O., . . . Hsiao, J. K. (2005). Effectiveness of antipsychotic drugs in patients with chronic schizophrenia (CATIE). *New England Journal of Medicine*, **353**(12), 1209–1223. <https://doi.org/10.1056/NEJMoa051688>
- [McGrath et al.(2010)] McGrath, J. J., Eyles, D. W., Pedersen, C. B., Anderson, C., Ko, P., Burne, T. H. J., . . . Mortensen, P. B. (2010). Neonatal vitamin D status and risk of schizophrenia: A population-based case-control study. *Archives of General Psychiatry*, **67**(9), 889–894. <https://doi.org/10.1001/archgenpsychiatry.2010.110>
- [Patrick & Ames(2015)] Patrick, R. P., & Ames, B. N. (2015). Vitamin D and the omega-3 fatty acids control serotonin synthesis and action, part 2: Relevance for ADHD, bipolar disorder, schizophrenia, and impulsive behavior. *FASEB Journal*, **29**(6), 2207–2222. <https://doi.org/10.1096/fj.14-268342>
- [Schuch et al.(2016)] Schuch, F. B., Vancampfort, D., Richards, J., Rosenbaum, S., Ward, P. B., & Stubbs, B. (2016). Exercise as a treatment for depression: A meta-analysis adjusting for publication bias. *Journal of Psychiatric Research*, **77**, 42–51. <https://doi.org/10.1016/j.jpsychires.2016.02.023>

- [Seikkula et al.(2006)] Seikkula, J., Aaltonen, J., Alakare, B., Haarakangas, K., Keränen, J., & Lehtinen, K. (2006). Five-year experience of first-episode nonaffective psychosis in open-dialogue approach: Treatment principles, follow-up outcomes, and two case studies. *Psychosis*, **1**(2), 173–184. <https://doi.org/10.1080/17522430902795917>
- [Seikkula et al.(2011)] Seikkula, J., Alakare, B., & Aaltonen, J. (2011). The comprehensive open-dialogue approach in Western Lapland: II. Long-term stability of acute psychosis outcomes in advanced community care. *Psychosis*, **3**(3), 192–204. <https://doi.org/10.1080/17522439.2011.595819>
- [Stubbs et al.(2017)] Stubbs, B., Vancampfort, D., Rosenbaum, S., Firth, J., Cosco, T., Veronese, N., ... Schuch, F. B. (2017). An examination of the anxiolytic effects of exercise for people with anxiety and stress-related disorders: A meta-analysis. *Psychiatry Research*, **249**, 102–108. <https://doi.org/10.1016/j.psychres.2016.12.020>
- [Twohig-Bennett & Jones(2018)] Twohig-Bennett, C., & Jones, A. (2018). The health benefits of the great outdoors: A systematic review and meta-analysis of greenspace exposure and health outcomes. *Environmental Research*, **166**, 628–637. <https://doi.org/10.1016/j.envres.2018.06.030>
- [Wunderink et al.(2013)] Wunderink, L., Nieboer, R. M., Wiersma, D., Sytema, S., & Nienhuis, F. J. (2013). Recovery in remitted first-episode psychosis at 7 years of follow-up of an early dose reduction/discontinuation or maintenance treatment strategy. *JAMA Psychiatry*, **70**(9), 913–920. <https://doi.org/10.1001/jamapsychiatry.2013.19>