

CS 260: Foundations of Data Science

Prof. Thao Nguyen

Fall 2024



HAVERFORD
COLLEGE

Admin

- Lab 4 due tonight



Admin

- **Lab 3** grades & feedback will be posted on Wednesday
- **Midterm 1** will be handed out on Wednesday (due the following Wednesday – take in a 3 hour block)
- **Tuesday + Wednesday**: review sessions

Midterm 1 Notes

- Handed out in class this Wednesday, due at the beginning of class the following Wednesday.
- Timed exam: **3 hour limit**. DO NOT open the exam until you are ready to take it for 3 hours!
- You may use a one page (front and back) “study sheet”, handwritten, created by you
- You may also use a regular calculator
- Outside of your “study sheet” and calculator, **no other notes or resources**
- **As per the Honor Code, all work must be your own**

Outline for today

- Go over Lab 2
- Intro to probability
- Intro to Bayesian models

Outline for today

- Go over Lab 2
- Intro to probability
- Intro to Bayesian models

Lab 2: not posted online

Outline for today

- Go over Lab 2
- **Intro to probability**
- Intro to Bayesian models

Intro to Probability

- The **probability** of an **event** e has a number of epistemological interpretations
- Assuming we have **data**, we can count the number of times e occurs in the dataset to estimate the probability of e , $P(e)$.

$$P(e) = \frac{\text{count}(e)}{\text{count}(\text{all events})}.$$

- If we put all events in a bag, shake it up, and choose one at random (called **sampling**), how likely are we to get e ?

Intro to Probability



- Suppose we have a fair 6-sided die.
- What's the probability of getting "1"?

$$\frac{\textit{count}(s)}{\textit{count}(1) + \textit{count}(2) + \textit{count}(3) + \cdots + \textit{count}(6)} = \frac{1}{1 + 1 + 1 + 1 + 1 + 1} = \frac{1}{6}$$

Intro to Probability



- What about a die with only three numbers $\{1, 2, 3\}$, each of which appears twice?
- What's the probability of getting "1"?

Intro to Probability



- What about a die with only three numbers $\{1, 2, 3\}$, each of which appears twice?
- What's the probability of getting "1"?

$$P(e = 1) = \frac{\text{count}(1)}{\text{count}(1) + \text{count}(2) + \text{count}(3)} = \frac{2}{2 + 2 + 2} = \frac{1}{3}.$$

Intro to Probability



- The set of all probabilities for an event e is called a **probability distribution**
- Each coin toss is an independent event (Bernoulli trial).

Intro to Probability



- Which is greater, $P(HHHHHH)$ or $P(HHTHHH)$?

Intro to Probability



- Which is greater, $P(HHHHHH)$ or $P(HHTHHH)$?
- Since the events are independent, they're equal

Intro to Probability

Probability Axioms

1. Probabilities of events must be no less than 0. $P(e) \geq 0$ for all e .
2. The sum of all probabilities in a distribution must sum to 1. That is,
 $P(e_1) + P(e_2) + \dots + P(e_n) = 1$. Or, more succinctly,

$$\sum_{e \in E} P(e) = 1.$$

Intro to Probability

Joint Probability


The probability that two independent events e_1 and e_2 *both* occur is given by their product.

$$P(e_1 \wedge e_2) = P(e_1 \cap e_2) = P(e_1)P(e_2) \text{ when } e_1 \cap e_2 = \emptyset$$

- Intuitively, think of every probability as a *scaling factor*.
- You can think of a probability as the fraction of the probability space occupied by an event e_1 .
 - $P(e_1 \wedge e_2)$ is the fraction of e_1 's probability space wherein e_2 also occurs.
 - So, if $P(e_1) = \frac{1}{2}$ and $P(e_2) = \frac{1}{3}$, then $P(e_2, e_1)$ is a third of a half of the probability space or $\frac{1}{3} \times \frac{1}{2}$.

Intro to Probability

Joint Probability




Intro to Probability

Conditional Probability

- A **conditional probability** is the probability that one event occurs given that we take another for granted.
- The probability of e_2 given e_1 is $P(e_2 \mid e_1)$.
- This is the probability that e_2 will occur given that we take for granted that e_1 occurs.

Intro to Probability

Conditional Probability




example $R = \text{rain}$
 $U = \text{umbrella}$

If $P(R) = 20\%$ and
 $P(R, U) = 15\%$, and (joint prob)

what is $\underline{P(U|R)}$?
given
(conditional probability)

Bayes rule


$$P(U|R) = \frac{P(R, U)}{P(R)} = \frac{0.15}{0.2} = \boxed{0.75}$$


$\boxed{\text{diagonal}} = 0.2$
 $\boxed{\text{diagonal}} = 0.15$
 $P(R, U)$

Bayes' Theorem

- $P(A,B) = P(A | B)P(B)$
- $P(A,B) = P(B | A)P(A)$

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

Independence: $P(A,B) = P(A)P(B)$  not true in general!!!

Conditional Independence

$$P(A | B, C) = P(A | C)$$

↑ ↑ ↑
thunder rain lightning

"Thunder is independent of rain
give lightning."

$P(B|A)$

~~$P(A|B) = P(A)$~~ } standard independence

Intro to Probability

Marginal Probability Distributions

Given a discrete joint probability distribution function $P(X, Y)$, how would we find $P(X)$?

- "Marginalize out" the Y (sum over all $y \in Y$).
- Discrete Case: $p(x) = \sum_{y \in Y} P(x, y)$
- Continuous Case: $p(x) = \int p(x, y) dy$

Marginalizing

$$P(A) = \sum_{b \in \text{val}(B)} P(A, B=b)$$

$$P(u) = P(R, u) + P(\bar{R}, u)$$

Example

$$P(\text{spam} | \underbrace{\text{words}}_{\text{email}}) = \frac{p(\text{spam}, \text{words})}{\underbrace{p(\text{words})}_{\text{very difficult!}}}$$

posterior

"data"

$$p(\text{spam}, \text{words})$$

$$p(\text{spam}, \text{words}) + p(\bar{\text{spam}}, \text{words})$$

$$= \frac{p(\text{spam}) p(\text{words} | \text{spam})}{p(\text{spam}) p(\text{words} | \text{spam}) + p(\bar{\text{spam}}) p(\text{words} | \bar{\text{spam}})}$$

Prior

evidence

likelihood
(generative)

X

Handout 9

Handout 9

①

$$P(A|B) = \frac{P(A, B)}{P(B)} = \frac{P(A)P(B|A)}{P(B)}$$

$$P(D|pos) = \frac{\overbrace{P(D)}^{\text{prior}} \overbrace{P(pos|D)}^{\text{likelihood}}}{\underbrace{P(pos)}_{\text{evidence}}}$$

$$= \frac{P(D)P(pos|D)}{P(D)P(pos|D) + \underbrace{P(H)P(pos|H)}}_{\text{evidence}}$$

neg pos

true H -

9/10	1/10
1/10	9/10

D +

$$\frac{\frac{1}{100} \cdot \frac{9}{10}}{\frac{1}{100} \cdot \frac{9}{10} + \frac{99}{100} \cdot \frac{1}{10}} = \frac{9}{9 + 99} = \frac{9}{108}$$

$$= \frac{1}{12}$$

≈ 0.0825 8%

Outline for today

- Go over Lab 2
- Intro to probability
- **Intro to Bayesian models**


Components of a Bayesian Model

- Identify the evidence, prior, posterior, and likelihood in the equation below

$$p(y = k|\mathbf{x}) = \frac{p(y = k)p(\mathbf{x}|y = k)}{p(\mathbf{x})}$$

Components of a Bayesian Model

- Identify the evidence, prior, posterior, and likelihood in the equation below

$$p(y = k|\mathbf{x}) = \frac{p(y = k)p(\mathbf{x}|y = k)}{p(\mathbf{x})}$$


- Evidence:** this is the data (features) we actually observe, which we think will help us predict the outcome we're interested in

Components of a Bayesian Model

- Identify the evidence, prior, posterior, and likelihood in the equation below

$$p(y = k|\mathbf{x}) = \frac{p(y = k)p(\mathbf{x}|y = k)}{p(\mathbf{x})}$$

- Prior**: without seeing any evidence (data), what is our prior believe about each outcome (intuition: what is the outcome in the population as a whole?)

Components of a Bayesian Model

- Identify the evidence, prior, posterior, and likelihood in the equation below

$$p(y = k|\mathbf{x}) = \frac{p(y = k)p(\mathbf{x}|y = k)}{p(\mathbf{x})}$$

- Posterior**: this is the quantity we are actually interested in. **Given** the evidence, what is the probability of the outcome?

Components of a Bayesian Model

- Identify the evidence, prior, posterior, and likelihood in the equation below

$$p(y = k|\mathbf{x}) = \frac{p(y = k)p(\mathbf{x}|y = k)}{p(\mathbf{x})}$$

- Likelihood**: given an outcome, what is the probability of observing this set of features?

Examples

- Computing the probability an email message is **spam**, given the **words** of the email
- Another example: what is the probability of **Trisomy 21** (Down Syndrome), given the **amount of sequencing of each chromosome?**

Bayesian Model for Trisomy 21 (T_{21})

Input data are read counts for each chromosome (1,2,...,n):

$$q_1, q_2, \dots, q_n = \vec{q}$$

Bayesian Model for Trisomy 21 (T_{21})

Input data are read counts for each chromosome (1,2,...,n):

$$q_1, q_2, \dots, q_n = \vec{q}$$

Goal:

$$\begin{aligned}\mathbb{P}(T_{21}|\vec{q}) &= \frac{\mathbb{P}(\vec{q} | T_{21}) \cdot \mathbb{P}(T_{21})}{\mathbb{P}(\vec{q})} \\ &= \frac{\mathbb{P}(\vec{q} | T_{21}) \cdot \mathbb{P}(T_{21})}{\mathbb{P}(\vec{q} | T_{21}) \cdot \mathbb{P}(T_{21}) + \mathbb{P}(\vec{q} | T_{21}^C) \cdot \mathbb{P}(T_{21}^C)}\end{aligned}$$

Bayesian Model for Trisomy 21 (T_{21})

Input data are read counts for each chromosome (1,2,...,n):

$$q_1, q_2, \dots, q_n = \vec{q}$$

Goal:

$$\mathbb{P}(T_{21} | \vec{q}) = \frac{\mathbb{P}(\vec{q} | T_{21}) \cdot \mathbb{P}(T_{21})}{\mathbb{P}(\vec{q})}$$

Prior probability of T_{21}

$$= \frac{\mathbb{P}(\vec{q} | T_{21}) \cdot \mathbb{P}(T_{21})}{\mathbb{P}(\vec{q} | T_{21}) \cdot \mathbb{P}(T_{21}) + \mathbb{P}(\vec{q} | T_{21}^C) \cdot \mathbb{P}(T_{21}^C)}$$

Prior:

$P(T_{21})$

Maternal Age	Trisomy 21	All Trisomies
20	1 in 1,667	1 in 526
21	1 in 1,429	1 in 526
22	1 in 1,429	1 in 500
23	1 in 1,429	1 in 500
24	1 in 1,250	1 in 476
25	1 in 1,250	1 in 476
26	1 in 1,176	1 in 476
27	1 in 1,111	1 in 455
28	1 in 1,053	1 in 435
29	1 in 1,000	1 in 417
30	1 in 952	1 in 384
31	1 in 909	1 in 384
32	1 in 769	1 in 323
33	1 in 625	1 in 286
34	1 in 500	1 in 238
35	1 in 385	1 in 192
36	1 in 294	1 in 156
37	1 in 227	1 in 127
38	1 in 175	1 in 102
39	1 in 137	1 in 83
40	1 in 106	1 in 66
41	1 in 82	1 in 53
42	1 in 64	1 in 42
43	1 in 50	1 in 33
44	1 in 38	1 in 26
45	1 in 30	1 in 21
46	1 in 23	1 in 16
47	1 in 18	1 in 13
48	1 in 14	1 in 10
49	1 in 11	1 in 8

Anonymous feedback form