

# Assignment 2

David Gyarakı, Thao Le

## Contents

<b>1</b>	<b>Question 1</b>	<b>2</b>
1.1	(i) . . . . .	2
1.2	(ii) . . . . .	4
1.3	(iii) . . . . .	8
<b>2</b>	<b>Question 2</b>	<b>14</b>
2.1	(i) . . . . .	14

```
# load packages
if(!require(pacman)){install.packages("pacman")}

p_load(devtools,tidyverse,dplyr,ggplot2,latex2exp,
       sampleSelection, quantreg)


#load data
dfData = read.csv("assignment2a_2023.csv")
attach(dfData)
```

## 1 Question 1

### 1.1 (i)


```
# Get the quantile values
quant=quantile(lntotexp, seq(0.1, 0.9, by=.4))
n = length(lntotexp)

# Histogram of log of total medical expenditure
hist(lntotexp)
```



```
# Quantile plot of log of total medical expenditure
plot((1:n - 1)/(n - 1), sort(lntotexp), type="l",
main = "Quantiles for log of total medical expenditure",
xlab = "Sample Fraction",
ylab = "Sample Quantile") + abline(h=quant, col = c("dark green","red",
↪ "blue"))
```

## Quantiles for log of total medical expenditure



`integer(0)`

In the quantile plot, the median is indicated by the red line, the 10<sup>th</sup> and 90<sup>th</sup> quantile are indicated by the blue and green lines.

We can see from the distribution of log of total medical expenditure that there are few values from 0 to 4. Thus, the quantile plot increases quickly in this region. From 4 to 6, we see an increase frequencies of observations, thus, the quantile plot increases slower. The most rapid increase in the quantile plot is observed between 6 and 10, which makes sense because that is the region where most observations lie. After 10, there are less observations and the quantile plot increases rapidly again.

Although the quantile plot increases rapidly in both regions from 0 to 4 and 10 to 12, we observed a much steeper increase from 0 to 4, thus, we can say that the distribution of log total medical expenditure is left-skewed. This is confirmed by looking at its histogram.

### 1.2 (ii)

```
# Quantile regression
q= c(0.1,0.25,0.5,0.75,0.9)
quant_reg = rq(lntotexp ~ . , tau = q, data = dfData)
```

```
summary(quant_reg)
```

```
Call: rq(formula = lntotexp ~ ., tau = q, data = dfData)
```

```
tau: [1] 0.1
```

```
Coefficients:
```

	Value	Std. Error	t value	Pr(> t )
(Intercept)	3.86704	0.48065	8.04549	0.00000
age	0.01927	0.00601	3.20732	0.00135
female	-0.01273	0.07579	-0.16794	0.86664
white	0.07344	0.19533	0.37597	0.70697
totchr	0.53919	0.02534	21.27920	0.00000
suppins	0.39572	0.07851	5.04027	0.00000

```
Call: rq(formula = lntotexp ~ ., tau = q, data = dfData)
```

```
tau: [1] 0.25
```

```
Coefficients:
```

	Value	Std. Error	t value	Pr(> t )
(Intercept)	4.74732	0.30724	15.45160	0.00000
age	0.01551	0.00399	3.88410	0.00010
female	-0.01623	0.05328	-0.30462	0.76068
white	0.33775	0.09662	3.49570	0.00048
totchr	0.45918	0.01833	25.04804	0.00000
suppins	0.38584	0.05992	6.43964	0.00000

```
Call: rq(formula = lntotexp ~ ., tau = q, data = dfData)
```

```
tau: [1] 0.5
```

```
Coefficients:
```

	Value	Std. Error	t value	Pr(> t )
(Intercept)	5.61116	0.35187	15.94656	0.00000
age	0.01487	0.00406	3.66512	0.00025
female	-0.08810	0.05406	-1.62961	0.10329
white	0.53648	0.19319	2.77697	0.00552
totchr	0.39427	0.01846	21.35942	0.00000
suppins	0.27698	0.05347	5.18025	0.00000

```
Call: rq(formula = lntotexp ~ ., tau = q, data = dfData)
```

```
tau: [1] 0.75
```

```
Coefficients:
```

	Value	Std. Error	t value	Pr(> t )
(Intercept)	6.59997	0.42690	15.46027	0.00000
age	0.01825	0.00475	3.83862	0.00013
female	-0.12194	0.06060	-2.01231	0.04428
white	0.19319	0.25684	0.75219	0.45200
totchr	0.37354	0.02286	16.33884	0.00000
suppins	0.14885	0.06203	2.39991	0.01646

```
Call: rq(formula = lntotexp ~ ., tau = q, data = dfData)
```

```
tau: [1] 0.9
```

```
Coefficients:
```

	Value	Std. Error	t value	Pr(> t )
(Intercept)	8.32264	0.54599	15.24309	0.00000
age	0.00592	0.00651	0.91022	0.36278
female	-0.15763	0.08914	-1.76831	0.07711
white	0.30522	0.24260	1.25811	0.20845
totchr	0.35795	0.03310	10.81289	0.00000
suppins	-0.01428	0.08642	-0.16527	0.86874

Looking at the results, we observe different coefficients across the different quantiles. Quite expectedly, we have increasing intercept coefficients, however the interesting part is the different significance of the coefficients in the different quantile regressions. We observe that for the 0.1 quantile, the female and white dummies are insignificant, for the 0.25 and 0.5 quantiles only the female dummy is insignificant, for the 0.75, interestingly the white dummy is insignificant while the female dummy turns out to be significant, and for the 0.9 quantile, only the chronic illness variable seems to be strongly significant with the female dummy slightly (at 10% level) significant too. These trends will lead to the conclusion that the different predictors likely have different dynamics across the groups of patients when ordered by medical expenditure. Being white significantly increases medical expenditure in the mid-groups but not in the tails of the expenditure distribution. Age and extra insurance are associated with significant increase in costs for low spending groups but not for the highest spenders, and gender comes into influence for the highest spenders only. Let us then look at the OLS results, coefficients and their significance levels.

```
# OLS Regression
OLS_reg = lm(lntotexp ~ . , data = dfData)
summary(OLS_reg)
```

Call:

```
lm(formula = lntotexp ~ . , data = dfData)
```

Residuals:

Min	1Q	Median	3Q	Max
-6.2474	-0.7666	-0.0032	0.7827	3.8516

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )	
(Intercept)	5.898155	0.295694	19.947	< 2e-16	***
age	0.012656	0.003595	3.520	0.000437	***
female	-0.076517	0.046110	-1.659	0.097132	.
white	0.317811	0.141360	2.248	0.024635	*
totchr	0.445272	0.017549	25.374	< 2e-16	***
suppins	0.256811	0.046450	5.529	3.51e-08	***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.227 on 2949 degrees of freedom

Multiple R-squared: 0.1969, Adjusted R-squared: 0.1955

F-statistic: 144.6 on 5 and 2949 DF, p-value: < 2.2e-16

When one looks at the OLS regression results, the model shows that most variables are statistically significant for explaining the logarithm of medical expenditure, except for the female dummy variable. The variables *age*, *totchr* and *suppins* all have positive effect on medical expenditure with less than 0.001 significance, and the variable *white* has a positive effect as well on 5% significance level. The interpretation of the coefficients can also be given as one unit increase in the independent variables (keeping all else equal) increases the medical expenditure by  $(\exp(\beta_k) - 1) * 100$  percentage. We can see below, that a year of age increase will result in an estimated 1.274% increase in medical expenses. Similarly, being female reduces the expenses by -7.366% (although this is only significant at 10% level in the OLS model), being white is associated with 37.412% increase in medical expenses, an additional chronic illness will increase expenditure by 56.091% and having a supplementary private insurance will result in 29.280% increase in medical expenses.

```
(exp(OLS_reg$coefficients)-1)*100
```

(Intercept)	age	female	white	totchr	suppins
36336.450509	1.273661	-7.366254	37.411599	56.091416	29.280045

### 1.3 (iii)

```
# Quantile regression in increments of 0.05
q_005 = seq(0.05, 0.95, length.out=19)
quant_reg_005 = rq(lntotexp ~ ., tau = q_005, data = dfData)
summary(quant_reg_005)
```

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.05

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	3.36557	0.68439	4.91765	0.00000
age	0.01977	0.00893	2.21353	0.02694
female	0.12068	0.10803	1.11704	0.26407
white	-0.23365	0.23069	-1.01282	0.31123
totchr	0.63345	0.02977	21.27576	0.00000
suppins	0.41912	0.11495	3.64608	0.00027

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.1

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	3.86704	0.48065	8.04549	0.00000
age	0.01927	0.00601	3.20732	0.00135
female	-0.01273	0.07579	-0.16794	0.86664
white	0.07344	0.19533	0.37597	0.70697
totchr	0.53919	0.02534	21.27920	0.00000
suppins	0.39572	0.07851	5.04027	0.00000

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.15



Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	4.15640	0.41748	9.95605	0.00000
age	0.01865	0.00537	3.47031	0.00053
female	0.02271	0.07068	0.32138	0.74795
white	0.15737	0.13749	1.14459	0.25247
totchr	0.51204	0.02432	21.05569	0.00000
suppins	0.39942	0.06989	5.71491	0.00000

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.2

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	4.47890	0.34615	12.93916	0.00000
age	0.01734	0.00454	3.81746	0.00014
female	-0.01323	0.06120	-0.21618	0.82886
white	0.25032	0.09454	2.64763	0.00815
totchr	0.48030	0.02012	23.86793	0.00000
suppins	0.40203	0.06042	6.65370	0.00000

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.25

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	4.74732	0.30724	15.45160	0.00000
age	0.01551	0.00399	3.88410	0.00010
female	-0.01623	0.05328	-0.30462	0.76068
white	0.33775	0.09662	3.49570	0.00048
totchr	0.45918	0.01833	25.04804	0.00000
suppins	0.38584	0.05992	6.43964	0.00000

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.3

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	5.18763	0.32873	15.78085	0.00000
age	0.01207	0.00428	2.82053	0.00483

female	-0.03342	0.05733	-0.58296	0.55996
white	0.47252	0.07958	5.93801	0.00000
totchr	0.42963	0.01802	23.84426	0.00000
suppins	0.28488	0.05991	4.75485	0.00000

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.35

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	5.14852	0.32570	15.80777	0.00000
age	0.01458	0.00420	3.46956	0.00053
female	-0.06382	0.05469	-1.16706	0.24328
white	0.52359	0.12196	4.29323	0.00002
totchr	0.41297	0.01906	21.66773	0.00000
suppins	0.29115	0.05391	5.40044	0.00000

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.4

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	5.34247	0.34784	15.35906	0.00000
age	0.01400	0.00414	3.38472	0.00072
female	-0.08100	0.05366	-1.50939	0.13131
white	0.54055	0.17574	3.07593	0.00212
totchr	0.41102	0.01960	20.97561	0.00000
suppins	0.28977	0.05397	5.36882	0.00000

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.45

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	5.53579	0.35381	15.64622	0.00000
age	0.01411	0.00407	3.46239	0.00054
female	-0.06450	0.05189	-1.24309	0.21393
white	0.49315	0.19768	2.49466	0.01266
totchr	0.40721	0.01893	21.50765	0.00000
suppins	0.25994	0.05275	4.92812	0.00000

```
Call: rq(formula = lntotexp ~ ., tau = q_005, data = dfData)
```

```
tau: [1] 0.5
```

```
Coefficients:
```

	Value	Std. Error	t value	Pr(> t )
(Intercept)	5.61116	0.35187	15.94656	0.00000
age	0.01487	0.00406	3.66512	0.00025
female	-0.08810	0.05406	-1.62961	0.10329
white	0.53648	0.19319	2.77697	0.00552
totchr	0.39427	0.01846	21.35942	0.00000
suppins	0.27698	0.05347	5.18025	0.00000

```
Call: rq(formula = lntotexp ~ ., tau = q_005, data = dfData)
```

```
tau: [1] 0.55
```

```
Coefficients:
```

	Value	Std. Error	t value	Pr(> t )
(Intercept)	5.82910	0.39492	14.76022	0.00000
age	0.01416	0.00407	3.48048	0.00051
female	-0.09861	0.05257	-1.87593	0.06076
white	0.54989	0.26352	2.08671	0.03700
totchr	0.38758	0.01961	19.76391	0.00000
suppins	0.23471	0.05495	4.27124	0.00002

```
Call: rq(formula = lntotexp ~ ., tau = q_005, data = dfData)
```

```
tau: [1] 0.6
```

```
Coefficients:
```

	Value	Std. Error	t value	Pr(> t )
(Intercept)	6.15907	0.44080	13.97262	0.00000
age	0.01506	0.00420	3.58836	0.00034
female	-0.10853	0.05583	-1.94395	0.05200
white	0.25683	0.31863	0.80602	0.42030
totchr	0.39562	0.02031	19.47553	0.00000
suppins	0.25798	0.05577	4.62590	0.00000

```
Call: rq(formula = lntotexp ~ ., tau = q_005, data = dfData)
```

```
tau: [1] 0.65
```

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	6.36258	0.40648	15.65275	0.00000
age	0.01487	0.00461	3.22352	0.00128
female	-0.12887	0.05958	-2.16293	0.03063
white	0.28299	0.23108	1.22462	0.22082
totchr	0.38288	0.02194	17.44947	0.00000
suppins	0.20693	0.06372	3.24745	0.00118

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.7

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	6.63358	0.40368	16.43281	0.00000
age	0.01444	0.00478	3.02030	0.00255
female	-0.12951	0.05988	-2.16259	0.03065
white	0.27653	0.21300	1.29828	0.19429
totchr	0.37716	0.02214	17.03824	0.00000
suppins	0.15564	0.06329	2.45903	0.01399

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.75

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	6.59997	0.42690	15.46027	0.00000
age	0.01825	0.00475	3.83862	0.00013
female	-0.12194	0.06060	-2.01231	0.04428
white	0.19319	0.25684	0.75219	0.45200
totchr	0.37354	0.02286	16.33884	0.00000
suppins	0.14885	0.06203	2.39991	0.01646

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.8

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	6.90999	0.36065	19.15991	0.00000

age	0.01785	0.00471	3.78762	0.00016
female	-0.15788	0.06144	-2.56945	0.01023
white	0.13863	0.11657	1.18927	0.23443
totchr	0.38143	0.02285	16.69225	0.00000
suppins	0.11425	0.06222	1.83630	0.06641

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.85

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	7.31366	0.46945	15.57926	0.00000
age	0.01407	0.00590	2.38227	0.01727
female	-0.18200	0.07945	-2.29064	0.02205
white	0.28563	0.16208	1.76226	0.07813
totchr	0.36909	0.02806	13.15508	0.00000
suppins	0.10036	0.08226	1.21999	0.22257

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)

tau: [1] 0.9

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	8.32264	0.54599	15.24309	0.00000
age	0.00592	0.00651	0.91022	0.36278
female	-0.15763	0.08914	-1.76831	0.07711
white	0.30522	0.24260	1.25811	0.20845
totchr	0.35795	0.03310	10.81289	0.00000
suppins	-0.01428	0.08642	-0.16527	0.86874

Call: rq(formula = lntotexp ~ ., tau = q\_005, data = dfData)


tau: [1] 0.95

Coefficients:

	Value	Std. Error	t value	Pr(> t )
(Intercept)	9.74213	0.57059	17.07369	0.00000
age	-0.00606	0.00560	-1.08127	0.27967
female	-0.25712	0.07341	-3.50255	0.00047
white	0.40026	0.36872	1.08554	0.27777
totchr	0.31566	0.02827	11.16644	0.00000

suppins       -0.04675   0.07189   -0.65032   0.51553

```
quantile_coeffs <- as.data.frame(quant_reg_005[["coefficients"]])  
#print(predict.rq(quant_reg_005, interval = c("confidence")))  
  
plot.rqs(quant_reg_005, ols=TRUE)
```



## 2 Question 2

### 2.1 (i)

When one takes the observation relative to the individual-level mean, we include the information present in all of the observations belonging to one panel group or time horizon. In this case, each observation's fitted value will consider information from the individual groups, thus controlling for group/time fixed effects.