

# THE METRIC SPACE OF CONNECTED SIMPLE GRAPHS

Kiran R. Bhutani

Bilal Khan

## Abstract

We introduce a sequence  $e_n^\ell$  ( $\ell = 1, 2, \dots$ ) of real-valued functions on  $\mathcal{G}_n \times \mathcal{G}_n$ , where  $\mathcal{G}_n$  is the set of all simple, connected, undirected graphs of order  $n$  up to isomorphism. The functions  $e_n^\ell$  arise naturally from the consideration of graph embeddings. The binary relations  $\succ_\ell$  on  $\mathcal{G}_n$  are then defined to be exactly the zeroes of  $e_n^\ell$ . We observe that the relation  $\succ_1$  coincides precisely with the classical subgraph relation, and that  $\succ_1, \succ_2, \dots$  is a sequence of weaker binary relations on  $\mathcal{G}_n$ . Motivated by this, we define  $d_n^\ell(H, K) = e_n^\ell(H, K) + e_n^\ell(K, H)$ , thereby obtaining a sequence of symmetric, real-valued functions  $d_n^\ell$  ( $\ell = 1, 2, \dots$ ) on  $\mathcal{G}_n \times \mathcal{G}_n$ . When  $\ell = 1$ ,  $(\mathcal{G}_n, d_n^1)$  is a totally disconnected metric space of graphs that embodies the classical notion of graph isomorphism:  $H \cong K \Leftrightarrow d_n^1(H, K) = 0$ ,  $H \not\cong K \Leftrightarrow d_n^1(H, K) = \infty$ . As  $\ell$  increases in the range  $2, \dots, n-2$ , the value of  $d_n^\ell$  decreases for every distinct pair of graphs in  $\mathcal{G}_n$  although distinct graphs in  $\mathcal{G}_n$  do not collapse. Further, we show that the graphs  $H$  for which  $d_n^\ell(H, K_n)$  is finite, are precisely those graphs  $H \in \mathcal{G}_n$  which have diameter  $\leq \ell$ . Finally, when  $\ell \geq n-1$ , the functions  $d_n^\ell$  form a stationary sequence whose elements equip  $\mathcal{G}_n$  once again with full metric structure. We refer to this space as  $(\mathcal{G}_n, d_n^*)$ , “The metric space of connected simple graphs.” Unlike  $(\mathcal{G}_n, d_n^1)$ , the space  $(\mathcal{G}_n, d_n^*)$  is connected. A number of open questions concerning the geodesic structure of  $(\mathcal{G}_n, d_n^*)$  are also presented.

## 1 Introduction

We begin with an informal, brief description of the concepts and motivation behind this work. The statements here will be formulated precisely in the subsequent sections.

Given two connected, simple graphs  $H, K$  of order  $n$ , begin by considering a one-to-one mapping  $\phi : V[K] \rightarrow V[H]$ . Upon fixing  $\phi$ , to each edge  $e = (u, v) \in E[K]$  we associate a walk  $p_e$  between  $\phi(u)$  and  $\phi(v)$  in  $H$ . We view  $\phi$  together with the chosen set of walks  $Q = \{p_e | e \in E[K]\}$  as a topological embedding  $\phi'$  of  $K$  into  $H$ . If all the walks in  $Q$  are of length  $\leq \ell$ , then  $\phi'$  is called an  $\ell$ -topological

---

Subject Classification: 05C99 (Graph theory)

## 5 Acknowledgements

The authors gratefully acknowledge the Center for Computational Science at the Naval Research Laboratory, Washington DC, where this work began, in the context of virtual path layout for high-speed computer networks. The authors would like to thank Daniel Tunkelang for many helpful discussions and comments.

## References

- [1] G. Chartrand, G. Kubicki, and M. Schultz. Graph similarity and distance in graphs. *Aequationes Mathematicae*, 55:129–145, 1998.
- [2] M. Coornaert, T. Delzant, and A. Papdopoulos. *Geometrie et theorie des groupes: Les groupes hyperboliques de Gromov (Lecture Notes in Mathematics No. 1441)*. Springer Verlag, 1990.
- [3] R. Diestel. *Graph Theory*. Springer Verlag, 1997.
- [4] O. Gerstel, I. Cidon, and S. Zaks. The layout of virtual paths in ATM networks. *IEEE/ACM Transactions on Networking*, 4(6):873–884, 1996.
- [5] F. Harary. *Graph Theory*. Addison-Wesley, 1969.
- [6] S. Zaks. Path layout in ATM networks—a survey. *The DIMACS Workshop on Networks in Distributed Computing, DIMACS Center, Rutgers University*, October 1997.

Kiran R. Bhutani  
Department of Mathematics, The Catholic University of America.  
Washington D.C. 20064, U.S.A.

Bilal Khan  
Department of Mathematics, City University of New York Graduate Center,  
365 5th Avenue, New York City, NY 10016, U.S.A.